

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 January 2003 (16.01.2003)

PCT

(10) International Publication Number
WO 03/004621 A2

(51) International Patent Classification⁷: **C12N**

IMAI, Shin-Ichiro [JP/US]; 11019 Wellsley Court, St. Louis, MO 63146 (US). **GU, Wei** [US/US]; 75 Sidney Street, Cambridge, MA 02139-4169 (US).

(21) International Application Number: PCT/US02/21461

(22) International Filing Date: 8 July 2002 (08.07.2002)

(74) Agent: **MYERS, Louis**; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/303,370	6 July 2001 (06.07.2001)	US
60/303,456	6 July 2001 (06.07.2001)	US
Not furnished	5 July 2002 (05.07.2002)	US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:

US	Not furnished (CON)
Filed on	5 July 2002 (05.07.2002)

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **ELIXIR PHARMACEUTICALS, INC.** [US/US]; One Kendall Square, Building 1000-Fifth Floor, Cambridge, MA 02139 (US).

Published:

— *without international search report and to be republished upon receipt of that report*

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GUARENTE, Leonard** [US/US]; 72 Goodnough Road, Chestnut Hill, MA 02167 (US). **VAZIRI, Homayoun** [—/US]; 44 Clarendon Ave., Unit 2, Somerville, MA 02144 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SIR2 ACTIVITY

(57) Abstract: This invention relates to methods of screening compounds that modulate cellular and organismal processes by modification of the activity of SIR2 and/or transcription factors, e.g., p53, particularly methods of screening for compounds that modify lifespan and/or metabolism of a cell or an organism by modulation of the activity of SIR2 and/or transcription factors, e.g., p53, and more particularly to methods of screening for compounds that modulate the activity of Sir2 and/or transcription factors, e.g., p53. In particular, the present invention relates to a method for screening a compound, by providing a test mixture comprising a transcription factor, Sir2, and a Sir2 cofactor with the compound, and evaluating an activity of a component of the test mixture in the presence of the compound. The invention further relates to therapeutic uses of said compounds. The invention further relates to a method of modifying the acetylation status of a transcription factor binding site on histone or DNA by raising local concentrations of Sir2.



WO 03/004621 A2

SIR2 ACTIVITY

CLAIM OF PRIORITY

This application claims priority under 35 USC §119(e) to U.S. Patent Application Serial No. 60/303,370, filed on July 6, 2001, and U.S. Patent Application Serial No. 60/303,456, also filed on July 6, 2001, the entire contents of which are hereby incorporated by reference.

GOVERNMENT SUPPORT

The invention was supported, in whole or in part, by a grant RO1 CA78461 to RAW; NHLBI/NIH Fellowship to SKD KO8 HL04463. The U.S. Government has certain rights in the invention.

BACKGROUND

Regulation of the cell cycle is important in homeostasis of both cells and organisms (*e.g.*, mammalian cells or mammals). Disruptions in the normal regulation of the cell cycle can occur, for example, in tumors which proliferate uncontrollably, in response to DNA damage (*e.g.*, ionizing radiation) to the cell or organism, and under conditions of stress (*e.g.*, oxidative stress) in the cell or organism.

The p53 tumor suppressor protein exerts anti-proliferative effects, including growth arrest, apoptosis, and cell senescence, in response to various types of stress, *e.g.*, DNA damage (Levine, 1997; Giaccia and Kastan, 1998; Prives and Hall, 1999; Oren, 1999; Vogelstein *et al.*, 2000). Inactivation of p53 function appears to be critical to tumorigenesis (Hollstein *et al.*, 1999). Mutations in the p53 gene have been shown in more than half of all human tumors (Hollstein *et al.*, 1994). Accumulating evidence further indicates that, in the cells that retain wild-type p53, other defects in the p53 pathway also play an important role in tumorigenesis (Prives and Hall, 1999; Lohrum and Vousden, 1999; Vousden, 2000). The molecular function of p53 that is required for tumor suppression involves its ability to act as a transcriptional factor in regulating endogenous gene expression. A number of genes which are critically involved in either cell growth arrest or apoptosis have been identified as p53 direct targets, including p21CIP1/WAF1, Mdm2, GADD45, Cyclin G, 14-3-3F, Noxa, p53AIP1, PUMA and others

(Nakano and Vousden, 2001; Yu *et al.*, 2001; Oda *et al.*, 2000a, 2000b; El-Deriry *et al.*, 1993; Wu *et al.*, 1993; Barak *et al.*, 1993; Kastan *et al.*, 1992; Okamoto and Beach, 1994).

p53 is a short-lived protein whose activity is maintained at low levels in normal cells. Tight regulation of p53 is essential for its effect on tumorigenesis as well as maintaining normal cell growth. The precise mechanism by which p53 is activated by cellular stress is not completely understood. It is generally thought to involve primarily post-translational modifications of p53, including phosphorylation and acetylation (reviewed in Appella and Anderson, 2000; Giaccia and Kastan, 1998). Early studies demonstrated that CBP/p300, a histone acetyl-transferase (HAT), acts as a coactivator of p53 and potentiates its transcriptional activity as well as biological function *in vivo* (Gu *et al.*, 1997; Lill *et al.*, 1997; Avantaggiati *et al.*, 1997). Genetic studies have also revealed that p300 mutations are present in several types of tumors, and that mutations of CBP in human Rubinstein-Taybi syndrome as well as CBP knockout mice lead to higher risk of tumorigenesis, further supporting an important role for this interaction in the tumor suppressor pathway (reviewed in Goodman and Smolik, 2000; Gile *et al.*, 1998; Kung *et al.*, 2000; Gayther *et al.*, 2000). Significantly, the observation of functional synergism between p53 and CBP/p300 together with its intrinsic HAT activity led to the discovery of a novel FAT (Transcriptional factor acetyl-transferase) activity of CBP/p300 on p53 which suggests that acetylation represents a general functional modification for non-histone proteins *in vivo* (Gu and Roeder, 1997) which has been shown for other transcriptional factors (reviewed in Kouzarides, 2000; Sterner and Berger, 2000; Muth *et al.*, 2001).

p53 is specifically acetylated at multiple lysine residues (Lys 370, 371, 372, 381, 382) of the C-terminal regulatory domain by CBP/p300. The acetylation of p53 can dramatically stimulate its sequence-specific DNA binding activity, perhaps as a result of an acetylation-induced conformational change (Gu and Roeder, 1997; Sakaguchi *et al.*, 1998; Liu *et al.*, 1999). By developing site-specific acetylated p53 antibodies, CBP/p300 mediated acetylation of p53 was confirmed *in vivo* by a number of studies (reviewed in Chao *et al.*, 2000; Ito *et al.*, 2001). In addition, p53 can be acetylated at Lys320 by another HAT cofactor, PCAF, although the *in vivo* functional consequence needs to be further elucidated (Sakaguchi *et al.*, 1998; Liu *et al.*, 1999; Liu *et al.*, 2000). Steady-state levels of acetylated p53 are stimulated in response to various types of stress (reviewed in Ito *et al.*, 2001).

Recently, by introducing a transcription defective p53 mutant (p53Q25S26) into mice, it was found that the mutant mouse thymocytes and ES cells failed to undergo DNA damage-induced apoptosis (Chao *et al.*, 2000; Jimenez *et al.*, 2000). Interestingly, this mutant protein was phosphorylated normally at the N-terminus in response to DNA damage but could not be acetylated at the C-terminus (Chao *et al.*, 2000), supporting a critical role of p53 acetylation in transactivation as well as p53-dependent apoptotic response (Chao *et al.*, 2000; Luo *et al.*, 2000). Furthermore, it has been found that oncogenic Ras and PML upregulate acetylated p53 in normal primary fibroblasts, and induce premature senescence in a p53-dependent manner (Pearson *et al.*, 2000; Ferbeyre *et al.*, 2000). Additionally acetylation, not phosphorylation of the p53 C-terminus, may be required to induce metaphase chromosome fragility in the cell (Yu *et al.*, 2000). Thus, CBP/p300-dependent acetylation of p53 may be a critical event in p53-mediated transcriptional activation, apoptosis, senescence, and chromosome fragility.

In contrast, much less is known about the role of deacetylation in modulating p53 function. Under normal conditions, the proportion of acetylated p53 in cells remains low. This may reflect the action of strong deacetylase activities *in vivo*. The acetylation level of p53 is enhanced when the cells are treated with histone deacetylase (HDAC) inhibitors such as Trichostatin A (TSA). These observations led to identification of a HDAC1 complex which is directly involved in p53 deacetylation and functional regulation (Luo *et al.*, 2000; Juan *et al.*, 2000). PID/MTA2, a component of the HDAC1 complex, acts as an adaptor protein to enhance HDAC1-mediated deacetylation of p53 which is repressed by TSA (Luo *et al.*, 2000). In addition, Mdm2, a negative regulator of p53, actively suppresses CBP/p300-mediated p53 acetylation, and this inhibitory effect can be abrogated by tumor suppressor p19ARF. Acetylation may have a critical role in the p53-MDM2-p19ARF feed back loop (Ito *et al.*, 2001; Kobet *et al.*, 2000).

The Silent Information Regulator (SIR) family of genes represents a highly conserved group of genes present in the genomes of organisms ranging from archaeobacteria to a variety of eukaryotes (Frye, 2000). The encoded SIR proteins are involved in diverse processes from regulation of gene silencing to DNA repair. The proteins encoded by members of the SIR2 gene family show high sequence conservation in a 250 amino acid core domain. A well-characterized gene in this family is *S. cerevisiae* SIR2, which is involved in silencing HM loci that contain information specifying yeast mating type, telomere position effects and cell aging (Guarente,

1999; Kaeberlein *et al.*, 1999; Shore, 2000). The yeast Sir2 protein belongs to a family of histone deacetylases (reviewed in Guarente, 2000; Shore, 2000). The Sir2 homolog, CobB, in *Salmonella typhimurium*, functions as an NAD (nicotinamide adenine dinucleotide)-dependent ADP-ribosyl transferase (Tsang and Escalante-Semerena, 1998).

The Sir2 protein is a deacetylase which uses NAD as a cofactor (Imai *et al.*, 2000; Moazed, 2001; Smith *et al.*, 2000; Tanner *et al.*, 2000; Tanny and Moazed, 2001). Unlike other deacetylases, many of which are involved in gene silencing, Sir2 is insensitive to histone deacetylase inhibitors like trichostatin A (TSA) (Imai *et al.*, 2000; Landry *et al.*, 2000a; Smith *et al.*, 2000).

Deacetylation of acetyl-lysine by Sir2 is tightly coupled to NAD hydrolysis, producing nicotinamide and a novel acetyl-ADP ribose compound (1-O-acetyl-ADP-ribose) (Tanner *et al.*, 2000; Landry *et al.*, 2000b; Tanny and Moazed, 2001). The NAD-dependent deacetylase activity of Sir2 is essential for its functions which can connect its biological role with cellular metabolism in yeast (Guarente, 2000; Imai *et al.*, 2000; Lin *et al.*, 2000; Smith *et al.*, 2000). Mammalian Sir2 homologs have NAD-dependent histone deacetylase activity (Imai *et al.*, 2000; Smith *et al.*, 2000). Most information about Sir2 mediated functions comes from the studies in yeast (Gartenberg, 2000; Gottschling, 2000).

Among Sir2 and its homolog proteins (HSTs) in yeast, Sir2 is the only protein localized in nuclei, which is critical for both gene silencing and extension of yeast life-span (reviewed in Guarente, 2000). Based on protein sequence homology analysis, mouse Sir2 α and its human ortholog SIRT1 (or human Sir2 α or hSir2) are the closest homologs to yeast Sir2 (Imai *et al.*, 2000; Frye, 1999, 2000) and both exhibit nuclear localization (Figure 7C). Homologues of Sir2 have been identified in almost all organisms examined including bacteria, which has no histone proteins (reviewed in Gray and Ekstrom, 2001; Frye, 1999; 2000; Brachmann *et al.*, 1995). For this reason it is likely that Sir2 also targets non-histone proteins for functional regulation (Muth *et al.*, 2001).

The *S. cerevisiae* Sir2 is involved in DNA damage responses (Martin *et al.*, 1999; McAinsh *et al.*, 1999; Mills *et al.*, 1999). In mammalian cells, one of the primary mediators of the DNA damage response is the p53 protein (Levine, 1997; Oren, 1999; Vogelstein *et al.*, 2000). Following DNA damage, the p53 protein is protected from rapid degradation and acquires transcription-activating functions, these changes being achieved largely through post-

translational modifications (Abraham *et al.*, 2000; Canman *et al.*, 1998; Chehab *et al.*, 1999; Sakaguchi *et al.*, 1998; Shieh *et al.*, 2000; Siliciano *et al.*, 1997). Transcriptional activation of p53 protein in turn upregulates promoters of a number of genes including p21WAF1 (el-Deiry *et al.*, 1993) that promotes cell cycle exit or death-inducing proteins like PIDD (Lin *et al.*, 2000).

The p53 protein is phosphorylated in response to DNA damage (Siliciano *et al.*, 1997). There are at least 13 different residues both at the N and C terminal portions of p53 protein that are phosphorylated by various kinases (Appella and Anderson, 2000). For example, the ATM and ATR proteins phosphorylate p53 at residue Ser15 (Khanna *et al.*, 1998; Siliciano *et al.*, 1997; Tibbetts *et al.*, 1999) and Chk1/2 kinases at residue Ser20 (Chehab *et al.*, 1999; Shieh *et al.*, 2000).

Modification of Ser15 is important for the functional activation of the p53 protein. Phosphorylation of Ser15 may increase the affinity of the p300 acetylase for p53 (Dumaz and Meek, 1999; Lambert *et al.*, 1998).

p53 is acetylated *in vitro* by p300 at Lys 370-372, 381 and 382 (Gu and Roeder, 1997). In response to DNA damage, p53 is also acetylated *in vivo* at Lys 373 and Lys 382 (Abraham *et al.*, 2000; Sakaguchi *et al.*, 1998). Other factors that can affect acetylation of p53 include MDM2 protein, which is involved in the negative regulation of p53 (Oren, 1999) and can suppress acetylation of p53 protein by p300 (Ito *et al.*, 2001; Kobet *et al.*, 2000). While acetylation by p300 and deacetylation by the TSA-sensitive HDAC1 complex (Luo *et al.*, 2000) have been shown to be important in regulation of p53 protein activity, the remaining factors responsible for its regulation as a transcription factor remain elusive.

Analogues of NAD that inhibit endogenous ADP-ribosylases reduce induction of p21WAF1 in response to DNA damage and overcome p53-dependent senescence (Vaziri *et al.*, 1997). In addition, p53 protein can bind to the NAD-dependent poly-ADP-ribose polymerase.

The SIR complex in *Saccharomyces cerevisiae* was originally identified through its involvement in the maintenance of chromatin silencing at telomeres and at mating type loci. It is composed of four components, Sir1p, Sir2p, Sir3p, and Sir4p, that normally reside at yeast telomeres. In response to DNA damage, the SIR complexes relocate to the site of double-stranded breaks where they participate in the repair of the lesions by non-homologous end joining. This DNA damage response is dependent on the function of the MEC1/RAD9 DNA checkpoint pathway. MEC1 is a homolog of the ATM protein that coordinates the DNA damage

response in mammalian cells, in part by triggering the cascade of events that lead to the stabilization of the p53 protein (Canman *et al.*, 1998). Another major function of Sir2, gene silencing, is closely tied to the regulation of lifespan in *S. cerevisiae* (Guarente, 1999).

Double-strand breaks in the genome of mammals invoke a cascade of signaling events that ultimately cause phosphorylation and subsequent stabilization of p53 protein. In addition, these strand breaks lead to activation of p53 protein as a transcription factor. This activation may be due largely to its acetylation (Gu and Roeder, 1997; Sakaguchi *et al.*, 1998). The resulting stabilized, activated p53 protein contributes to the upregulation of cyclin-dependent kinase inhibitors such as p21 WAF1 and hence to the cytostatic effects of p53. Alternatively, depending on the cellular background or degree of damage, the apoptotic effects of p53 may predominate through its ability to induce expression of pro-apoptotic proteins such as PIDD (Lin *et al.*, 2000). These various phenomena indicate that specific components of the machinery that monitors the integrity of the genome are clearly able to alert p53 to the presence of genetic damage, leading to its functional activation. Conversely, in the event that damage has been successfully repaired, signals must be conveyed to p53 in order to deactivate it. Thus, a cell cycle advance that has been halted by p53 to enable repair to proceed should be relieved following completion of repair, enabling the cell to return to its active growth state. For this reason, the inactivation of p53 becomes as important physiologically as its activation.

In light of this information, modulators of Sir2 and/or p53 activity would be useful in modulating various cellular processes including, *e.g.*, repair of DNA damage, apoptosis, oncogenesis, gene silencing and senescence, *inter alia*.

SUMMARY

In one aspect, the present invention relates to methods and compositions employing p53 and Sir2 proteins. Cellular and organismal processes are regulated by modulating the activity of Sir2 and/or p53. In some cases the regulated processes control a program of regulated aging and/or metabolism of a cell or an organism. Compounds that regulate the activity of Sir2 and/or p53 can be identified, for example, by a method described herein.

As used herein, the term "Sir2" refers to a protein that is at least 25% identical to the 250 amino acid conserved Sir2 core catalytic domain, amino acids 258-451 of SEQ ID NO. 12. A

Sir2 protein can be for example, at least 30, 40, 50, 60, 70, 80, 85, 90, 95, 99% identical to amino acids 258-451 of SEQ ID NO. 12. For example, the Sir2 protein is human SIRT1, GenBank Accession No: AF083106. There are at least seven different Sir2 homologs present in mammalian cells (Frye, 1999, 2000; Imai *et al.*, 2000; Gray and Ekstrom, 2001). The mouse Sir2 α and human SIRT1, are preferred Sir2 proteins.

Sir2 can be a protein (*e.g.*, SEQ ID NOS. 8, 10, 12, 14, 16 or 18) or a fragment of the protein capable of deacetylating a substrate in the presence of NAD and/or an NAD analog and/or a fragment capable of binding to a target protein, *e.g.*, a transcription factor. Such functions can be evaluated by a method described herein. A Sir2 fragment can include a “domain” which is a structurally stable folded unit of the full-length protein. The Sir2 protein can be encoded by the nucleic acid sequence of SEQ ID NOS. 7, 9, 11, 13, 15 or 17. In a preferred embodiment, the Sir2 is a human Sir2. A model of the three-dimensional structure of a Sir2 protein has been determined (see, *e.g.*, Bedalov *et al.* (2001), Min *et al.* (2001), Finnin *et al.*, (2001)) and provides guidance for identifying domains of Sir2.

A “full length” Sir2 protein refers to a protein that has at least the length of a naturally-occurring Sir2 protein. A “full length” Sir2 protein or a fragment thereof can also include other sequences, *e.g.*, a purification tag, or other attached compounds, *e.g.*, an attached fluorophore, or cofactor.

The invention includes sequences and variants that include one or more substitutions, *e.g.*, between one and six substitutions, *e.g.*, with respect to a naturally-occurring protein. Whether or not a particular substitution will be tolerated can be determined by a method described herein. One or more or all substitutions may be conservative. A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine).

The terms “identical” or percent “identity,” in the context of two or more nucleic acids or

polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (i.e., about 50% identity, preferably 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or higher identity over a specified region (e.g., the *C. elegans* proteins provided herein), when compared and aligned for maximum correspondence over a comparison window or designated region) as measured using a sequence comparison methodology such as BLAST or BLAST 2.0 with default parameters described below, or by manual alignment and visual inspection. Such sequences are then said to be “substantially identical.” This definition also refers to, or may be applied to, the complement of a test nucleic acid sequence. The definition also includes sequences that have deletions and/or additions, as well as those that have substitutions. As described below, the preferred algorithms can account for gaps and the like. Preferably, identity exists over a region that is at least about 25 amino acids or nucleotides in length, or more preferably over a region that is at least 50 or 100 amino acids or nucleotides in length.

The p53 polypeptide can have greater than or equal to 25%, 50%, 75%, 80%, 90% overall identity or greater than or equal to 30%, 50%, 75%, 80%, 90% overall similarity to SEQ ID NO. 3. Preferably, the Sir2 or p53 polypeptide is a human protein (e.g., as described herein), although it may also be desirable to analyze Sir2 or p53 polypeptides isolated from other organisms such as yeast, worms, flies, fish, reptiles, birds, mammals (especially rodents), and primates using the methods of the invention.

In one aspect, the invention features a method of screening a compound. The method includes providing a reaction mixture including Sir2, a transcription factor, and the compound, and determining if the compound modulates Sir2 interaction with, e.g., binding, of the transcription factor. Determining if the compound modulates Sir2 binding may be accomplished by methods known in the art, including comparing the binding of Sir2 to the transcription factor at a first concentration of the compound and at a second concentration of the compound. In a further embodiment, either of the first or second concentration of the compound may be zero, e.g., as a reference or control.

In a further embodiment, the reaction mixture also includes a Sir2 cofactor, such as NAD or an NAD analog.

In a further embodiment, the transcription factor is p53 or a Sir-2 binding fragment thereof. The transcription factor, e.g., p53, or fragment thereof may be acetylated or labeled. In a preferred embodiment, the transcription factor is an acetylated p53 fragment, and the fragment includes lysine 382.

In a further embodiment, the Sir2 included in the reaction mixture is a Sir2 variant, e.g., a variant that has reduced deacetylase activity, such as the H363Y mutation. The Sir2 may be human, e.g., human SIRT1. Alternatively, the Sir2 may be murine, e.g., Sir2 α . In one embodiment of the inventions, the Sir2 is exogenous and expressed from a heterologous nucleic acid. Additionally, in a further embodiment, the transcription factor may be exogenous and expressed from a heterologous nucleic acid.

The method of screening can be used to identify compounds that modulate, e.g., increase or decrease, cell growth, modulate, e.g., slow or speed, aging, modulate, e.g., increase or decrease, lifespan, modulate cellular metabolism, e.g., by increasing or decreasing a metabolic function or rate.

In another aspect, the invention features a method of screening a compound by providing a reaction mixture comprising Sir2, a transcription factor, and the compound, and determining if the compound modulates Sir2-mediated deacetylation of the transcription factor. The step of determining if the compound modulates Sir2-mediated deacetylation of the transcription factor may be performed by methods known in the art, including comparing the binding of Sir2 to the transcription factor at a first concentration of the compound and at a second concentration of the compound. In a further embodiment, either of the first or second concentration of the compound may be zero, e.g., as a reference or control. In a further embodiment, the reaction mixture also includes a Sir2 cofactor, such as NAD or an NAD analog.

In a further embodiment, the transcription factor is p53 or a Sir-2 binding fragment thereof. The p53 or fragment thereof may be acetylated or labeled. In a preferred embodiment, the transcription factor is an acetylated p53 fragment, and the fragment includes lysine 382.

In a further embodiment, the Sir2 included in the reaction mixture is a Sir2 variant that has reduced deacetylase activity, such as the H363Y mutation. The Sir2 may be human, e.g., human SIRT1. Alternatively, the Sir2 may be murine, e.g., Sir2a. In one embodiment of the inventions, the Sir2 is exogenous and expressed from a heterologous nucleic acid. Additionally,

in a further embodiment, the transcription factor may be exogenous and expressed from a heterologous nucleic acid.

The method of screening can be used to identify compounds that modulate, e.g., increase or decrease, cell growth, modulate, e.g., slow or speed, aging, modulate, e.g., increase or decrease, lifespan, modulate cellular metabolism, e.g., by increasing or decreasing a metabolic function or rate.

The present invention also relates to a method of screening a compound by providing a compound that interacts with Sir2, e.g., a compound that binds Sir2; contacting the compound with a cell or a system; and determining if the compound modulates transcription of a p53-regulated gene. Determining if the compound modulates transcription of a p53-regulated gene may be by any of the methods known in the art, including comparing the modulation of transcription of a p53-regulated gene at a first concentration of the compound and at a second concentration of the compound. In a further embodiment, either of the first or second concentration of the compound may be zero, e.g., as a reference or control.

In a related aspect, the invention features a method of evaluating a compound, the method comprising: contacting Sir2 or a transcription factor, e.g., p53, with a test compound; evaluating an interaction between the test compound and the Sir2 or the transcription factor, e.g., p53; contacting a cell or organism that produces the Sir2 or transcription factor polypeptide with the test compound; and evaluating the effect of the test compound on the rate of aging on the cell or organism. The interaction can, for example, be a physical interaction, e.g., a direct binding interaction, a covalent change in one or both of the test compound or the Sir2 or transcription factor, a change in location of the test compound (e.g., a change in subcellular localization), or a functional interaction (e.g., an alteration in activity, stability, structure, or activity of the polypeptide).

In some embodiments, the method is repeated one or more times such that, e.g., a library of test compounds can be evaluated. In an related embodiment, the evaluating of the interaction with the test compound and the Sir2 or the transcription factor, e.g., p53, is repeated, and the evaluating of the rate of aging is selectively used for compounds for which an interaction is

detected. Possible test compounds include, e.g., small organic molecules, peptides, antibodies, and nucleic acid molecules.

In some embodiments, the interaction between the test compound and the Sir2 or transcription factor, e.g., p53, is evaluated in vitro, e.g., using an isolated polypeptide. The Sir2 or transcription factor, e.g., p53, polypeptide can be in solution (e.g., in a micelle) or bound to a solid support, e.g., a column, agarose beads, a plastic well or dish, or a chip (e.g., a microarray). Similarly, the test compound can be in solution or bound to a solid support.

In other embodiments, the interaction between the test compound and the Sir2 or transcription factor, e.g., p53, is evaluated using a cell-based assay. For example, the cell can be a yeast cell, an invertebrate cell (e.g., a fly cell), or a vertebrate cell (e.g., a *Xenopus* oocyte or a mammalian cell, e.g., a mouse or human cell). In preferred embodiments, the cell-based assay measures the activity of the Sir2 or transcription factor, e.g., p53, polypeptide.

In preferred embodiments, the effect of the test compound on the rate of aging of a cell or animal is evaluated only if an interaction between the test compound and the Sir2 or transcription factor, e.g., p53, is observed.

In some embodiments, the cell is a transgenic cell, e.g., a cell having a transgene. In some embodiments, the transgene encodes a protein that is normally exogenous to the transgenic cell. In some embodiments, the transgene encodes a human protein, e.g., a human Sir2 or transcription factor, e.g., p53, polypeptide. In some embodiments, the transgene is linked to a heterologous promoter. In other embodiments, the transgene is linked to its native promoter. In some embodiments, the cell is isolated from an organism that has been contacted with the test compound. In other embodiments, the cell is contacted directly with the test compound.

In other embodiments, the rate of aging of an organism, e.g., an invertebrate (e.g., a worm or a fly) or a vertebrate (e.g., a rodent, e.g., a mouse) is determined. The rate of aging of an organism can be determined by a variety of methods, e.g., by one or more of: a) assessing the life span of the cell or the organism; (b) assessing the presence or abundance of a gene transcript or gene product in the cell or organism that has a biological age-dependent expression pattern; (c) evaluating resistance of the cell or organism to stress, e.g., genotoxic stress (e.g., etoposide, UV irradiation, exposure to a mutagen, and so forth) or oxidative stress; (d) evaluating one or more metabolic parameters of the cell or organism; (e) evaluating the proliferative capacity of the cell or a set of cells present in the organism; (f) evaluating physical appearance or behavior

of the cell or organism, and (g) assessing the presence or absence of a gene transcript or gene product in the cell or organism that has a p53-regulation-dependent expression pattern. In one example, evaluating the rate of aging includes directly measuring the average life span of a group of animals (e.g., a group of genetically matched animals) and comparing the resulting average to the average life span of a control group of animals (e.g., a group of animals that did not receive the test compound but are genetically matched to the group of animals that did receive the test compound). Alternatively, the rate of aging of an organism can be determined by measuring an age-related parameter. Examples of age-related parameters include: appearance, e.g., visible signs of age; the expression of one or more genes or proteins (e.g., genes or proteins that have an age-related expression pattern); resistance to oxidative stress; metabolic parameters (e.g., protein synthesis or degradation, ubiquinone biosynthesis, cholesterol biosynthesis, ATP levels, glucose metabolism, nucleic acid metabolism, ribosomal translation rates, etc.); and cellular proliferation (e.g., of retinal cells, bone cells, white blood cells, etc.). In some embodiments, the organism is a transgenic animal. The transgenic animal can include a transgene that encodes, e.g., a copy of a Sir2 or transcription factor protein, e.g., a p53 protein, e.g., the Sir2 or transcription factor, e.g., a p53 polypeptide that was evaluated for an interaction with the test compound. In some embodiments, the transgene encodes a protein that is normally exogenous to the transgenic animal. For example, the transgene can encode a human protein, e.g., a human Sir2 or transcription factor, e.g., p53, polypeptide. In some embodiments, the transgene is linked to a heterologous promoter. In other embodiments, the transgene is linked to its native promoter. In some embodiments, the transgenic animal further comprises a genetic alteration, e.g., a point mutation, insertion, or deficiency, in a gene encoding an endogenous Sir2 or transcription factor, e.g., p53, protein, such that the expression or activity of the endogenous Sir2 or transcription factor protein is reduced or eliminated.

In some embodiments, the organism is on a calorically rich diet, while in other embodiments the organism is on a calorically restricted diet.

In some embodiments, a portion of the organism's life, e.g., at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more, of the expected life span of the organism, has elapsed prior to the organism being contacted with the test compound.

In another aspect, the invention features a method of evaluating a protein, comprising: identifying or selecting a candidate protein, wherein the candidate protein is a Sir2 or transcription factor, e.g. p53, polypeptide; altering the sequence, expression or activity of the candidate protein in a cell or in one or more cells of an organism; and determining whether the alteration has an effect on the interaction, e.g., binding, of Sir2 with a transcription factor, e.g. p53, or on the deacetylation of transcription factor, e.g. p53.

In some embodiments, the candidate protein is identified by amplification of the gene or a portion thereof encoding the candidate protein, e.g., using a method described herein, e.g., PCR amplification or the screening of a nucleic acid library. In preferred embodiments, the candidate protein is identified by searching a database, e.g., searching a sequence database for protein sequences homologous to Sir2 or a transcription factor, e.g., p53.

In preferred embodiments, the candidate protein is a human protein. In other embodiments, the candidate protein is a mammalian protein, e.g., a mouse protein. In other embodiments, the protein is a vertebrate protein, e.g., a fish, bird or reptile protein, or an invertebrate protein, e.g., a worm or insect protein. In still other embodiments, the protein is a eukaryotic protein, e.g., yeast protein.

In another aspect, the invention features method of evaluating a protein, the method comprising a) identifying or selecting a candidate protein, wherein the candidate protein is Sir2 or a transcription factor, e.g., p53; b) identifying one or more polymorphisms in a gene, e.g., one or more SNPs that encodes the candidate protein; and c) assessing correspondence between the presence of one or more of the polymorphisms and an interaction, e.g., binding, of Sir2 with the transcription factor, e.g., p53, or with the deacetylation of the transcription factor, e.g., p53. The polymorphisms can be naturally occurring or laboratory induced. In one embodiment, the organism is an invertebrate, e.g., a fly or nematode; in another embodiment the organism is a mammal, e.g., a rodent or human. A variety of statistical and genetic methods can be used to assess correspondence between a polymorphism and longevity. Such correlative methods include determination of linkage disequilibrium, LOD scores, and the like.

In another aspect, the invention features a method of modulating cell growth in an animal, e.g., a mammal, by modulating the Sir2-mediated deacetylation of a transcription factor in the animal.

In one embodiment, the method includes modulating cell growth by increasing acetylation of p53. In a further embodiment, the method includes inactivating Sir2, e.g., by the use of antisense, RNAi, antibodies, intrabodies, NAD depletion, a dominant negative mutant of Sir2, or by the addition of Sir2 cofactor-analogs, e.g., NAD analogs such as those described in Vaziri et al. (1997) or nicotinamide. In a further embodiment, the method includes introducing a deacetylation-resistant form of p53. In still another embodiment, the invention is a method for treating a mammal, e.g., a mammal having a disease characterized by unwanted cell proliferation, e.g., cancer, accelerated senescence-related disorders, inflammatory and autoimmune disorders, Alzheimer's disease, and aging-related disorders.

In another embodiment, the method includes modulating cell growth by decreasing acetylation of p53. In a further embodiment, the method includes increasing NAD concentrations. In a further embodiment, the method includes increasing Sir2 concentrations, e.g. by addition of purified Sir2, by expression of Sir2 from heterologous genes, or by increasing the expression of endogenous Sir2, or by the addition of Sir2 cofactor-analogs, e.g., NAD analogs such as those described in Vaziri et al. (1997).

The present invention also relates to a method of modulating the growth of a cell *in vivo* or *in vitro* by modulating the Sir2-mediated deacetylation of a transcription factor in the cell.

In one embodiment, the method includes modulating the growth of a cell by increasing acetylation of p53, thereby decreasing cell growth. In a further embodiment, the method includes inactivating Sir2, e.g., by the use of antisense, RNAi, antibodies, intrabodies, NAD depletion, a dominant negative mutant of Sir2, or nicotinamide, or decreasing Sir2 activity by the addition of Sir2 cofactor-analogs, e.g., NAD analogs such as those described in Vaziri et al. (1997). In a further embodiment, the method includes introducing a deacetylation-resistant form of p53.

In one embodiment, the method includes modulating the growth of a cell by decreasing acetylation of p53, thereby increasing cell growth. In a further embodiment, the method includes increasing NAD concentrations. In a further embodiment, the method includes increasing Sir2

concentrations, e.g. by addition of purified Sir2, by expression of Sir2 from heterologous genes, or by increasing the expression of endogenous Sir2, or by the addition of Sir2 cofactor-analogs, e.g., NAD analogs such as those described in Vaziri et al. (1997).

In one aspect the invention features a method of directing Sir2 to a transcription factor binding site, e.g., a p53 binding site, and thereby modifying the acetylation status of the binding site on histone or DNA. The method includes providing a Sir2-transcription factor complex under conditions such that the transcription factor targets Sir2 to the transcription factor binding site, allowing the Sir 2 to modify the acetylation status of histones and DNA at the transcription factor binding site.

In a preferred embodiment, the method is performed *in vivo* or *in vitro*, e.g., in an animal or in a cell.

In a preferred embodiment, the Sir2-transcription factor complex is provided at a different stage of development of the cell or animal or at a greater concentration than occurs naturally.

In a preferred embodiment, the Sir2 or transcription factor or both is increased, e.g., by supplying exogenous Sir2 and/or transcription factor, e.g., p53, by supplying an exogenous nucleic acid encoding Sir2 or transcription factor, e.g., p53, or by inducing endogenous production of Sir2 or a transcription factor, e.g., p53.

In one embodiment, the present invention relates to a method of evaluating a compound, e.g., a potential modulator of Sir2 or transcription factor, e.g., p53 activity, comprising the steps of contacting the transcription factor, e.g., p53, Sir2, and NAD or an NAD analog with the compound; evaluating an interaction between the compound and one or more of the transcription factor, e.g., p53, Sir2, and a cofactor such as NAD or an NAD analog; contacting the compound with a cell or organism having transcription factor, e.g., p53 or Sir2 activity; and evaluating the rate of aging of the cell or organism. In a preferred embodiment, evaluating the rate of aging comprises one or more of:

- a) assessing the life span of the cell or organism;
- b) assessing the presence or absence of a gene transcript or gene product in the cell or organism that has a biological age-dependent expression pattern;

- c) evaluating resistance of the cell or organism to stress;
- d) evaluating one or more metabolic parameters of the cell or organism;
- e) evaluating the proliferative capacity of the cell or a set of cells present in the organism;
- f) evaluating physical appearance, behavior, or other characteristic of the cell or organism; and
- g) assessing the presence or absence of a gene transcript or gene product in the cell or organism that has a p53-regulation-dependent expression pattern.

The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

DESCRIPTION OF DRAWINGS

Figure 1. Interactions between p53 and mammalian Sir2 α both *in vitro* and *in vivo*.

(A) is an autoradiograph demonstrating direct interactions of Sir2 α with GST-p53. The GST-p53 full length protein (GST-p53) (lane 1), the N-terminus of p53 protein (1-73) (lane 2), the middle part of p53 (100-290) (lane 3), the C-terminus of p53 (290-393) (lane 4), and GST alone (lane 6) were used in GST pull-down assay with *in vitro* translated ³⁵S-labeled full length mouse Sir2 α . (B) is two western blots demonstrating p53 interactions with Sir2 α in H1299 cells. Western blot analyses of the indicated whole cell extract (WCE) (lanes 1, 3, 5, 7), or the p53 immunoprecipitates with M2 antibody (IP/Flag-p53) prepared from the transfected H1299 cells (lane 6, 8), or the Sir2 α immunoprecipitates (IP/Flag-Sir2 α) with M2 antibody prepared from the transfected H1299 cells (lanes 2, 4) with either anti-p53 monoclonal antibody (DO-1) (lanes 1-4), or anti-Sir2 α polyclonal antibody (lanes 5-8). The cells were either transfected with p53 (lanes 3, 4) or Sir2 α (lanes 7, 8) alone, or cotransfected with p53 and Sir2 α (lanes 1, 2, 5, 6). (C) is a schematic representation of the high homology regions between mouse Sir2 α and human SIRT1 (hSIRT1). The core domain represents the very conserved enzymatic domain among all Sir2 family proteins (Frye, 1999, 2000). (D) is a western blot demonstrating p53 interactions with human SIRT1 in H1299 cells. Western blot analyses of the indicated whole cell extract (WCE) (lanes 1, 3) or the Flag-hSIRT1 immunoprecipitates with M2 antibody (IP/hSIRT1)

(lanes 2, 4) prepared from either the hSIRT1 and p53 cotransfected H1299 cells (lanes 1, 2) or the p53 alone transfected cells (lanes 3, 4) with anti-p53 monoclonal antibody (DO-1).

Figure 2. P53 interacts with mammalian Sir2 α (mouse Sir2 α and hSIRT1) in normal cells.

(A) is two western blots demonstrating the interaction between p53 and hSIRT1 in H460 cells. (B) is two western blots demonstrating the interaction between p53 and Sir2 α in F9 cells. (C) The interaction between p53 and hSIRT1 in HCT116 cells either at the normal condition (lanes, 1-3) or after DNA damage treatment by etoposide (lanes, 4-6). Western blot analyses of the indicated whole cell extract (WCE) (lanes 1, 4), or immunoprecipitates with anti-Sir2 α antibody (IP/anti-Sir2 α) (lanes 2, 5) prepared from different cell extracts, or control immunoprecipitates with pre-immunoserum from the same extracts (lanes 3, 6), with anti-p53 monoclonal antibodies (DO-1 for human p53, 421 for mouse p53), or anti-Sir2 α antibody.

Figure 3. TSA-insensitive deacetylation of p53 by mammalian Sir2 α .

(A) Colloidal blue staining of a SDS-PAGE gel containing protein Marker (lane 1), a control eluate from M2 loaded with untransfected cell extract (lane 2), and 100 ng of the highly purified Flag-tagged Sir2 α recombinant protein (lane 3). (B) Deacetylation of p53 by Sir2 α , 2.5 μ g of ¹⁴C-labeled acetylated p53 (lane 1) was incubated with either the control eluate (lane 4), the purified 10 ng of Sir2 α (lanes 2 and 3), or the same amount of Sir2 α in the presence of 500 nM TSA (lane 5) for 60 min at 30°C. NAD (50 μ M) was also added in each reaction except lane 2. The proteins were analyzed by resolution on SDS-PAGE and autoradiography (upper) or Coomassie blue staining (lower). (C). Reduction of the steady-state levels of acetylated p53 by both mouse Sir2 α and human SIRT1 expression. Western blot analysis of H1299 cell extracts from the cells cotransfected with p53 and p300 (lane 1), or in combination with Sir2 α (lane 2), or in combination with hSIRT1 (lane 4), or in combination with Sir2 α H355A (lane 3), in combination with hSIRT5 (lane 5), or in combination with PARP (lane 6) by acetylated p53-specific antibody (upper) or DO-1 for total p53 (lower). (D) Deacetylation of p53 by Sir2 α in the presence of TSA. Western blot analysis of acetylated p53 levels in H1299 cells cotransfected with p53 and p300 (lanes 1, 3), or cotransfected with p53, p300 and Sir2 α (lanes 2, 4) by

acetylated p53-specific antibody (upper) or OF-1 for total p53 (lower). Cells were either not treated (lanes 1, 2) or treated with 500 nM TSA (lanes 3, 4).

Figure 4. Abrogation of mammalian Sir2 α mediated deacetylation of p53 by nicotinamide.

(A) Sir2 α -mediated deacetylation of p53 is inhibited by nicotinamide. 2.5 μ g of ¹⁴C-labeled acetylated p53 (lane 1) was incubated with 10 ng of purified Sir2 α and 50 μ M NAD alone (lane 2), or in the presence of either 5mM of nicotinamide (lane 3) or 3 mM of 3-AB (3-aminobenzamide) (lane 4) for 60 min at 30°C. The proteins were analyzed by resolution on SDS-PAGE and autoradiography (upper) or Coomassie blue staining (lower). (B) The Sir2 α -mediated deacetylation of endogenous p53 was abrogated in the presence of nicotinamide. Cell extracts from the mock-infected MEF p53 (+/+) cell (lanes 1-2, 5-6), or the p/Babe-Sir2 α infected cells (lanes 3-4, 7-8), either untreated (lanes 1, 3, 5, 7), or treated with etoposide and TSA (lane 2, 4), or in combination with nicotinamide (lanes 6, 8) for 6 hr were analyzed by Western blot with acetylated p53-specific antibody (upper) or DO-1 for total p53 (lower). (C) Synergistic induction of p53 acetylation levels by TSA and nicotinamide during DNA damage response. Western blot analysis of cell extracts from the H460 cells treated with etoposide alone (lane 2), or in combination with TSA (lane 3), or TSA and nicotinamide (lane 4), or TSA and 3-AB (lane 5) for 6 hr by acetylated p53-specific antibody (upper) or DO-1 for total p53 (lower). The cell extracts from untreated cells (lane 1), or treated with a proteasome inhibitor LLNL (50 μ M) were also included (lane 6).

Figure 5. Bar graphs illustrating repression of p53-mediated transcriptional activation by mammalian Sir2 α .

(A), (B) MEF (p53^{-/-}) cells were transiently transfected with 10 ng of CMV-p53 alone, or in combination with indicated Sir2 α constructs together with either the PG13-Luc reporter construct (A), or a control reporter construct (TK-Luc) (B) by calcium phosphate precipitation essentially as previously described (Luo *et al.*, 2000). (C), (D) MEF (p53^{-/-}) cells were transiently transfected with 10 ng of CMV-p53 alone, or in combination with 5 μ g of either CMV-Sir2 α , or CMV-hSIRT1, or CMV-hSIRT5 (C), or CMV-Sir2 α H355A as indicated (D) together with the PG13-Luc reporter construct. All transfections were done in duplicate and

representative experiments depict the average of three experiments with standard deviations indicated.

Figure 6. Inhibition of p53-dependent apoptosis by Sir2 α .

(A) H1299 cells were transfected with p53 alone, or cotransfected with p53 and Sir2 α , or cotransfected with p53 and Sir2 α H355A. After transfection, the cells were fixed, stained for p53 by FITC-conjugated α -p53 antibody, and analyzed by flow cytometry for apoptotic cells (subG1) according to DNA content (PI staining). (B) The experiments were repeated at least three times; this bar graph depicts the average of three experiments with standard deviations indicated.

Figure 7. Inhibition of p53-dependent apoptotic response to stress by mammalian Sir2 α .

(A) Repression of the apoptotic response to DNA damage by Sir2 α . Both mock infected cells and p/babe-Sir2 α infected MEF p53(+/-) cells were either not treated (1 and 2) or treated with either 20 μ M etoposide. The cells were analyzed by flow cytometry for apoptotic cells (subG1) according to DNA content (PI staining). (B) Similar results were obtained for three times, and this bar graph of representative data depicts the average of three experiments with standard deviations indicated (B).

Figure 8. Co-precipitation of hSir2 and p53 protein.

(A) Immunoprecipitation of hSir2 with a C-terminal polyclonal rabbit antibody followed by immunoblotting with the same antibody revealed the existence of a 120Kd protein in normal BJ fibroblasts (left panel), and increased levels in these cells expressing the wild type (middle panel) and HY mutant (right panel) of hSir2. (B) Immunofluorescence analysis of hSir2 indicated the existence of a nuclear protein with a punctuate staining pattern. (C) Nuclear lysates from H1299 cells ectopically expressing p53 and hSir2 were precipitated with the anti-hSir2 antibody. The blot was probed the anti-hSir2 antibody and a polyclonal sheep anti-p53 antibody (bottom panel). (D) p53 protein was immunoprecipitated with the Do-1 anti-p53 antibody from lysates of non-irradiated and irradiated (6Gy) BJT cells (expressing telomerase) that had been stably infected with pYESir2wt and pYESir2HY mutant vectors. The blot was probed with anti-hSir2 antibody and rabbit anti-p53 polyclonal antibodies (CM1+SC6243).

Figure 9. Effect of hSir2 expression on p53 acetylation *in vitro*.

The deacetylation activity of mSir2 on the human p53 C-terminal peptide (residues 368-386) di-acetylated at positions 373 and 382. (A, B) HPLC chromatograms of products of deacetylation assays with mSir2 and the indicated concentrations of NAD. Peaks 1 and 2 correspond to the monomeric and dimeric forms of the p53 peptide, respectively. Peak 3 corresponds to the singly deacetylated monomer identified by mass spectroscopy. (C-F) Amino-terminal Edman sequencing of peaks 1 and 3. Chromatograms of positions 373 and 382 are shown. Peaks of acetyl-lysine (AcK) and simple lysine (K) are indicated in each panel. Small peaks of lysine in panels C, D and F are due to residual fractions of previous lysines at positions 372 and 381.

Figure 10. hSir2 effects on p53 acetylation *in vivo*.

(A) Reconstitution of the acetylation and deacetylation cascade in immortal human epithelial H1299 cells by transient co-transfection of the indicated genes. After co-transfection of the mentioned constructs, the cellular lysates were analyzed by Western blot analysis, using Ab-1 to detect K382 p53, DO-1 for total p53 or β actin for loading control. Lane 3, co-transfection of CMVwtp53 and p300 generates acetylated p53 at K382, lane 4, co-transfection of the acetylation mutant K382R of p53 with p300. Lane 5, Same as 4 but with co-transfected wild type hSir2. Lanes 7-8, co-transfection of the acetylation mutant K320R with or without wild type hSir2. Lane 9, Co-transfection of CMVwtp53, CMVp300 and wild type hSir2.

(B) BJ cells expressing telomerase (BJT), were stably infected with either a wild type hSir2 or a mutant hSir2HY virus. The hSir2-expressing mass cultures were subjected to 6Gy of ionizing radiation in presence of low concentrations of TSA (0.1mg/ml) and the p53 acetylation was measured at indicated time points by immunoblotting with Ab-1 that recognizes specifically the deacetylated K382 p53 protein. The blots were subsequently probed with anti-p53, anti-p21, anti- β -actin and anti-hSir2 antibodies. Time (hrs) post 6 Gy of irradiation is shown inside the brackets.

(C) Deacetylation of p53 *in vivo* in MCF7 cells. Four-fold ectopic expression of wild type hSir2 or hSir2HY mutant in MCF7 cells radiated with 6Gy of ionizing radiation and its

effect on p53 acetylation at K382. The blot was probed for acetylation with Ab-1 and reprobed with other antibodies as in (B). Times shown are post irradiation in hours.

Figure 11. hSir2 expression and its influence on p53 activity

(A) is a bar graph depicting transcriptional activity of p53 protein, as measured in H1299 cells by co-transfection p53 with a p21WAF1 promoter-luciferase construct (p21Pluc). Transcriptional activity of p53 protein was measured upon ectopic expression of wild type hSir2, hSir2HY. (B) is a bar graph illustrating results from control SV40-Luciferase transfections with CMVp53 and increasing amounts of wild type hSir2 in to H1299 cells and luciferase activity was measured and expressed as Relative Light Unit (%RLU). (C) Is an immunoblot demonstrating levels of p21WAF1 in MCF73L cells expressing wt hSir2 or hSir2HY protein in response to 6Gy of ionizing radiation. The blot was probed with Do1 for detection of p53 and β actin for loading control.

Figure 12. Effects of hSir2 on p53-dependent apoptosis and radiosensitivity

(A) is a bar graph illustrating ectopic expression of hSir2wt and its influence on p53-dependent apoptosis in H1299 cells. H1299 cells were transfected with a wild type p53 expression construct to induce p53-dependent apoptosis. Annexin V positive and propidium iodide negative cells were measured.

(B) is a line graph comparison of gamma-ray survival. Dose-response curves are shown for different types of BJ cells treated with ionizing radiation while growing exponentially and asynchronously. Twelve days after radiation the colonies were counted and survival calculated as described previously (Dhar *et al.*, 2000). The ataxia-telangiectasia (A-T) cell line was used a positive control to indicate radiosensitivity in an exponentially growing population.

Figures 13A and 13B. The coding nucleic acid (SEQ ID NO. 2) and deduced amino acid (SEQ ID NO. 3) of human p53.

Figure 14. The nucleic acid (SEQ ID NO. 4) sequence of human p53 (GenBank Accession No: K03199).

Figures 15A, B, C and D. The nucleic acid (SEQ ID NO. 5) and deduced amino acid sequence (SEQ ID NO. 6) of mouse Sir2.

Figures 16A, B and C. The nucleic acid (SEQ ID NO. 7) and deduced amino acid sequence (SEQ ID NO. 8) of mouse Sir2 GenBank Accession No: AF214646.

Figures 17A and B. The nucleic acid (SEQ ID NO. 9) and deduced amino acid sequence (SEQ ID NO. 10) of human Sir2 SIRT2 GenBank Accession No: AF083107.

Figures 18A, B and C. The nucleic acid (SEQ ID NO. 11) and deduced amino acid sequence (SEQ ID NO. 12) of human Sir2 SIRT1 GenBank Accession No: AF083106.

Figure 19. The nucleic acid (SEQ ID NO. 13) and deduced amino acid sequence (SEQ ID NO. 14) of human Sir2 SIRT3 GenBank Accession No: AF083108.

Figures 20A and B. The nucleic acid (SEQ ID NO. 15) and deduced amino acid sequence (SEQ ID NO. 16) of human Sir2 SIRT4 GenBank Accession No: AF083109.

Figures 21A and B. The nucleic acid (SEQ ID NO. 17) and deduced amino acid sequence (SEQ ID NO. 18) of human Sir2 SIRT5 GenBank Accession No: AF083110.

DETAILED DESCRIPTION

As described below, hSir2 directly binds the human p53 protein both *in vitro* and *in vivo* and can deacetylate p53, e.g., at the K382 residue of p53. A functional consequence of this deacetylation is an attenuation of the p53 protein's activity, e.g., as a transcription factor operating at a cellular promoter, e.g., the p21WAF1 promoter. In another cellular context, in which the DNA damage response leads to apoptosis, hSir2 activity attenuates the p53-dependent apoptotic response. Hence, hSir2 can negatively regulate a program of cellular death.

Sir2 proteins can also deacetylate histones. For example, Sir2 can deacetylate lysines 9 or 14 of histone H3. Histone deacetylation alters local chromatin structure and consequently can

regulate the transcription of a gene in that vicinity. Sir2 proteins can bind to a number of other proteins, termed “Sir2-binding partners.” For example, hSIRT1 binds to p53. In many instances the Sir-2 binding partners are transcription factors, e.g., proteins that recognize specific DNA sites. Interaction between Sir2 and Sir2-binding partners delivers Sir2 to specific regions of a genome and can result in local modification of substrates, e.g., histones and transcription factors localized to the specific region. Accordingly, cellular processes can be regulated by compounds that alter (e.g., enhance or diminish) the ability of a Sir2 protein to interact with a Sir2-binding partner or that alter that ability of a Sir2 protein to modify a substrate. While not wishing to be bound by theory, a Sir2-transcription factor complex may be directed to a region of DNA with a transcription factor binding site; once there, Sir2 may alter the acetylation status of the region, e.g., by deacetylating histones, non-histone proteins, and/or DNA. This would locally raise the concentration of Sir2 and may potentially result in the Sir2-mediated silencing of genes located at or near transcription-factor binding sites. Certain organismal programs such as aging or metabolism and disorders such as cancer can be controlled using such compounds.

While not wishing to be bound by theory, in mammalian cells, signals indicating the successful completion of DNA repair may be relayed via hSir2 to acetylated proteins like p53 that have been charged with the task of imposing a growth arrest following DNA damage. These signals enable hSir2 to reverse part or all of the damage-induced activation of p53 as a transcription factor by deacetylating the K382 residue of p53. By doing so, hSir2 reduces the likelihood of subsequent apoptosis and, at the same time, makes it possible for cells to re-enter the active cell cycle, enabling them to return to the physiological state that they enjoyed prior to sustaining damage to their genomes.

Inactivation of the p53 signaling pathway is involved in the pathogenesis of most if not all human tumors (Hollstein *et al.*, 1994; Lohrum and Vousden, 1999). In about half of these tumors, mutation of the p53 gene itself suffices to derail function. In some of the remaining tumors, loss of p14^{ARF}, which acts to down-regulate p53 protein levels, has been implicated (Lohrum and Vousden, 1999; Prives and Hall, 1999). The present invention is related to the discovery of a novel mode by which an incipient cancer cell attenuate at least some p53 functions via modulation of the activity of hSir2, which, like the other two genetic strategies, may result in the inactivation of both the cytostatic and pro-apoptotic functions of p53.

The invention is thus based in part on the discovery of the existence of a p53 regulatory pathway that is regulated by mammalian Sir2 α . Sir2 α is involved in gene silencing and extension of life span in yeast and *C. elegans* (reviewed in Guarente, 2000; Shore, 2000; Kaeberlein *et al.*, 1999; Tissenbaun and Guarente, 2001). p53 binds to mouse Sir2 α as well as its human ortholog hSIRT1 both *in vitro* and *in vivo*. p53 is a substrate for the NAD-dependent deacetylase activity of mammalian Sir2 α . Sir2 α -mediated deacetylation antagonizes p53-dependent transcriptional activation and apoptosis. Sir2 α -mediated deacetylation of p53 is inhibited by nicotinamide both *in vitro* and *in vivo*. Sir2 α specifically inhibits p53-dependent apoptosis in response to DNA damage and/or oxidative stress, but not p53-independent, Fas-mediated cell death. Accordingly, compounds that alter (e.g., decrease or enhance) the interaction between Sir2 and p53 can be used to regulate processes downstream of p53, e.g., apoptosis. Such compounds may alter the catalytic activity of Sir2 for a substrate such as p53 or may alter the interaction between Sir2 and p53.

The present invention relates to the discovery that p53 is a binding partner of mammalian Sir2 α , which physically binds to p53 both *in vitro* and *in vivo*. In some cases, p53 is also a substrate of Sir2. Sir2 α specifically represses p53-mediated functions including p53-dependent apoptotic response to stress.

p53 can be, for example, the mature protein (e.g., SEQ ID NO. 3) or a fragment thereof. The p53 protein can be encoded by the nucleic acid sequence of SEQ ID NOS. 2 and/or 4). In a preferred embodiment, p53 is the human p53. Deacetylation of p53 can be mediated by Sir2, e.g., in combination with a cofactor, such as NAD and/or an NAD analog.

The phrase "deacetylating p53" refers to the removal of one or more acetyl groups (e.g., CH₃CO²⁻) from p53 that is acetylated on at least one amino acid residue. In a preferred embodiment, p53 is deacetylated at a lysine of p53 selected from the group consisting of lysine 370, lysine 371, lysine 372, lysine 381 and lysine 382 of SEQ ID NO. 3. p53 can be deacetylated in the presence or absence of DNA damage or oxidative cellular stress. The DNA damage can be caused by, for example, ionizing radiation (e.g., 6 Gy of ionizing radiation), or a tumor or some other uncontrolled cell proliferation. p53 is deacetylated in the presence of DNA damage or oxidative stress by combining p53, Sir2, NAD and/or an NAD analog.

Sir2 can be the mature protein (e.g., SEQ ID NOS. 8, 10, 12, 14, 16 or 18) or a fragment of the mature protein capable of deacetylating p53 in the presence or NAD and/or an NAD

analog. The Sir2 protein can be encoded by the nucleic acid sequence of SEQ ID NOS. 7, 9, 11, 13, 15 or 17). In a preferred embodiment, the Sir2 is human Sir2.

In one embodiment, the invention is a method of deacetylating p53 comprising the step of combining Sir2 and NAD and/or an NAD analog with p53. The combination can be performed in the presence or the absence of cells. Such combinations can be in tissue culture (*e.g.*, BJT cells, MCF-7 cells) or in an organism (*e.g.*, a mammal, *e.g.*, as a human). Combination of p53, Sir2 and NAD and/or an NAD analog can be any placement of p53, Sir 2 and NAD or a NAD analog in sufficient proximity to cause Sir2 to deacetylate p53 that is acetylated on at least one amino acid residue, which deacetylation by Sir2 requires the presence of NAD and/or an NAD analog.

“NAD” refers to nicotinamide adenine dinucleotide. An “NAD analog” as used herein refers to a compound (*e.g.*, a synthetic or naturally occurring chemical, drug, protein, peptide, small organic molecule) which possesses structural similarity to component groups of NAD (*e.g.*, adenine, ribose and phosphate groups) or functional similarity (*e.g.*, deacetylates p53 in the presence of Sir2). For example, an NAD analog can be 3-aminobenzamide or 1,3-dihydroisoquinoline (H. Vaziri *et al.*, EMBO J. 16:6018-6033 (1997), the entire teachings of which are hereby incorporated by reference).

“p53 activity” refers to one or more activity of p53, *e.g.*, p-53 mediated apoptosis, cell cycle arrest, and/or senescence,

“Modulating p53 activity” refers to increasing or decreasing p53 activity, *e.g.*, p-53 mediated apoptosis, cell cycle arrest, and/or senescence, *e.g.* by altering the acetylation or phosphorylation status of p53.

“Acetylation status” refers to the presence or absence of one or more acetyl groups (*e.g.*, CH_3CO^2^-) at one or more lysine (K) residues, *e.g.*, K370, K371, K372, K381, and/or K382 of SEQ ID NO. 3. “Altering the acetylation status” refers to adding or removing one or more acetyl groups (*e.g.*, CH_3CO^2^-) at one or more lysine (K) residues, *e.g.*, K370, K371, K372, K381, and/or K382 of SEQ ID NO. 3, *e.g.*, by modulating Sir2 activity.

Similarly, “phosphorylation status” refers to the presence or absence of one or more phosphate groups (PO_3^-) at one or more residues, *e.g.*, serine 15 and/or serine 20 of SEQ ID NO. 3. “Altering the phosphorylation status” refers to adding or removing one or more phosphate groups (PO_3^-) at one or more residues, *e.g.*, serine 15 and/or serine 20 of SEQ ID NO. 3.

“Sir2 activity” refers to one or more activity of Sir2, *e.g.*, deacetylation of p53 or histone proteins.

“Modulating Sir2 activity” refers to increasing or decreasing one or more activity of Sir2, *e.g.*, deacetylation of p53 or histone proteins, *e.g.*, by altering the binding affinity of Sir2 and p52, introducing exogenous Sir2 (*e.g.*, by expressing or adding purified recombinant Sir2), increasing or decreasing levels of NAD and/or an NAD analog (*e.g.*, 3-aminobenzamide, 1,3-dihydroxyisoquinoline), and/or increasing or decreasing levels of a Sir2 inhibitor, *e.g.*, nicotinamide and/or a nicotinamide analog. Additionally or alternatively, modulating Sir2 activity can be accomplished by expressing, *e.g.* by transfection, a dominant negative gene of Sir2 (*e.g.*, SirHY). The dominant negative gene can, for example, reduce the activity of endogenous Sir2 on p53 deacetylation thereby modulating the activity of Sir2.

A “nicotinamide analog” as used herein refers to a compound (*e.g.*, a synthetic or naturally occurring chemical, drug, protein, peptide, small organic molecule) which possesses structural similarity to component groups of nicotinamide or functional similarity (*e.g.*, reduces Sir2 deacetylation activity of p53).

The Sir2 α -mediated pathway is critical for cells under stress

It is believed that there are multiple pathways in cells for regulation of p53 function (Prives and Hall, 1999; Giaccia and Kastan, 1998; Ashcroft *et al.*, 2000). In normal cells, Mdm2 is the major negative regulator for p53, and Mdm2-mediated repression appears sufficient to downregulate p53 activity. Sir2 regulation of p53 may be an Mdm2-independent, negative regulatory pathway for p53. Interestingly, while no obvious effect by Sir2 α expression was observed in cells at normal conditions, Sir2 α became critical in protecting cells from apoptosis when cells were either treated by DNA damage or under oxidative stress (Figure 7). Thus, Sir2 α -mediated pathway can be critical for cell survival when the p53 negative-control mediated by Mdm2 is severely attenuated in response to DNA damage or other types of stress.

p53 is often found in latent or inactive forms and the levels of p53 protein are very low in unstressed cells, mainly due to the tight regulation by Mdm2 through functional inhibition and protein degradation mechanisms (reviewed in Freedman *et al.*, 1999). However, in response to DNA damage, p53 is phosphorylated at multiple sites at the N-terminus; these phosphorylation

events contribute to p53 stabilization and activation by preventing Mdm2 binding to p53 (reviewed in Appella and Anderson, 2000; Giaccia and Kastan, 1998; Shieh *et al.*, 1997, 2000; Unger *et al.*, 1999; Hirao *et al.*, 2000). Mdm2 itself is also phosphorylated by ATM during DNA damage response, and this modification attenuates its inhibitory potential on p53 (Maya *et al.*, 2001). Furthermore, while p53 is strongly stabilized and highly acetylated in stressed cells, acetylation of the C-terminal multiple lysine sites may occur at the same sites responsible for Mdm2-mediated ubiquitination (Rodriguez *et al.*, 2000; Nakamura *et al.*, 2000), and the highly acetylated p53 may not be effectively degraded by Mdm2 without deacetylation (Ito *et al.*, 2001). Thus, in contrast to unstressed cells, the main p53 negative regulatory pathway mediated by Mdm2 is blocked at several levels in response to DNA damage (Maya *et al.*, 2001). Under these circumstances, Sir2 α -mediated regulation may become a major factor in controlling p53 activity, making it possible for cells to adjust p53 activity to allow time for DNA repair before committing to apoptosis.

In oncogene-induced premature senescence of cells, the p53 negative regulatory pathway controlled by Mdm2 may be blocked (reviewed in Sherr and Weber, 2000; Sharpless and Depinho, 1999; Serrano *et al.*, 1997). However, in contrast to DNA damage response, the Mdm2-mediated pathway is abrogated by induction of p14^{ARF} (or mouse p19^{ARF}) in these cells (Honda and Yasuda, 1999; Weber *et al.*, 1999; Tao *et al.*, 1999a, 1999b; Zhang *et al.*, 1998; Pomerantz *et al.*, 1998). Furthermore, when primary fibroblasts undergo senescence, a progressive increase of the p53 acetylation levels was observed in serially passaged cells (Pearson *et al.*, 2000). Oncogenic Ras and PML induced p53-dependent premature senescence, and upregulated the p53 acetylation levels in both mouse and human normal fibroblasts (Pearson *et al.*, 2000; Ferbeyre *et al.*, 2000). Thus, mammalian Sir2 α -mediated regulation may also play an important role in oncogene-induced premature senescence.

Attenuation of p53-mediated transactivation by Sir2 α

Earlier studies indicated that p53-mediated transcriptional activation is sufficient and also absolutely required for its effect on cell growth arrest, while both transactivation-dependent and -independent pathways are involved in p53-mediated apoptosis (reviewed in Prives and Hall, 1999; Vousden, 2000). p53 may be effective to induce apoptosis by activating pro-apoptotic genes *in vivo* (reviewed in Nakano and Vousden, 2001; Yu *et al.*, 2001). Thus, tight regulation

of p53-mediated transactivation is critical for its effect on both cell growth and apoptosis (Chao *et al.*, 2000; Jimenez *et al.*, 2000).

Recent studies indicate that the intrinsic histone deacetylase activity of Sir2 α is essential for its mediated functions (reviewed in Gurante, 2000). Reversible acetylation was originally identified in histones (reviewed in Cheung *et al.*, 2000; Wolffe *et al.*, 2000); however, accumulating evidence indicates that transcriptional factors are also functional targets of acetylation (reviewed in Serner and Berger, 2000; Kouzarides, 2000). Thus, the transcriptional attenuation mediated by histone deacetylases may act through the effects on both histone and non-histone transcriptional factors (Serner and Berger, 2000; Kuo and Allis, 1998). Microarray surveys for transcriptional effects of Sir2 in yeast revealed that Sir2 appears to repress amino acid biosynthesis genes, which are not located at traditional "silenced" loci (Bernstein *et al.*, 2000). Thus, in addition to silencing (repression) at telomeres, mating type loci and ribosomal DNA (reviewed in Guarente, 2000; Shore, 2000), Sir2 may also be targeted to specific endogenous genes for transcriptional regulation in yeast.

In contrast to the yeast counterpart Sir2, the mouse Sir2 α protein does not colocalize with nucleoli, telomeres or centromeres by co-immunofluorescence assay, indicating that this protein is not associated with the most highly tandemly repeated DNA in the mouse genome. The immunostaining pattern of human SIRT1 as well as mouse Sir2 α indicates that mammalian Sir2 α is, similar to HDAC1, broadly localized in the nucleus, further supporting the notion that mammalian Sir2 α may be recruited to specific target genes for transcriptional regulation *in vivo*.

Mammalian Sir2 α may inhibit p53-mediated functions by attenuation of the transcriptional activation potential of p53. Since deacetylation of p53 is critical, but may not be the only function mediated by this Sir2 α -p53 interaction, additional functions mediated by Sir2 α , such as histone deacetylation, may also contribute to this regulation. As one theory, not meant to be limiting, p53 and Sir2 α may strongly interact to deacetylate p53 and possibly recruit the p53-Sir2 α complex to the target promoter. The subsequent transcription repression may act both through decreasing p53 transactivation capability and through Sir2 α -mediated histone deacetylation at the target promoter region. In contrast to HDAC1-mediated effect, this transcriptional regulation is not affected by TSA treatment. Other cellular factors may use a similar mechanism to recruit Sir2 α for TSA-insensitive transcriptional regulation in mammalian cells.

Novel implications for cancer therapy

Inactivation of p53 functions has been well documented as a common mechanism for tumorigenesis (Hollstein *et al.*, 1999; Vogelstein *et al.*, 2000). Many cancer therapy drugs have been designed based on either reactivating p53 functions or inactivating p53 negative regulators. Since p53 is strongly activated in response to DNA damage, mainly through attenuation of the Mdm2-mediated negative regulatory pathway (Maya *et al.*, 2001), many DNA damage-inducing drugs such as etoposide are very effective antitumor drugs in cancer therapy (reviewed in Chresta and Hickman, 1996; Lutzker and Levine, 1996). Maximum induction of p53 acetylation in normal cells, however, requires both types of deacetylase inhibitors in addition to DNA damage, and there may be at least three different p53 negative regulatory pathways in mammalian cells. Inhibitors for HDAC-mediated deacetylases, including sodium butyrate, TSA, SAHA and others, have been also proposed as antitumor drugs (Butler *et al.*, 2000; Finnin *et al.*, 1999; Taunton *et al.*, 1996; Yoshida *et al.*, 1995; Buckley *et al.*, 1996). Combining DNA damage drugs, HDAC-mediated deacetylase inhibitors, and Sir2 α -mediated deacetylase inhibitors, may have synergistic effects in cancer therapy for maximally activating p53.

In contrast to PID/HDAC1-mediated p53 regulation (Luo *et al.*, 2000), the invention shows that mammalian Sir2 α -mediated effect on p53 is NAD-dependent, indicating that this type of regulation is closely linked to cellular metabolism (reviewed Guarente 2000; Alfred, 2000; Campisi, 2000; Min *et al.*, 2001). In fact, null mutants of NPT1, a gene that functions in NAD synthesis, show phenotypes similar to that of Sir2 mutants in gene silencing (Smith *et al.*, 2000) and in life extension in response to caloric restriction in yeast (Lin *et al.*, 2000). Thus, metabolic rate may play a role in Sir2 α -mediated regulation of p53 function and, perhaps, modulate the sensitivity of cells in p53-dependent apoptotic response.

In yet another embodiment, the invention is a method of modulating p53-mediated apoptosis by modulating Sir2 activity. Sir2 activity can be modulated as described herein (*e.g.*, overexpressing Sir2, transfecting a cell with a dominant negative regulating gene). An increase in Sir2 activity (*e.g.*, by overexpressing Sir2) can result in a decrease in p53-mediated apoptosis.

A decrease in Sir2 activity (*e.g.*, transfecting a cell with a dominant negative gene) can result in an increase in p53-mediated apoptosis.

In still another embodiment, the invention is a method of screening for a compound (*e.g.*, a small organic or inorganic molecule) which modulates (*e.g.*, increases or decreases) Sir2-mediated deacetylation of p53. In the method, Sir2, p53, NAD and/or an NAD analog, and the compound to be tested are combined, the Sir2-mediated deacetylation of p53 is measured and compared to the Sir2-mediated deacetylation of p53 measured in the absence of the compound. An increase in the Sir2-mediated deacetylation of p53 in the presence of the compound being tested compared to the Sir2-mediated deacetylation of p53 in the absence of the compound indicates that the compound increases Sir2 deacetylation of p53. Likewise, a decrease in the Sir2-mediated deacetylation of p53 in the presence of the compound being tested compared to the Sir2-mediated deacetylation of p53 in the absence of the compound indicates that the compound decreases deacetylation of p53 by Sir2. As used herein, "Sir2-mediated deacetylation" refers to the NAD-dependent removal of acetyl groups which requires Sir2.

In another embodiment, the present invention relates to a method of screening a compound by providing an *in vitro* test mixture comprising a transcription factor or a fragment thereof, Sir2, and a Sir2 cofactor with the compound, evaluating an activity of a component of the test mixture in the presence of the compound, and comparing the activity in the presence of the compound to a reference obtained in the absence of the compound.

In another embodiment, the present invention relates to a method of screening a compound that is a potential NAD analog by providing an *in vitro* test mixture comprising a transcription factor or a fragment thereof, Sir2, and the compound, evaluating an activity of a component of the test mixture in the presence of the compound, and comparing the activity in the presence of the compound to a reference obtained in the absence of the compound,

In one embodiment the Sir2 is human, *e.g.*, human SIRT1. In another embodiment, the Sir2 is murine, *e.g.*, murine Sir2 α .

In one embodiment the Sir2 cofactor is NAD or an NAD analog.

In another embodiment the transcription factor is p53 or a fragment thereof, and it may be acetylated and/or labeled.

In a further embodiment, the evaluated activity is Sir2 activity, *e.g.*, deacetylation of a protein, *e.g.*, deacetylation of a histone protein, and/or deacetylation of the transcription factor,

e.g., deacetylation of p53. The Sir2 activity may also be binding of a protein, e.g., binding of a histone protein and/or binding of the transcription factor, e.g., binding of p53. The Sir2 activity may be evaluated by detecting production of nicotinamide.

In a further embodiment, the evaluated activity is p53 activity. The p53 activity may be evaluated by detecting cell cycle arrest, apoptosis, senescence, and/or a change in the levels of transcription or translation products of a gene regulated by p53. Methods for detecting such changes and genes regulated by p53 are known in the art and include those methods and genes disclosed in U.S. Pat. No. 6,171,789, which is incorporated herein by reference in its entirety.

In one embodiment, the test mixture is provided in a cell-free system.

In another embodiment, the test mixture is provided in a cell-based system, wherein one of the components is exogenous. The term "exogenous" refers to a component that is either added directly, or expressed from a heterologous DNA source, such as transfected DNA. Many methods are known in the art for expression of heterologous or exogenous gene products.

In a further embodiment, the evaluated activity is an effect on the rate of aging of a cell or organism. Such an effect may be evaluated by contacting the compound with a cell or organism having p53 or Sir2 activity, e.g., endogenous or exogenous p53 or Sir2 activity; and evaluating the rate of aging of the cell or organism. The rate of aging may be evaluated by several methods, including:

- a) assessing the life span of the cell or organism;
- b) assessing the presence or absence of a gene transcript or gene product in the cell or organism that has a biological age-dependent expression pattern;
- c) evaluating resistance of the cell or organism to stress;
- d) evaluating one or more metabolic parameters of the cell or organism;
- e) evaluating the proliferative capacity of the cell or a set of cells present in the organism;
- f) evaluating physical appearance, behavior, or other characteristic of the cell or organism; and
- (g) assessing the presence or absence of a gene transcript or gene product in the cell or organism that has a p53-regulation-dependent expression pattern.

The compounds identified by the methods of the invention can be used, for example, to treat cancer (e.g., a compound which decreases Sir2-mediated deacetylation of p53) or prevent

p53-mediated apoptosis (*e.g.*, a compound which increases Sir2-mediated deacetylation of p53). The compounds can be used in methods of treating a cell or an organism, *e.g.*, a cell or organism that has been exposed to DNA-damaging ionizing radiation, by modulating Sir2 activity in the cell. In the method of treating cancer in a mammal, Sir2 activity can be reduced. In a preferred embodiment, Sir2 activity is reduced by nicotinamide or a nicotinamide analog.

In yet another embodiment, the invention is a method of screening for analogs of NAD. In the method, Sir2, p53 and a compound to be tested as an analog of NAD (*e.g.*, a small organic or inorganic molecule) are combined. Deacetylation of the p53 by the Sir2 is measured and compared to the measured deacetylation of p53 by Sir2 in the presence of NAD. A compound which, for example, promotes Sir2-mediated deacetylation of p53 when combined with Sir2 and p53, is an NAD analog and can be used in place of NAD, for example, as a cofactor with Sir2 to prevent or decrease p53-mediated apoptosis.

In a further embodiment, the invention is a method of treating cancer in a mammal comprising the step of modulating Sir2 activity in tumor cells to cause an increase in p53 activity. The Sir2 activity can be modulated as described herein (*e.g.*, overexpression of Sir2, transfection of a cell with a dominant negative regulatory gene, or nicotinamide or a nicotinamide analog).

In another embodiment, the invention includes a method of treating a cell that has been exposed to ionizing radiation, the method comprising modulating Sir2 activity in the cell. In a particular embodiment, in a cell which has undergone DNA damage or oxidative stress, Sir2 activity can be modulated to reduce Sir2 activity (*e.g.*, by transfecting a cell with a dominant negative regulatory gene, or by addition or expression of nicotinamide or a nicotinamide analog) which can result in the arrest of the growth cycle of the cell, allowing the cell to repair at least a portion of the DNA damage caused by the ionizing radiation. Once the cell has repaired a portion of the DNA damage, the reduction in Sir2 activity can be removed and the cell cycle of the cell resumed.

In still another embodiment, the invention includes an isolated protein complex of Sir2 and acetylated p53. p53 can also be phosphorylated (*e.g.*, on one or both of serine 15 or serine 20 of SEQ ID NO. 3).

The compounds or NAD analogs identified by the methods of the invention can be used in the treatment of diseases or conditions such as cancer, or following DNA damage or oxidative

stress. The compounds or NAD analogs can be administered alone or as mixtures with conventional excipients, such as pharmaceutically, or physiologically, acceptable organic, or inorganic carrier substances such as water, salt solutions (*e.g.*, Ringer's solution), alcohols, oils and gelatins. Such preparations can be sterilized and, if desired, mixed with lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, and/or aromatic substances and the like which do not deleteriously react with the NAD analogs or compounds identified by the methods of the invention.

The dosage and frequency (single or multiple doses) of the compound or NAD analog administered to a mammal can vary depending upon a variety of factors, including the duration of DNA damage, oxidative stress or cancer condition.

In some embodiments of the present invention, the rate of aging of a cell, *e.g.*, a yeast cell, invertebrate cell (*e.g.*, fly cell), or vertebrate cell (*e.g.*, mammalian cell, *e.g.*, human or mouse cell) is determined. For example, the rate of aging of the cell can be evaluated by measuring the expression of one or more genes or proteins (*e.g.*, genes or proteins that have an age-related expression pattern), by measuring the cell's resistance to stress, *e.g.*, genotoxic stress or oxidative stress, by measuring one or more metabolic parameters (*e.g.*, protein synthesis or degradation, ubiquinone biosynthesis, cholesterol biosynthesis, ATP levels within the cell, glucose metabolism, nucleic acid metabolism, ribosomal translation rates, etc.), by measuring cellular proliferation, or any combination of measurements thereof.

In other embodiments, the rate of aging of an organism, *e.g.*, an invertebrate (*e.g.*, a worm or a fly) or a vertebrate (*e.g.*, a rodent, *e.g.*, a mouse) is determined. The rate of aging of an organism can be determined by directly measuring the average life span of a group of animals (*e.g.*, a group of genetically matched animals) and comparing the resulting average to the average life span of a control group of animals (*e.g.*, a group of animals that did not receive the test compound but are genetically matched to the group of animals that did receive the test compound). Alternatively, the rate of aging of an organism can be determined visually, *e.g.*, by looking for visible signs of age (*e.g.*, physical appearance or behavior), by measuring the expression of one or more genes or proteins (*e.g.*, genes or proteins that have an age-related expression pattern), by measuring the cell's resistance to genotoxic (*e.g.*, caused by exposure to etoposide, UV irradiation, mutagens, etc.) or oxidative stress, by measuring one or more metabolic parameters (*e.g.*, protein synthesis or degradation, ubiquinone biosynthesis, cholesterol

biosynthesis, ATP levels, glucose metabolism, nucleic acid metabolism, ribosomal translation rates, etc.), by measuring cellular proliferation (*e.g.*, of retinal cells, bone cells, white blood cells, etc.), or any combination of measurements thereof. In one embodiment, the visual assessment is for evidence of apoptosis, *e.g.*, nuclear fragmentation.

All animals typically go through a period of growth and maturation followed by a period of progressive and irreversible physiological decline ending in death. The length of time from birth to death is known as the life span of an organism, and each organism has a characteristic average life span. Aging is a physical manifestation of the changes underlying the passage of time as measured by percent of average life span.

In some cases, characteristics of aging can be quite obvious. For example, characteristics of older humans include skin wrinkling, graying of the hair, baldness, and cataracts, as well as hypermelanosis, osteoporosis, cerebral cortical atrophy, lymphoid depletion, thymic atrophy, increased incidence of diabetes type II, atherosclerosis, cancer, and heart disease. Nehlin *et al.* (2000), *Annals NY Acad Sci* 980:176-79. Other aspects of mammalian aging include weight loss, lordokyphosis (hunchback spine), absence of vigor, lymphoid atrophy, decreased bone density, dermal thickening and subcutaneous adipose tissue, decreased ability to tolerate stress (including heat or cold, wounding, anesthesia, and hematopoietic precursor cell ablation), liver pathology, atrophy of intestinal villi, skin ulceration, amyloid deposits, and joint diseases. Tyner *et al.* (2002), *Nature* 415:45-53.

Careful observation reveals characteristics of aging in other eukaryotes, including invertebrates. For example, characteristics of aging in the model organism *C. elegans* include slow movement, flaccidity, yolk accumulation, intestinal autofluorescence (lipofuscin), loss of ability to eat food or dispel waste, necrotic cavities in tissues, and germ cell appearance.

Those skilled in the art will recognize that the aging process is also manifested at the cellular level, as well as in mitochondria. Cellular aging is manifested in loss of doubling capacity, increased levels of apoptosis, changes in differentiated phenotype, and changes in metabolism, *e.g.*, decreased levels of protein synthesis and turnover.

Given the programmed nature of cellular and organismal aging, it is possible to evaluate the “biological age” of a cell or organism by means of phenotypic characteristics that are correlated with aging. For example, biological age can be deduced from patterns of gene expression, resistance to stress (*e.g.*, oxidative or genotoxic stress), rate of cellular proliferation,

and the metabolic characteristics of cells (*e.g.*, rates of protein synthesis and turnover, mitochondrial function, ubiquinone biosynthesis, cholesterol biosynthesis, ATP levels within the cell, levels of a Krebs cycle intermediate in the cell, glucose metabolism, nucleic acid metabolism, ribosomal translation rates, etc.). As used herein, “biological age” is a measure of the age of a cell or organism based upon the molecular characteristics of the cell or organism. Biological age is distinct from “temporal age,” which refers to the age of a cell or organism as measured by days, months, and years.

Described below are exemplary methods for identifying compounds that can reduce the rate of aging of an organism and thereby slow or ameliorate the pathologies associated with increased temporal age. Activation of p53 may lead to cell cycle arrest or to apoptosis; Sir2 can suppress this effect by deacetylating p53. Accordingly, the expression or activity of p53 and/or Sir2 gene products in an organism can be a determinant of the rate of aging and life span of the organism. Reduction in the level and/or activity of such gene products would reduce the rate of aging and may ameliorate (at least temporarily) the symptoms of aging. A variety of techniques may be utilized to inhibit the expression, synthesis, or activity of such target genes and/or proteins. Such molecules may include, but are not limited to small organic molecules, peptides, antibodies, antisense, ribozyme molecules, triple helix molecules, and the like.

The following assays provide methods (also referred to herein as “evaluating a compound” or “screening a compound”) for identifying modulators, *i.e.*, candidate or test compounds (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) which modulate Sir2 or p53 activity, *e.g.*, have a stimulatory or inhibitory effect on, for example, Sir2 or p53 expression or activity, or have a stimulatory or inhibitory effect on, for example, the expression or activity of a Sir2 or p53 substrate. Such compounds can be agonists or antagonists of Sir2 or p53 function. These assays may be performed in animals, *e.g.*, mammals, in organs, in cells, in cell extracts, *e.g.*, purified or unpurified nuclear extracts, intracellular extracts, in purified preparations, in cell-free systems, in cell fractions enriched for certain components, *e.g.*, organelles or compounds, or in other systems known in the art. Given the teachings herein and the state of the art, a person of ordinary skill in the art would be able to choose an appropriate system and assay for practicing the methods of the present invention.

Some exemplary screening assays for assessing activity or function include one or more of the following features:

- use of a transgenic cell, *e.g.*, with a transgene encoding Sir2 or p53 or a mutant thereof;
- use of a mammalian cell that expresses Sir2 or p53;
- detection of binding of a labeled compound to Sir2 or a transcription factor where the compound is, for example, a peptide, protein, antibody or small organic molecule; *e.g.*, the compound interferes with or disrupts an interaction between Sir2 and a transcription factor
- use of proximity assays that detect interaction between Sir2 and a transcription factor (*e.g.*, p53), or fragments thereof, for example, fluorescence proximity assays..
- use of a two hybrid assay to detect interaction between Sir2 and a transcription factor (*e.g.*, p53) or fragments thereof. In some instances, the two hybrid assay can be evaluated in the presence of a test compound, *e.g.*, to determine if the test compound disrupts or interferes with an interaction. Two hybrid assays can, for example, be conducted using yeast or bacterial systems.
- use of radio-labelled substrates, *e.g.* ^{35}S , ^3H , ^{14}C , *e.g.*, to determine acetylation status, metabolic status, rate of protein synthesis, *inter alia*.
- use of antibodies specific for certain acetylated or de-acetylated forms of the substrate.

One embodiment herein accordingly comprises methods for the identification of small molecule drug candidates from large libraries of compounds that appear to have therapeutic activity to affect metabolic maintenance and/or to reverse or prevent cell death and thus exhibits potential therapeutic utility, such as the ability to enhance longevity. Small organic molecules and peptides having effective inhibitory activity may be designed *de novo*, identified through assays or screens, or obtained by a combination of the two techniques. Non-protein drug design may be carried out using computer graphic modeling to design non-peptide, organic molecules able to bind to p53 or Sir2. The use of nuclear magnetic resonance (NMR) data for modeling is also known in the art, as described by Lam *et al.*, *Science* 263: 380, 1994, using information from x-ray crystal structure studies of p53 or Sir2, such as that described in Min, J. *et al.*, *Cell* 105:269-279, 2001.

Small molecules may also be developed by generating a library of molecules, selecting for those molecules which act as ligands for a specified target, (using protein functional assays, for example), and identifying the selected ligands. See, *e.g.*, Kohl *et al.*, *Science* 260: 1934,

1993. Techniques for constructing and screening combinatorial libraries of small molecules or oligomeric biomolecules to identify those that specifically bind to a given receptor protein are known. Suitable oligomers include peptides, oligonucleotides, carbohydrates, nonoligonucleotides (*e.g.*, phosphorothioate oligonucleotides; see Chem. and Engineering News, page 20, 7 Feb. 1994) and nonpeptide polymers (see, *e.g.*, "peptoids" of Simon *et al.*, *Proc. Natl. Acad. Sci. USA* 89 9367, 1992). See also U.S. Pat. No. 5,270,170 to Schatz; Scott and Smith, *Science* 249: 386-390, 1990; Devlin *et al.*, *Science* 249: 404-406, 1990; Edgington, *BIO/Technology*, 11: 285, 1993. Libraries may be synthesized in solution on solid supports, or expressed on the surface of bacteriophage viruses (phage display libraries).

Known screening methods may be used by those skilled in the art to screen combinatorial libraries to identify active molecules. For example, an increase (or decrease) in p53 or Sir2 activity due to contact with an agonist or antagonist can be monitored.

In one embodiment, assays for screening candidate or test compounds that are substrates of a Sir2 or p53 protein or polypeptide or biologically active portion thereof are provided. In another embodiment, assays for screening candidate or test compounds which bind to or modulate the activity of a Sir2 or p53 protein or polypeptide or biologically active portion thereof, *e.g.*, modulate the ability of Sir2 or p53 to interact with a ligand, are provided. In still another embodiment, assays for screening candidate or test compounds for the ability to bind to or modulate the activity of a Sir2 or p53 protein or polypeptide and to also alter the rate of aging of a cell or an organism are provided.

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 90: 6909, 1993; Erb. *et al.*, *Proc. Natl. Acad. Sci. USA* 91: 11422, 1994; Zuckermann *et al.*, *J. Med. Chem.* 37: 2678, 1994; Cho *et al.*, *Science* 261: 1303, 1993; Carrell *et al.*, *Angew. Chem. Int. Ed. Engl.* 33: 2059, 1994; Carell *et al.*, *Angew. Chem. Int. Ed. Engl.* 33: 2061, 1994; and in Gallop *et al.*, *J. Med. Chem.* 37:1233, 1994.

Libraries of compounds may be presented in solution (*e.g.*, Houghten, *Biotechniques* 13: 412-421, 1992), or on beads (Lam, *Nature* 354: 82-84, 1991), chips (Fodor, *Nature* 364: 555-556, 1993), bacteria (Ladner U.S. P.N. 5,223,409), spores (Ladner U.S. P.N. '409), plasmids (Cull *et al.*, *Proc Natl Acad Sci USA* 89: 1865-1869, 1992) or on phage (Scott and Smith, *Science* 249:

386-390, 1990); (Devlin, *Science* 249: 404-406, 1990); (Cwirla *et al.*, *Proc. Natl. Acad. Sci U.S.A.* 87: 6378-6382, 1990); (Felici, *J. Mol. Biol.* 222: 301-310, 1991); (Ladner *supra.*).

The compounds tested as modulators of Sir2 or p53 can be any small chemical compound, or a biological entity, such as a protein, *e.g.*, an antibody, a sugar, a nucleic acid, *e.g.*, an antisense oligonucleotide or a ribozyme, or a lipid. Alternatively, modulators can be genetically altered versions of Sir2 or p53. Typically, test compounds will be small chemical molecules and peptides, or antibodies, antisense molecules, or ribozymes. Essentially any chemical compound can be used as a potential modulator or ligand in the assays of the invention, although most often compounds that can be dissolved in aqueous or organic (especially DMSO-based) solutions are used. The assays are designed to screen large chemical libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (*e.g.*, in microtiter formats on microtiter plates in robotic assays). It will be appreciated that there are many suppliers of chemical compounds, including Sigma (St. Louis, MO), Aldrich (St. Louis, MO), Sigma-Aldrich (St. Louis, MO), Fluka Chemika-Biochemica Analytika (Buchs Switzerland) and the like.

In one preferred embodiment, high throughput screening methods known to one of ordinary skill in the art involve providing a combinatorial chemical or peptide library containing a large number of potential therapeutic compounds (potential modulator or ligand compounds). Such “combinatorial chemical libraries” or “ligand libraries” are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional “lead compounds” or can themselves be used as potential or actual therapeutics.

A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining a number of chemical “building blocks” such as reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (*i.e.*, the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks. Moreover, a combinatorial library can

be designed to sample a family of compounds based on a parental compound, *e.g.*, based on the chemical structure of NAD or nicotinamide.

Preparation and screening of combinatorial chemical libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (*see, e.g.*, U.S. Patent 5,010,175, Furka, *Int. J. Pept. Prot. Res.* 37:487-493 (1991) and Houghton *et al.*, *Nature* 354:84-88 (1991)). Other chemistries for generating chemical diversity libraries can also be used. Such chemistries include, but are not limited to: peptoids (*e.g.*, PCT Publication No. WO 91/19735), encoded peptides (*e.g.*, PCT Publication No. WO 93/20242), random bio-oligomers (*e.g.*, PCT Publication No. WO 92/00091), benzodiazepines (*e.g.*, U.S. Pat. No. 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs *et al.*, *Proc. Nat. Acad. Sci. USA* 90:6909-6913 (1993)), vinylogous polypeptides (Hagihara *et al.*, *J. Amer. Chem. Soc.* 114:6568 (1992)), nonpeptidal peptidomimetics with glucose scaffolding (Hirschmann *et al.*, *J. Amer. Chem. Soc.* 114:9217-9218 (1992)), analogous organic syntheses of small compound libraries (Chen *et al.*, *J. Amer. Chem. Soc.* 116:2661 (1994)), oligocarbamates (Cho *et al.*, *Science* 261:1303 (1993)), and/or peptidyl phosphonates (Campbell *et al.*, *J. Org. Chem.* 59:658 (1994)), nucleic acid libraries (*see* Ausubel, Berger and Sambrook, *all supra*), peptide nucleic acid libraries (*see, e.g.*, U.S. Patent 5,539,083), antibody libraries (*see, e.g.*, Vaughn *et al.*, *Nature Biotechnology*, 14(3):309-314 (1996) and PCT/US96/10287), carbohydrate libraries (*see, e.g.*, Liang *et al.*, *Science*, 274:1520-1522 (1996) and U.S. Patent 5,593,853), small organic molecule libraries (*see, e.g.*, benzodiazepines, Baum C&EN, Jan 18, page 33 (1993); isoprenoids, U.S. Patent 5,569,588; thiazolidinones and metathiazanones, U.S. Patent 5,549,974; pyrrolidines, U.S. Patents 5,525,735 and 5,519,134; morpholino compounds, U.S. Patent 5,506,337; benzodiazepines, 5,288,514, and the like).

Devices for the preparation of combinatorial libraries are commercially available (*see, e.g.*, 357 MPS, 390 MPS, Advanced Chem Tech, Louisville KY, Symphony, Rainin, Woburn, MA, 433A Applied Biosystems, Foster City, CA, 9050 Plus, Millipore, Bedford, MA). In addition, numerous combinatorial libraries are themselves commercially available (*see, e.g.*, ComGenex, Princeton, N.J., Asinex, Moscow, Ru, Tripos, Inc., St. Louis, MO, ChemStar, Ltd, Moscow, RU, 3D Pharmaceuticals, Exton, PA, Martek Biosciences, Columbia, MD, etc.).

In one embodiment, the invention provides solid phase based *in vitro* assays in a high throughput format, *e.g.*, where each assay includes a cell or tissue expressing Sir2 and/or p53. In

a high throughput assays, it is possible to screen up to several thousand different modulators or ligands in a single day. In particular, each well of a microtiter plate can be used to run a separate assay against a selected potential modulator, or, if concentration or incubation time effects are to be observed, every 5-10 wells can test a single modulator. Thus, a single standard microtiter plate can assay about 96 modulators. If 1536 well plates are used, then a single plate can easily assay from about 100- about 1500 different compounds. It is possible to assay many plates per day; assay screens for up to about 6,000, 20,000, 50,000, or 100,000 or more different compounds are possible using the integrated systems of the invention.

Candidate Sir2- or p53-interacting molecules encompass many chemical classes. They can be organic molecules, preferably small organic compounds having molecular weights of 50 to 2,500 Daltons. The candidate molecules comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, for example, carbonyl, hydroxyl, and carboxyl groups. The candidate molecules can comprise cyclic carbon or heterocyclic structures and aromatic or polyaromatic structures substituted with the above groups. In one embodiment, the candidate molecules are structurally and/or chemically related to NAD or to nicotinamide.

Other techniques are known in the art for screening synthesized molecules to select those with the desired activity, and for labeling the members of the library so that selected active molecules may be identified, as in U.S. P.N. 5,283,173 to Fields *et al.*, (use of genetically altered *Saccharomyces cerevisiae* to screen peptides for interactions). As used herein, "combinatorial library" refers to collections of diverse oligomeric biomolecules of differing sequence, which can be screened simultaneously for activity as a ligand for a particular target. Combinatorial libraries may also be referred to as "shape libraries", *i.e.*, a population of randomized fragments that are potential ligands. The shape of a molecule refers to those features of a molecule that govern its interactions with other molecules, including Van der Waals, hydrophobic, electrostatic and dynamic.

Nucleic acid molecules may also act as ligands for receptor proteins. See, *e.g.*, Edgington, *BIO/Technology* 11: 285, 1993. U.S. P.N. 5,270,163 to Gold and Tuerk describes a method for identifying nucleic acid ligands for a given target molecule by selecting from a library of RNA molecules with randomized sequences those molecules that bind specifically to the target molecule. A method for the in vitro selection of RNA molecules immunologically cross-reactive with a specific peptide is disclosed in Tsai *et al.*, *Proc. Natl. Acad. Sci. USA* 89:

8864, (1992); and Tsai *et al. Immunology* 150:1137, (1993). In the method, an antiserum raised against a peptide is used to select RNA molecules from a library of RNA molecules; selected RNA molecules and the peptide compete for antibody binding, indicating that the RNA epitope functions as a specific inhibitor of the antibody-antigen interaction.

Antibodies that are both specific for a target gene protein and that interfere with its activity may be used to inhibit target gene function. Such antibodies may be generated using standard techniques, against the proteins themselves or against peptides corresponding to portions of the proteins. Such antibodies include but are not limited to polyclonal, monoclonal, Fab fragments, single chain antibodies, chimeric antibodies, and the like. Where fragments of the antibody are used, the smallest inhibitory fragment which binds to the target protein's binding domain is preferred. For example, peptides having an amino acid sequence corresponding to the domain of the variable region of the antibody that binds to the target gene protein may be used. Such peptides may be synthesized chemically or produced via recombinant DNA technology using methods well known in the art (*e.g.*, see Sambrook *et al.*, Eds., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, (1989), or Ausubel, F. M. *et al.*, eds. *Current Protocols in Molecular Biology* (1994).

Alternatively, single chain neutralizing antibodies that bind to intracellular target gene epitopes may also be administered. Such single chain antibodies may be administered, for example, by expressing nucleotide sequences encoding single-chain antibodies within the target cell population by utilizing, for example, techniques such as those described in Marasco *et al.*, *Proc. Natl. Acad. Sci. USA* 90: 7889-7893 (1993).

Also encompassed are assays for cellular proteins that interact with Sir2 or p53. Any method suitable for detecting protein-protein interactions may be used. The traditional methods that may be used include, for example, co-immunoprecipitation, crosslinking, and co-purification through gradients or chromatographic columns. For these assays, Sir2 or p53 can be a full-length protein or an active fragment. Additional methods include those methods that allow for the simultaneous identification of genes that encode proteins that interact with Sir2 or p53. These methods include, for example, probing expression libraries using a labeled Sir2 or p53 protein, Sir2 or p53 fragment, or Sir2 or p53 fusion protein.

One method to detect protein-protein interaction in vivo is the two-hybrid system, see, for example, Chien *et al.*, *Proc. Natl. Acad. Sci. USA* 88: 9578-9582 (1991). In brief, the two-hybrid

system utilizes plasmids constructed to encode two hybrid proteins: one plasmid comprises the nucleotides encoding the DNA binding domain of a transcriptional activator protein fused to the Sir2 or p53 nucleotide sequence encoding the Sir2 or p53 polypeptide, and the other plasmid comprises the nucleotides encoding the transcriptional activator protein's activation domain fused to a cDNA encoding an unknown protein that has been recombined into the plasmid from a cDNA library. The DNA binding domain fusion plasmid and the cDNA fusion protein library plasmids are transformed into a strain of yeast that contains a reporter gene, for example lacZ, whose regulatory region contains the activator's binding site. Either hybrid protein alone cannot activate translation of the reporter gene because it is lacking either the DNA binding domain or the activator domain. Interaction of the two hybrid proteins, however, reconstitutes a functional activator protein and results in activation of the reporter gene that is detected by an assay for the reporter gene product. The colonies that reconstitute activator activity are purified and the library plasmids responsible for reporter gene activity are isolated and sequenced. The DNA sequence is then used to identify the protein encoded by the library plasmid.

Macromolecules that interact with Sir2 or p53 are referred to as Sir2 or p53 binding partners. Sir2 or p53 binding partners are likely to be involved in the regulation of Sir2 or p53 function. Therefore, it is possible to identify compounds that interfere with the interaction between Sir2 or p53 and its binding partners. The basic principle of assay systems used to identify compounds that interfere with the interaction of Sir2 or p53 and a binding partner is to prepare a reaction mixture containing Sir2 or p53 or a Sir2 or p53 fragment and the binding partner under conditions that allow complex formation. The reaction mixture is prepared in the presence or absence of the test compound to test for inhibitory activity. The test compound may be added prior to or subsequent to Sir2/ or p53/binding partner complex formation. The formation of a complex in a control but not with the test compound confirms that the test compound interferes with complex formation. The assay can be conducted either in the solid phase or in the liquid phase.

In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing Sir2 or p53 with a test compound and determining the ability of the test compound to modulate (*e.g.* stimulate or inhibit) the activity of Sir2 or p53. A preferred activity is the deacetylation function of Sir2 on p53; a further preferred activity is the ability of p53 to cause ERU cycle arrest or apoptosis. Determining the ability of the test compound to modulate the

activity of Sir2 or p53 can be accomplished, for example, by determining the ability of Sir2 or p53 to bind to or interact with the test molecule, or by determining the ability of the test molecule to stimulate or inhibit the activity of Sir2 or p53. Cell-based systems can be used to identify compounds that inhibit Sir2 or p53. Such cells can be recombinant or non-recombinant, such as cell lines that express the Sir2 or p53 gene. Preferred systems are mammalian or yeast cells that express Sir2 or p53. In utilizing such systems, cells are exposed to compounds suspected of ameliorating body weight disorders or increasing lifespan. After exposure, the cells are assayed, for example, for expression of the Sir2 or p53 gene or activity of the Sir2 or p53 protein. Alternatively, the cells are assayed for phenotypes such as those resembling body weight disorders or lifespan extension. The cells may also be assayed for the inhibition of the deacetylation function of Sir2 on p53, or the apoptotic or cytostatic function of p53.

Another preferred cell for a cell-based assay comprises a yeast cell transformed with a vector comprising the Sir2 or p53 gene. One use for a yeast cell expressing Sir2 or p53 is to mutagenize the yeast and screen for yeast that will survive only when the Sir2 or p53 polypeptide is functioning normally. Synthetic lethal screens are described in Holtzman *et al.* (1993), *J. Cell Bio.* 122: 635-644. The yeast that require Sir2 or p53 function for survival can then be used to screen test compounds for those that inhibit Sir2 or p53 activity. Test compounds that results in a decrease in yeast survival are likely inhibitors of Sir2 or p53 in this system.

In yet another embodiment, an assay is a cell-free assay in which Sir2 or p53 protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to the Sir2 or p53 protein or biologically active portion thereof is determined. Binding of the test compound to the Sir2 or p53 protein can be determined either directly or indirectly as described above. In a preferred embodiment, the assay includes contacting the Sir2 or p53 protein or biologically active portion thereof with a known compound which binds Sir2 or p53 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an Sir2 or p53 protein, wherein determining the ability of the test compound to interact with an Sir2 or p53 protein comprises determining the ability of the test compound to preferentially bind to Sir2 or p53 or a biologically active portion thereof as compared to the known compound.

In yet another embodiment, an assay is a cell-free system in which Sir2 protein or biologically active portion thereof is contacted with p53 protein or biologically active portion

thereof, to form a mixture comprising a detectable amount bound p53:Sir complex. And a test compound is contacted with the mixture, and the ability of the compound to effect the stability or formation of the p53:Sir2 complex is determined. Interaction of the test compound with the p53:Sir2 complex may be determined directly or by methods known in the art. In a preferred embodiment, the method comprises contacting p53 with Sir2 to form a mixture comprising the p53:Sir2 complex, further contacting the mixture with a compound to be tested, and evaluating the binding kinetics of p53:Sir2 complex both in the presence and the absence of the test compound to directly bind the p53:Sir2 complex is evaluated. The cell-free assays are amenable to use of both soluble and/or membrane-bound forms of proteins. In the case of cell-free assays in which a membrane-bound form of a protein is used it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton X-100, Triton X-114, Thesit, Isotridecypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylammonio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylammonio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl,N,N-dimethyl-3-amino-1-propane sulfonate.

In more than one embodiment of the above assay methods, it may be desirable to immobilize either Sir2 or p53 or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to an Sir2 or p53 protein, or interaction of an Sir2 or p53 protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase/Sir2 or p53 fusion proteins or glutathione-S-transferase/target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or Sir2 or p53 protein, and the mixture incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter

plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of Sir2 or p53 binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either a Sir2 or p53 protein or a Sir2 or p53 target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated Sir2 or p53 protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with Sir2 or p53 protein or target molecules but which do not interfere with binding of the Sir2 or p53 protein to its target molecule can be derivatized to the wells of the plate, and unbound target Sir2 or p53 protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the Sir2 or p53 protein or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the Sir2 or p53 protein or target molecule.

In addition to cell-based and *in vitro* assay systems, non-human organisms, *e.g.*, transgenic non-human organisms, can also be used. A transgenic organism is one in which a heterologous DNA sequence is chromosomally integrated into the germ cells of the animal. A transgenic organism will also have the transgene integrated into the chromosomes of its somatic cells. Organisms of any species, including, but not limited to: yeast, worms, flies, fish, reptiles, birds, mammals (*e.g.*, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, and goats), and non-human primates (*e.g.*, baboons, monkeys, chimpanzees) may be used in the methods of the invention.

Accordingly, in another embodiment, the invention features a method of identifying a compound that alters the rate of aging of a cell or an organism, comprising: contacting a Sir2 or p53 polypeptide with a test compound; evaluating an interaction between the test compound and the Sir2 or p53 polypeptide; and further evaluating the effect of the test compound on the rate of aging of a cell or organism.

The interaction between a test compound and the Sir2 or p53 polypeptide can be performed by any of the methods described herein, *e.g.*, using cell-based assays or cell-free *in vitro* assays. Whether the interaction between the test compound and the Sir2 or p53 polypeptide is evaluated prior to the evaluation of the effect of the test compound on the rate of aging of a cell or organism is not critical to the method. However, it is preferable to evaluate the interaction between the test compound and Sir2 or p53 polypeptide first, so that test compounds that do not interact with the Sir2 or p53 polypeptide do not have to be tested for their effect upon the rate of aging. It can also be preferable to use an assay for evaluating the interaction between the test compound and the Sir2 or p53 polypeptide that can be adapted for high throughput screening, thus making it possible to screen one or more libraries of test compounds. Possible test compounds include, *e.g.*, small organic molecules, peptides, antibodies, and nucleic acid molecules, as described above.

The rate of aging of an organism can be determined using methods known in the art. For example, the rate of aging of an organism can be determined by directly measuring the life span of the organism. Preferably, a statistical measure, *e.g.*, an average or median value, of the life span of a group of animals, *e.g.*, a group of genetically matched animals, will be determined and the resulting statistical value compared to an equivalent statistical value, *e.g.*, an average or median value, of the life span of a control group of animals, *e.g.*, a group of animals that did not receive the test compound but are genetically matched to the group of animals that did receive the test compound. Such methods are suitable for organisms that have a short life span, such as worms or flies. See, for example, Rogina *et al.* (2000), *Science* 290:2137-40. Direct measurement of life span can also be performed with other organisms such as rodents, as discussed, for example, in Weindruch *et al.* (1986), *Journal of Nutrition* 116(4):641-54. Those skilled in the art will recognize that there are many ways of measuring the statistical difference (*e.g.*, using the Student's T test) between two sets of data, any of which may be suitable for the methods of the invention.

To reduce the time that it takes to measure a change in the rate of aging using data on the life span of the organisms treated with the test compound, various modifications or treatments of the organisms can be implemented. For example, animals fed on a calorically rich diet tend to live shorter lives, thus reducing the time that needs to elapse to determine when the average life span of the test group of animals has exceeded the average life span of the control group of

animals. Alternatively, the test compound can be administered to test animals that have already lived for 50%, 60%, 70%, 80%, 90%, or more of their expected life span. Thus, the test compound can be administered to an adult organism, or even an old adult organism. Other possibilities include the use of genetically modified organisms. For example, the organisms could harbor mutations (e.g., a *Hyperkinetic*¹ or *Shaker*⁵ mutation in *Drosophila*, or a mutation in a silent information regulator gene (e.g., *Sir2*), or a catalase or superoxide dismutase gene) or transgenes (e.g., encoding a transporter protein (e.g., a carboxylate transport protein such as INDY) or a protein involved in insulin signaling and metabolic regulation (e.g., IGF-1)) that reduce their average life span. See Rogina *et al.* (1997), *Proc. Natl. Acad. Sci., USA* 94:6303-6; Rogina and Helfand (2000), *Biogerontology* 1:163-9; and Guarente and Kenyon (2000), *Nature* 408:255-62. Those skilled in the art will understand that it may also be desirable to practice the methods of the invention using organisms that are long-lived, such as calorically restricted animals, or animals carrying mutations or transgenes that increase their life span.

A proxy for rate of aging of a cell or an organism can be determined using biomarkers that are indicative of the biological age of the organism (*i.e.*, age-related parameters). Using biomarkers for determining biological age can greatly facilitate screens for compounds that alter the rate of aging, as they bypass the requirement of waiting for the animal to die in order to determine the rate of aging. Biomarkers suitable for use in the present invention include, but are not limited to, levels of protein modification, e.g., accumulation of glycosylated proteins, rates or levels of protein turnover, levels or composition of T-cell populations, protein activity, physical characteristics, macular degeneration, and/or increased copper and zinc concentrations in neuronal tissues. The expression of genes whose regulation is biological age-dependent is a particularly preferred biomarker for use in the methods of the invention. Numerous genes are known to be expressed in a biological age-dependent manner. In *Drosophila*, for example, such genes include *wingless* and *engrailed*. See Rogina and Helfand (1997), *Mechanisms of Development* 63:89-97. In mice, the expression of the *ras* oncogene is elevated in older animals. See Hass *et al.* (1993), *Mutat. Res.* 295(4-6):281-9. Similarly, in rodents and worms, genes that are differentially expressed in young and old organisms have been identified by transcriptional profiling using microarrays. See, e.g., Lee *et al.* (1999), *Science* 285:1390-93; WO 01/12851; and Hill *et al.* (2000), *Science* 290:809-812. For example, Hill *et al.* (2000) *Science* 90:809 discloses genes whose transcripts are up-regulated in nematodes that are at 2 weeks in

development. Examples of such genes include the genes described in cluster (4,1):69 of Hill, *supra*. Any gene whose regulation is biological age-dependent is suitable for the methods of the invention. Preferably, more than one gene is analyzed so as to improve the accuracy of the determination. Analysis of gene expression can be performed by any technique known in the art, including Northern, in-situ hybridization, quantitative PCR, and transcriptional profiling using microarrays. Methods of determining biological age based on gene expression patterns are described in WO 01/12851.

Metabolic parameters can also be used to evaluate the rate of aging of a cell or organism. For example, the rate of protein synthesis and degradation decreases in biologically aged cells, and the levels proteins having advanced glycosylation end product modifications increases. See, Lambert and Merry (2000), *Exp. Gerontol* 35(5):583-94; and WO 01/79842. In addition, animals that harbor mutations conferring longer life span (and thus a reduced rate of aging) can show defects in ubiquinone biosynthesis, mitochondrial biogenesis, glucose metabolism, nucleic acid metabolism, ribosomal translation rates, and cholesterol biosynthesis. See, for example, WO 98/17823 and WO 99/10482. Thus, by measuring any of these parameters or some combination thereof, it is possible to indirectly evaluate the rate of aging of a cell or an organism. Methods of analyzing protein synthesis, degradation, and modification with advanced glycosylation end products are known in the art, as described in Lambert and Merry (2000), *Exp. Gerontol* 35(5):583-94 and WO 01/79842. Similarly, methods of analyzing ubiquinone biosynthesis, mitochondrial biogenesis, and glucose metabolism are known in the art (see, *e.g.*, Marbois *et al. J. Biol. Chem.* 271:2995; Proft *et al. EMBO J.* 14:6116; and WO 98/17823), as are methods of analyzing nucleic acid metabolism, ribosomal translation rates, and cholesterol biosynthesis (see, *e.g.*, WO 99/10482).

Cellular proliferation is another parameter that can be used to evaluate the biological age of a cell or organism. Cells from biologically aged organisms demonstrate reduced proliferative capacity as compared to the cells of a corresponding younger organism. See Li *et al.* (1997), *Invest. Ophthalmol.* 38(1):100-7; and Wolf and Pendergrass (1999), *J Gerontol. A Biol. Sci. Med. Sci.* 54(11):B502-17. It will be understood by one skilled in the art that there are many methods for evaluating the proliferative capacity of cells that are suitable for use in the methods of the invention. For example, cells can be labeled in vitro (or in vivo) with BrdU to determine the percent of dividing cells or evaluated using a colony forming assay, as described in Li *et al.*

(1997), *supra*. Cells suitable for the analysis of proliferative capacity include cells grown in tissue culture, cells isolated from an animal that has been treated with a test compound, cells that are part of a live animal, or cells that are part of a tissue section obtained from an animal. With respect to cells present in an animal or tissue section thereof, preferable cells include lens epithelial cells, osteoblasts, osteoclasts, and lymphoid cells.

Basically, any biomarker that is altered in a biological age-dependent manner has the potential to be used to evaluate the effect of a test compound upon the rate of aging of a cell or an organism. Thus, additional biomarkers include visual appearance, resistance to oxidative stress, cellular transformation (the ability to adopt a transformed (*i.e.*, cancerous or malignant) phenotype), or DNA methylation (*e.g.*, of a ras oncogene). See, for example, Finkel and Holbrook (2000), *Nature* 408:239-47; Kari *et al.* (1999), *J Nutr. Health Aging* 3(2):92-101; and Hass *et al.* (1993), *Mutat. Res.* 295(4-6):281-9.

A cell used in the methods of the invention can be from a stable cell line or a primary culture obtained from an organism, *e.g.*, a organism treated with the test compound.

A transgenic cell or animal used in the methods of the invention can include a transgene that encodes, *e.g.*, a copy of a Sir2 or p53 protein, *e.g.*, the Sir2 or p53 polypeptide that was evaluated for an interaction with the test compound. The transgene can encode a protein that is normally exogenous to the transgenic cell or animal, including a human protein, *e.g.*, a human Sir2 or p53 polypeptide. The transgene can be linked to a heterologous or a native promoter.

Transgenic Organisms

This disclosure further relates to a method of producing transgenic animals, *e.g.*, mice or flies. In one embodiment, the transgenic animal is engineered to express, overexpress or ectopically express Sir2 or p53, which method comprises the introduction of several copies of a segment comprising at least the polynucleotide sequence encoding SEQ ID NO. 2 with a suitable promoter into the cells of an embryo at an early stage. Techniques known in the art may be used to introduce the Sir2 or p53 transgene into animals to produce the founder line of animals. Such techniques include, but are not limited to: pronuclear microinjection (U.S. P.N. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten *et al.*, *Proc. Natl. Acad. Sci. USA* 82: 6148-6152, 1985; gene targeting in embryonic stem cells (Thompson *et al.*, *Cell* 56: 313-321, 1989; electroporation of embryos (Lo, *Mol. Cell Biol.* 3: 1803-1814, 1983; and sperm-

mediated gene transfer (Lavitano, *et al.*, *Cell* 57: 717-723, 1989; etc. For a review of such techniques, see Gordon, *Intl. Rev. Cytol.* 115: 171-229, 1989.

Gene targeting by homologous recombination in embryonic stem cells to produce a transgenic animal with a mutation in the Sir2 or p53 gene ("knock-out" mutation) can also be performed. In such so-called "knock-out" animals, there is inactivation of the Sir2 or p53 gene or altered gene expression, such that the animals can be useful to study the function of the Sir2 or p53 gene, thus providing animals models of human disease, which are otherwise not readily available through spontaneous, chemical or irradiation mutagenesis.

A particularly useful transgenic animal is one in which the Sir2 or p53 homolog has been disrupted or knocked out.

Transgenic animals such as mice, for example, may be used as test substrates for the identification of drugs, pharmaceuticals, therapies and interventions that can be used for the ameliorating or slowing the effects of aging.

Accordingly, the invention features a transgenic organism that contains a transgene encoding a Sir2 or p53 polypeptide. In preferred embodiments, the Sir2 or p53 polypeptide is a human Sir2 or p53 polypeptide. The Sir2 or p53 polypeptide can be exogenous to (*i.e.*, not naturally present in) the transgenic organism.

The transgenic organism can be a yeast cell, an insect, *e.g.*, a worm or a fly, a fish, a reptile, a bird, or a mammal, *e.g.*, a rodent.

The transgenic organism can further comprise a genetic alteration, *e.g.*, a point mutation, insertion, or deficiency, in an endogenous gene. The endogenous gene harboring the genetic alteration can be a gene involved in the regulation of life span, *e.g.*, a gene in the insulin signaling pathway, a gene encoding a Sir2 or transcription factor protein, or both. In cases where the genetically altered gene is a Sir2 or transcription factor, *e.g.*, p53, polypeptide, it is preferable that the expression or activity of the endogenous Sir2 or transcription factor, *e.g.*, p53, protein is reduced or eliminated.

Therapeutic Uses

In another embodiment, the invention features a method of altering the expression or activity of a Sir2 or p53 polypeptide, comprising administering to a cell or an organism a compound that increases or decreases the expression or activity of the Sir2 or p53 polypeptide in an amount effective to increase or decrease the activity of the Sir2 or p53 polypeptide.

The Sir2 or p53 polypeptide can also be a yeast, invertebrate (*e.g.*, worm or fly), or vertebrate (*e.g.*, fish, reptile, bird, or mammal (*e.g.*, mouse)) protein.

The cell to which the compound is administered can be an invertebrate cell, *e.g.*, a worm cell or a fly cell, or a vertebrate cell, *e.g.*, a fish cell (*e.g.*, zebrafish cell), a bird cell (*e.g.*, chicken cell), a reptile cell (*e.g.*, amphibian cell, *e.g.*, *Xenopus* cell), or a mammalian cell (*e.g.*, mouse or human cell). Similarly, the organism to which the compound is administered can be an invertebrate, *e.g.*, a worm or a fly, or a vertebrate, *e.g.*, a fish (*e.g.*, zebrafish), a bird (*e.g.*, chicken), a reptile (*e.g.*, amphibian, *e.g.*, *Xenopus*), or a mammal (*e.g.*, rodent or a human). When the organism is a human, it is preferred that the human is not obese or diabetic.

The compound that is administered to the cell or organism can be an agonist that increases the expression or activity of the Sir2 or p53 polypeptide or an antagonist that decreases the expression or activity of the Sir2 or p53 polypeptide. Whether agonist or antagonist, the compound can be a small organic compound, an antibody, a polypeptide, or a nucleic acid molecule.

The agonist or antagonist can alter the concentration of metabolites, *e.g.*, Krebs Cycle intermediates, *e.g.*, succinate, citrate, or α -keto-glutarate, within the cell or within one or more cells of the organism. Such action is expected to alter the cell's or the organism's resistance to oxidative stress. For example, an antagonist could increase the cell's or the organism's resistance to oxidative stress. In addition, the agonist or antagonist can alter one or more aging-related parameters, *e.g.*, the expression of one or more genes or proteins (*e.g.*, genes or proteins that have an age-related expression pattern), or the value of one or more metabolic parameters (*e.g.*, one or more metabolic parameters that reflect the rate of aging of the cell or organism). , the agonist or antagonist alters the rate of aging of the cell or organism.

Ideally, the compound reduces, *e.g.*, partially reduces, the expression of the Sir2 or p53 polypeptide. For example, anti-sense RNA, or ribozymes can be used to reduce the expression of the Sir2 or p53 polypeptide. Double-stranded inhibitory RNA is particularly useful as it can be used to selectively reduce the expression of one allele of a gene and not the other, thereby achieving an approximate 50% reduction in the expression of the Sir2 or p53 polypeptide. See Garrus *et al.* (2001), *Cell* 107(1):55-65.

In one embodiment, treatment of aging comprises modulating the expression of a Sir2 or p53 polypeptide. A cell or subject can be treated with a compound that modulates the expression

of a Sir2 or p53 gene. These compounds can be nucleic acid molecules substantially complementary to a Sir2 or p53 gene. Such approaches include oligonucleotide-based therapies such as antisense, ribozymes, and triple helices .

Oligonucleotides may be designed to reduce or inhibit mutant target gene activity. Techniques for the production and use of such molecules are well known to those of ordinary skill in the art. Antisense RNA and DNA molecules act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. With respect to antisense DNA, oligodeoxyribonucleotides derived from the translation initiation site, *e.g.*, between the -10 and +10 regions of the target gene nucleotide sequence of interest, are preferred. Antisense oligonucleotides are preferably 10 to 50 nucleotides in length, and more preferably 15 to 30 nucleotides in length. An antisense compound is an antisense molecule corresponding to the entire Sir2 or p53 mRNA or a fragment thereof.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage. The composition of ribozyme molecules includes one or more sequences complementary to the target gene mRNA, and includes the well known catalytic sequence responsible for mRNA cleavage disclosed, for example, in U.S. P.N. 5,093,246. Within the scope of this disclosure are engineered hammerhead motif ribozyme molecules that specifically and efficiently catalyze endonucleolytic cleavage of RNA sequences encoding target gene proteins. Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the molecule of interest for ribozyme cleavage sites that include the sequences GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for predicted structural features, such as secondary structure, that may render the oligonucleotide sequence unsuitable. The suitability of candidate sequences may also be evaluated by testing their accessibility to hybridization with complementary oligonucleotides, using ribonuclease protection assays.

Nucleic acid molecules used in triple helix formation for the inhibition of transcription should be single stranded and composed of deoxyribonucleotides. The base composition of these oligonucleotides are designed to promote triple helix formation via Hoogsteen base pairing rules,

which generally require sizeable stretches of either purines or pyrimidines to be present on one strand of a duplex. Nucleotide sequences may be pyrimidine-based, which will result in TAT and CGC triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich molecules provide base complementarity to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, nucleic acid molecules may be chosen that are purine-rich, for example, containing a stretch of G residues. These molecules will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.

Alternatively, the potential sequences targeted for triple helix formation may be increased by creating a "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

The antisense, ribozyme, and/or triple helix molecules described herein may reduce or inhibit the transcription (triple helix) and/or translation (antisense, ribozyme) of mRNA produced by both normal and mutant target gene alleles. If it is desired to retain substantially normal levels of target gene activity, nucleic acid molecules that encode and express target gene polypeptides exhibiting normal activity may be introduced into cells via gene therapy methods that do not contain sequences susceptible to whatever antisense, ribozyme, or triple helix treatments are being utilized. Alternatively, it may be preferable to coadminister normal target gene protein into the cell or tissue in order to maintain the requisite level of cellular or tissue target gene activity.

Antisense RNA and DNA, ribozyme, and triple helix molecules may be prepared by any method known in the art for the synthesis of DNA and RNA molecules. These include techniques for chemically synthesizing oligodeoxyribonucleotides and oligoribonucleotides, for example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety of vectors that incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters.

Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines. Various well-known modifications to the DNA molecules may be introduced as a means of increasing intracellular stability and half-life. Possible modifications include but are not limited to the addition of flanking sequences of ribonucleotides or deoxyribonucleotides of the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the oligodeoxyribonucleotide backbone.

Modulators of Sir2 or p53 expression can be identified by a method wherein a cell is contacted with a candidate compound and the expression of Sir2 or p53 mRNA or protein in the cell is determined. The level of expression of Sir2 or p53 mRNA or protein in the presence of the candidate compound is compared to the level of expression of mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of Sir2 or p53 expression based on this comparison. For example, when expression of Sir2 or p53 mRNA or protein is greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of Sir2 or p53 mRNA or protein expression. Alternatively, when expression of Sir2 or p53 mRNA or protein is less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of Sir2 or p53 mRNA or protein expression. The level of Sir2 or p53 mRNA or protein expression in the cells can be determined by methods described herein for detecting Sir2 or p53 mRNA or protein.

Delivery of antisense, triplex agents, ribozymes, and the like can be achieved using a recombinant expression vector such as a chimeric virus or a colloidal dispersion system or by injection. Useful virus vectors include adenovirus, herpes virus, vaccinia, and/or RNA virus such as a retrovirus. The retrovirus can be a derivative of a murine or avian retrovirus such as Moloney murine leukemia virus or Rous sarcoma virus. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. The specific nucleotide sequences that can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector containing an antisense oligonucleotide can be determined by one of skill in the art.

Another delivery system for polynucleotides is a colloidal dispersion system. Colloidal dispersion systems include macromolecular complexes, nanocapsules, microspheres, beads, and

lipid-based systems including oil-in-water emulsions, micelles, mixed micelles and liposomes. A preferred colloidal delivery system is a liposome, an artificial membrane vesicle useful as in vivo or in vitro delivery vehicles. The composition of a liposome is usually a combination of phospholipids, usually in combination with steroids, particularly cholesterol.

The Sir2 or p53 gene may also be underexpressed.

Methods whereby the level of Sir2 or p53 gene activity may be increased to levels wherein disease symptoms are ameliorated also include increasing the level of gene activity, for example by either increasing the level of Sir2 or p53 gene present or by increasing the level of gene product which is present.

For example, a target gene protein, at a level sufficient to ameliorate metabolic imbalance symptoms, may be administered to a patient exhibiting such symptoms. One of skill in the art will readily know how to determine the concentration of effective, non-toxic doses of the normal target gene protein. Additionally, RNA sequences encoding target gene protein may be directly administered to a patient exhibiting disease symptoms, at a concentration sufficient to produce a level of target gene protein such that the disease symptoms are ameliorated. Administration may be by a method effective to achieve intracellular administration of compounds, such as, for example, liposome administration. The RNA molecules may be produced, for example, by recombinant techniques such as those described above.

Further, patients may be treated by gene replacement therapy. One or more copies of a normal target gene, or a portion of the gene that directs the production of a normal target gene protein with target gene function, may be inserted into cells using vectors that include, but are not limited to adenovirus, adenoma-associated virus, and retrovirus vectors, in addition to other particles that introduce DNA into cells, such as liposomes. Additionally, techniques such as those described above may be utilized for the introduction of normal target gene sequences into human cells.

Cells, preferably autologous cells, containing and expressing normal target gene sequences may then be introduced or reintroduced into the patient at positions which allow for the amelioration of metabolic disease symptoms. Such cell replacement techniques may be preferred, for example, when the target gene product is a secreted, extracellular gene product.

In instances where the target gene protein is extracellular, or is a transmembrane protein, any of the administration techniques described, below which are appropriate for peptide

administration may be utilized to effectively administer inhibitory target gene antibodies to their site of action.

The identified compounds that inhibit target gene expression, synthesis and/or activity can be administered to a patient at therapeutically effective doses to treat or ameliorate or delay the symptoms of aging. A therapeutically effective dose refers to that amount of the compound sufficient to result in amelioration or delay of symptoms of aging.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50 / ED50. Compounds that exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects. The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

Pharmaceutical compositions may be formulated in conventional manner using one or more physiologically acceptable carriers or excipients. Thus, the compounds and their physiologically acceptable salts and solvates may be formulated for administration by inhalation or insufflation (either through the mouth or the nose) or oral, buccal, parenteral or rectal administration.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.*, pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (*e.g.*, lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (*e.g.*, magnesium stearate, talc or silica); disintegrants (*e.g.*, potato starch or sodium starch glycolate); or wetting agents (*e.g.*, sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups, or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (*e.g.*, sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (*e.g.*, lecithin or acacia); non-aqueous vehicles (*e.g.*, almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (*e.g.*, methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavoring, coloring, and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound. For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner. For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.* gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch. The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing, and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-

free water, before use. The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides. In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

All references cited herein are incorporated by reference in their entirety. The invention is illustrated by the following non-limiting examples.

Materials and Methods

Plasmids and antibodies

To construct mSir2 α expression constructs, the full-length cDNA was subcloned from pET28a-Sir2 α (Imai *et al.*, 2000) into pcDNA3 or pBabepuro vector. Site-directed mutation was generated in the plasmid pRS305-Sir2 α using the Gene Edit system (Promega). To construct the human SIRT1 expression construct, DNA sequences corresponding to the full-length hSIRT1 (Frye, 1999) were amplified by PCR from Marathon-Ready Hela cDNA (Clontech), and initially subcloned into pcDNA3.1/V5-His-Topo vector (Invitrogen), and then subcloned with a Flag-tag into a pCIN4 vector for expression (Gu *et al.*, 1999). To prepare the Sir2 α antibody that can recognize both human and mouse Sir2 α , a polyclonal antibody against the highly conserved C-terminus of Sir2 α was generated. DNA sequences corresponding to this region (480-737) were amplified by PCR and subcloned into pGEX-2T (Pharmacia). α -Sir2 α antisera was raised in rabbits against the purified GST-Sir2 α (480-737) fusion protein (Covance), and further affinity-purified on both protein-A and antigen columns. By Western blot analysis and immunofluorescent staining, this antibody can detect both mouse Sir2 α and human SIRT1 proteins.

To construct hSir2 expression constructs, BamHI/SnaBI fragment of hSIR2SIRT1 cDNA was inserted into pBabe-Y-Puro. The resulting plasmid was designated pYESir2-puro. Similarly a BamHI/SnaBI fragment of hSir2 that was mutated at residue 363 from Histidine (H) to Tyrosine (Y) by site-directed mutagenesis (Stratagene) was used to create the retroviral vector pYESir2HY. pBabe-hTERT-hygro contained an EcoRI/SalI fragment of hTERT cloned into

EcoRI/SalI site of pBabe-Hygro. pCMVwtp53, pCMVK382R and pCMVK320R were a gift from Dr. E. Appella (NIH).

Cell culture and derivation of cell lines

All cells were grown in presence of 20% O₂ and 5% CO₂ at 37°C in humidified chambers. Human diploid fibroblast BJ cells, human epithelial breast carcinoma cell line MCF7 and H1299 human epithelial carcinoma cell lines were grown in DME +10% FCS. PBS(-/-) (phosphate buffered saline) without magnesium or calcium was used for washing cells and other applications described herein.

Amphotrophic viruses were produced by transient co-transfection of pCL-pCL-Ampho with the LTR containing pBabe vectors (Morgenstern and Land, 1990), pYESir2 or pYESir2HY in to 293T cell line using Eugene6 (Roche). Three days post transfection supernatants were collected and filtered with 0.4 micron filters. Primary BJ cells or MCF7 cells were infected with retrovirus containing media in presence of 8 mg/ml of polybrene overnight and 48 hours later cells were selected in puromycin at 1 mg/ml.

Following selection and during the experimentation all the mass cultures were maintained in presence of puromycin. These selected BJ cells were subsequently infected and selected with a pBabe-hTERT virus carrying the hygromycin resistance gene (200mg/ml). The resulting cells were: BJT (carrying pYE-Puro backbone and pBabe-hTERT-hygro), BJThSir2wt (carrying pYESir2 wild type hSir2 and pBabe-hTERT hygro) and BJThSir2HY (pYESir2HY mutant hSir2 and pBabe-hTERT-hygro). MCF7 cells were transfected with the vector p21P-luc (Vaziri *et al.*, 1997) and pCMVneo, clones were selected in 500 mg/ml of G418 and the clone designated MCF73L was selected that was able to upregulate the p21WAF1 promoter-luciferase in response to treatment with 6 Gy of ionizing radiation. MCF7 cells or MCF73L were infected with the same viruses as described before to yield the following cell lines: MCF73LP (carrying pBabe Y-puro backbone), MCF73L-hSir2wt and MCF73L-hSir2HY. Cells were kept under appropriate selection throughout experiments.

***In vitro* p53 deacetylation Assay**

The Flag-tagged Sir2 α -expressing cells were established and expanded in DMEM medium, and cell extracts were prepared essentially as previously described (Luo *et al.*, 2000; Gu *et al.*, 1999; Ito *et al.*, 1999). The proteins were purified under a very high stringency

condition (300 mM NaCl and 0.5 % NP-40). The eluted proteins were resolved by a SDS-PAGE gel and analyzed by colloidal blue staining (Novex). Acetylated GST-p53 was prepared by p53 acetylation assay as previously described (Gu and Roeder, 1997) and further purified on glutathione-Sepharose (Luo *et al.*, 2000). The ^{14}C -labeled acetylated p53 (2.5 μg) was incubated with purified Sir2 α (10 ng) at 30 °C for 1 hr either in the presence of 50 μM NAD or as indicated. The reactions were performed in a buffer containing 50 mM Tris-HCl (pH 9.0), 50mM NaCl, 4 mM MgCl_2 , 0.5 mM DTT, 0.2 mM PMSF, 0.02% NP-40 and 5% glycerol. The reactions were resolved on SDS-PAGE and analyzed by Coomassie blue staining and autoradiography.

Immunoprecipitation and Immunofluorescence

H1299 cells transiently expressing p53 and hSir2 were lysed using the NP40 buffer and lysates described above and immunoprecipitated with 1ul of anti-hSir2 antibody overnight. Protein G-sepharose beads (50 ml) were added to the lysates and rotated at 4°C for 3hrs. The immune complexes were collected, washed 3 times, and resolved using the Nupage gradient 4-12% Bis-Tris MOPS (3-N-morpholino propane sulfonic acid) protein gel (Novex) in the presence of provided anti-oxidant (Novex).

The gels used were transferred to nitrocellulose and probed with anti-p53 antibody (pAb7 sheep anti human polyclonal antibody, Oncogene Science), signal detected using a goat anti-sheep HRP secondary antibody. The membranes were subsequently washed and reprobed with anti-hSir2 antibody.

For immunoprecipitation in BJ cells, 1mg of protein per reaction were incubated with 1ul of Ab-6(anit-p53 monoclonal, Oncogene Science) and immunoprecipitation was performed as described above except that the time of incubation in primary antibody was 2hrs and 4 times higher concentrations of protease inhibitors were used, due to the observed high instability of p53 protein in BJ cells. Immune complexes were resolved as previously described using the Novex system (Invitrogen) and membranes were exposed to a mix of polyclonal antibodies at 1:1000 dilution (SC6432, polyclonal rabbit and CM1, polyclonal rabbit). A secondary goat anti-rabbit HRP was used at 1:30,000 concentration for detection. Membranes were subsequently blocked again and re-probed with anti-hSir2 antibody.

Immunofluorescence of U20S and BJ cells was undertaken by fixing the cells in microchamber slides (LabTek) in 70% Ethanol and subsequent staining with anti-hSir2 antibody at 1:500 dilution. A secondary goat anti-rabbit FITC antibody at 0.5 ug/ml was used for detection of signal.

GST pull-down assay and co-immunoprecipitation assay

GST fusion proteins were expressed in *E. coli*, extracted with buffer BC500 (20 mM Tris-HCl, pH 8.0, 0.5 mM EDTA, 20% glycerol, 1mM DTT and 0.5 mM PMSF) containing 50 mM KCl and 1% NP-40, and purified on glutathione-sepharose (Pharmacia). ³⁵S-labeled Sir2 α was *in vitro* translated by a TNT kit (Promega) using pcDNA3-Sir2 α as a template. 5 μ l of ³⁵S-labeled Sir2 α were incubated at 4°C for 60 min with each of the different immobilized GST fusion proteins in BC200 buffer containing 200 mM KCl and 0.2% NP-40. Beads were then washed five times in 0.5 ml of the same buffer. Bound proteins were eluted with an equal volume of SDS sample buffer, resolved by SDS-PAGE, and analyzed by Coomassie blue staining and autoradiography.

The co-immunoprecipitation assay was performed essentially as described previously (Luo *et al.*, 2000). Cells were extracted with lysis buffer (25mM HEPES-KOH, pH 8.0, 150 mM KCl, 2mM EDTA, 1mM DTT, 1mM PMSF, 10 μ g/ml aprotinin, 10 μ g/ml leupeptin, 1 μ g/ml pepstatin A, 20 mM NaF, 0.1% NP-40). After centrifugation, the supernatants were incubated with M2 beads (Sigma) for 4 hr at 4°C. The M2 beads were washed five times with 0.5 ml lysis buffer, after which the associated proteins were eluted with Flag peptides to avoid the cross-reaction from the mouse IgG in western blot analysis. In the case of the co-immunoprecipitation in normal cells, 50 million cells were extracted in the same lysis buffer. The supernatants were incubated with 20 μ g α -Sir2 α antibody or pre-immune antiserum from the same rabbit and 40 μ l protein A/G plus-agarose (Santa Cruz) for overnight. The agarose beads were washed five times with 0.5 ml of lysis buffer, after which the associated proteins were eluted with BC1000 (20 mM Tris-HCl, pH 8.0, 0.5 mM EDTA, 20% glycerol, 1mM DTT and 0.5 mM PMSF) containing 1 M NaCl, 1% NP-40, 0.5% Deoxycholic Acid. The eluted proteins were resolved on 8% SDS PAGE and Western blot with α -Sir2 α antibody and α -p53 (DO-1) for human cells and α -p53 (421) for mouse cell.

Immunoblot analysis

For detection of acetylated forms of p53 in BJ cells and MCF7 cells, equal numbers of cells were plated 24 hrs before the experiment. 1.5×10^6 BJ cells or 10^7 MCF7 cells exponentially growing phase in 150 cm² dishes were exposed to 6Gy of ionizing radiation (137Cesium gamma source at dose rate of 1 Gy/min). At the appropriate time point, cells were washed and harvested by trypsinization and subsequent neutralization with 10% serum. After washing the cells once in PBS(-/-), cell pellets were frozen on dry ice instantly at the appropriate time point. Once all time points were collected, cell pellets were all lysed on ice at once by adding 0.5% NP40, 150 mM NaCl (in the presence of complete protease inhibitor mix, Roche), for 30 minutes and vortexing. Cell lysates were prepared by centrifugation for 20 minutes at 4°C. Protein content of lysates were measured using Lowry based assay (BioRad DC protein assay). Protein (300 mg) was resolved on gradient 4-20% criterion Tris-HC gels (Biorad), transferred to nitrocellulose and blocked in 10% skim milk.

The resulting membrane was incubated overnight in 1:400 dilution of Ab-1 (Oncogene Science, peptide based rabbit polyclonal anti K382 p53). This membrane was then washed twice in PBS(-/-) containing 0.05% Tween 20 for 15 minutes. Secondary Goat anti-rabbit antibody conjugated to HRP (Pierce) was used at a concentration of 1:30,000 for 1hr in 1% Milk. Membrane was subsequently washed twice for 30 minutes total time.

The membrane was incubated with Supersignal west femto maximum substrate (Pierce) for 2 minutes and exposed to X-OMAT sensitive film (Kodak) for up to 30 minutes. The membrane was subsequently blotted with a monoclonal p21 WAF1 antibody (F5, Santa Cruz Biotech), p53 antibody (SC6243, polyclonal rabbit, Santa Cruz) (Ab-6, Oncogene Science), anti-hSir2 (polyclonal rabbit). β -actin was used (Abcam) for loading control. 9671S is an anti-acetyl H3 Lys9 was a monoclonal antibody (Cell Signaling).

Virus infection and stress response

All MEF cells were maintained in DMEM medium supplemented with 10% fetal bovine serum, and the IMR-90 cells were maintained in Eagle's minimal essential medium supplemented with 10% fetal bovine serum and non-essential amino acids. The virus infection and selection were essentially as described previously (Ferbeyre *et al.*, 2000). After one-week selection, the cells were either frozen for stock or immediately used for further analysis. About

500,000 MEF cells were plated on a 10-cm dish 24 hr before treatment. The cells were then exposed to etoposide (20 μ m) for 12 hr. After treatment, the cells were washed with PBS and fed with normal medium. Another 36 hrs later, the cells were stained with PI and analyzed by flow cytometric analysis for apoptotic cells (SubG1) according to DNA content. In case of the Fas-mediated apoptosis assay, the cells were treated with actinomycin D (0.25 μ g/ml) and Fas antibody (100 ng/ml) as previously described (Di Cristofano *et al.*, 1999). In the case of oxidative stress response, the IMR-90 cells were treated with H₂O₂ (200 μ M) for 24 hrs.

Luciferase and apoptosis assays

H1299 cells were transfected using the Fugene6 protocols (Roche) with pCMVwtp53 in presence or absence of pCMVp300 and 5 μ g of p21P-Luc (containing a 2.4kb fragment of p21 linked to luciferase gene) as previously described (Vaziri *et al.*, 1997). All experiments were performed in triplicates.

Apoptosis was measured at approximately 48 hrs post transfection using the annexin V antigen and propidium iodide exclusion (Clontech laboratories).

Radiation survival curves of BJ cells were performed as described previously (Dhar *et al.*, 2000; Vaziri *et al.*, 1999).

FACS analysis for apoptosis assay

Both adherent and floating cells were combined and washed in cold PBS. For SubG1/FACs analysis, cells were fixed in methanol for 2 hr at -20 °C, rehydrated in PBS for 1 hr at 4 °C, and then reacted with the primary antibody (DO-1) for 30 min at room temperature. Cells were washed twice in PBS and incubated with a goat anti-mouse FITC-conjugated secondary antibody for 30 min at room temperature. Following incubation, cells were washed in PBS and treated with RNase A (50 μ g/ml) for 30 min at room temperature. Propidium iodide (PI: 2.5 μ g/ml) was added to the cells, and samples were then analyzed in a FACSCalibur (BD). A region defining high FITC fluorescence was determined, and the cells falling into this region were collected separately. The PI staining was recorded simultaneously in the red channel.

Immunofluorescence Assay

Immunofluorescence was performed essentially as the standard protocol (Guo *et al.*, 2000). After fixation, cells were exposed to two primary antibodies: p53 monoclonal antibody DO-1 (Santa Cruz) and α -Sir2 α for 1 hr at room temperature. The cells were washed three times with 1% BSA plus 0.2% Tween-20 in PBS and then treated with two secondary antibodies [a goat anti-rabbit IgG conjugated to Alexa 568 (Molecular Probes), and anti-mouse IgG-FITC (Santa-Cruz)]. DAPI was used for counter-staining to identify nuclei. The cells were further washed four times. Images were acquired from a Nikon Eclipse E600 fluorescent microscope (Hamamatsu Photonics).

Detecting acetylation levels of p53 in cells

The cells (human lung carcinoma cell lines H460 (wild-type p53) and H1299 (p53-null), human colon carcinoma HCT116 (wild-type p53), mouse embryonal carcinoma cell line F9 (wild-type p53), mouse embryonic fibroblast MEFs or others) were maintained in DMEM medium supplemented with 10% fetal bovine serum. For DNA damage response, about 1 million cells were plated on a 10-cm dish 24 hr before treatment. The cells were then exposed to etoposide (20 μ M) and or other drugs (0.5 μ M of TSA, 5 mM of nicotinamide, and 50 μ M of LLNL) as indicated for 6 hr.

After treatment, the cells were harvested for Western blot analysis. The rabbit polyclonal antibody specific for p300-mediated acetylated p53 [α -p53(Ac)-C] was raised and purified against the acetylated human p53 C-terminal peptide [p53 (Ac)-C: H-S55GQSTSRH55LMF-OH SEQ. ID No:1 (5 = acetylated Lysine)] as described before (Luo *et al.*, 2000).

This antibody recognizes the p300-mediated acetylated forms of both human and mouse p53. In the case of cotransfection assays testing for p53 acetylation levels, H1299 cells were transfected with 5 μ g of CMV-p53 plasmid DNA, 5 μ g of CMV-p300 plasmid DNA, and 10 μ g of pcDNA2-Sir2 α plasmid DNA as indicated. 24 hr after the transfection, the cells were lysed in a Flag-lysis buffer (50 mM Tris, 137 mM NaCl, 10 mM NaF, 1mM EDTA, 1% Triton X-100 and 0.2% Sarkosyl, 1 mM DTT, 10% glycerol, pH 7.8) with fresh proteinase inhibitors, 10 μ M TSA and 5mM nicotinamide (Sigma). The cell extracts were resolved by either 8% or 4-20% SDS-PAGE gels (Novex) and analyzed by Western blot with α -p53 (Ac)-C and α -p53 (DO-1).

Deacetylation assay of the p53 C-terminal peptide

The human p53 C-terminal peptide (residues 368-386+Cys; HLKSK(AcK)GQSTSRHK(AcK)LMFKC); (SEQ ID NO. 1) di-acetylated at positions 373 and 382 was synthesized and purified with HPLC. Deacetylation assays of this peptide by Sir2 and analyses of the reaction products were performed as described previously (Imai *et al.*, 2000).

EXAMPLES

Example 1. Mammalian Sir2 α interacts with p53 both *in vitro* and *in vivo*.

Mouse Sir2 α interacts with p53. The p53 protein can be divided into three distinct functional domains (Gu and Roeder, 1997): an amino-terminus that contains the transcriptional activation domain (NT: residues 1-73), a central core that contains the sequence-specific DNA-binding domain (M: residues 100-300), and the multifunctional carboxyl-terminus (CT: residues 300-393). The GST-p53 fusion proteins containing each domain as well as the full-length protein were expressed in bacteria and purified to near homogeneity on glutathione-agarose beads. As shown in Figure 1A, ³⁵S-labeled *in vitro* translated Sir2 α strongly bound to immobilized GST-p53 but not to immobilized GST alone (lane 1 vs. 6). Sir2 α was tightly bound to the C-terminal domain of p53 (GST-p53CT) (lane 4, Figure 1A), also bound to the central DNA-binding domain (GST-p53M), but showed no binding to the N-terminal domain of p53 (GST-p53NT) (lane 3 vs. 2, Figure 1A).

To test for the interactions between p53 and Sir2 α in cells, extracts from transiently-transfected p53-null cells (H1299) were immunoprecipitated with anti-Flag monoclonal antibody (M2). As shown in Figure 1B, p53 was detected in the immunoprecipitate obtained from H1299 cells cotransfected with constructs encoding Flag-Sir2 α and p53 (lane 2), but not from cells transfected with the p53 construct alone (lane 4). Conversely, Sir2 α was detected in the immunoprecipitates obtained from H1299 cells cotransfected with constructs encoding Sir2 α and Flag-p53 (lane 6, Figure 1B), but not from cells transfected with the Sir2 α construct alone (lane 8, Figure 1B). p53 interacts similarly with human SIRT1 (hSIRT1) (Figure 1C, D), the human ortholog of mouse Sir2 α (Frye, 1999; 2000), showing that p53 and mammalian Sir2 α interact.

Since mouse Sir2 α shares a highly conserved region at the C-terminus with human SIRT1 (Figure 1C), but not with any other mammalian Sir2 homologs (Frye, 1999; 2000), a polyclonal antibody against the C-terminus (amino acid 480-737) of mouse Sir2 α was developed. Anti-Sir2 α antisera (α -Sir2 α) was raised in rabbits against the purified GST-

Sir2 α (480-737) fusion protein. As shown in Western blots, this antibody can detect both mouse Sir2 α and human SIRT1 proteins, but not other human Sir2 homologs (Figure 2A, B).

p53 interaction with Sir2 α or hSIRT1 in normal cells without overexpression was studied employing this antibody. Cell extracts from human (H460) and mouse cells (F9), which express wild-type p53 proteins, were immunoprecipitated with α -Sir2 α , or with the pre-immune serum. Western blot analysis revealed that this antibody immunoprecipitated both Sir2 α and hSIRT1 (lower panels, Figure 2A, 2B). Human and mouse p53 were detected in the respective α -Sir2 α immunoprecipitations from cell extracts, but not in the control immunoprecipitations with the preimmune serum, showing that p53 interacts with mammalian Sir2 α in normal cells. In contrast to abrogation of the Mdm2-p53 interaction by DNA damage as previously reported (Shieh *et al.*, 1997), this interaction was stronger in cells after DNA damage treatment (Figure 2C), which shows mammalian Sir2 α is involved in regulating p53 during the DNA-damage response. Thus, p53 interacts with mammalian Sir2 α both *in vitro* and *in vivo*.

Example 2. Deacetylation of p53 by mammalian Sir2 α

p53 was deacetylated by mammalian Sir2 α *in vitro*. Mouse Sir2 α protein was expressed with the N-terminal Flag epitope in cells and purified to near homogeneity on the M2-agrose affinity column (lane 3, Figure 3A to determine). The GST-p53 fusion protein was acetylated by p300 in the presence of [14 C]-acetyl-CoA, and the acetylated p53 protein was purified on the GST affinity column. These highly purified recombinant proteins were used in this assay in order to avoid possible contamination by either inhibitory factors or other types of deacetylases.

As shown in Figure 3B, 14 C-labeled acetylated p53 was efficiently deacetylated by purified Sir2 α (lane 3), but not by a control eluate (lane 4). NAD is required for Sir2 α -mediated deacetylation of p53 (lane 2 vs. 3, Figure 3B). Further, the deacetylase inhibitor TSA, which significantly abrogates HDAC1-mediated deacetylase activity on p53 (Luo *et al.*, 2000), had no apparent effect on Sir2 α -mediated p53 deacetylation (lane 5, Figure 3B). These results show that Sir2 α can strongly deacetylate p53 *in vitro*, and that this activity depends on NAD.

A role for mammalian Sir2 α in deacetylating p53 in cells was established using acetylated p53-specific antibody to monitor the steady-state levels of acetylated p53 *in vivo* (Luo *et al.*, 2000). As shown in Figure 3C, a high level of acetylated p53 was detected in the cells cotransfected with p300 and p53 (lane 1). However, p53 acetylation levels were significantly

abolished by expression of either Sir2 α or hSIRT1 (lanes 2, 4). In contrast, a Sir2 α mutant (Sir2 α H355A) containing a point mutation at the highly conserved core domain causing defective histone deacetylase activity *in vitro* had almost no effect (lane 3 vs. 2, Figure 3C). Furthermore, neither SIRT5, another human Sir2 homolog, nor poly (ADP-ribose) polymerase (PARP), whose activity is also NAD-dependent (reviewed in Vaziri *et al.*, 1997), had any significant effect on p53 acetylation (lanes 5, 6, Figure 3C). In addition, in contrast to HDAC-mediated deacetylation of p53 (Luo *et al.*, 2000) Sir2 α still strongly deacetylated p53 in the presence of TSA (lane 4 vs. 3, Fig. 3D) even though the steady state level of acetylated p53 was elevated when the cells were treated with TSA (lane 3 vs. 1, Fig. 3D). Thus, mammalian Sir2 α has robust TSA-independent p53 deacetylation activity.

Example 3. Inhibition of Sir2 α -mediated p53 deacetylation by nicotinamide

Sir2 α -mediated deacetylase activity of p53 can be inhibited. Deacetylation of acetylated lysine by Sir2 α is tightly coupled to NAD hydrolysis, producing nicotinamide and a novel acetyl-ADP-ribose compound (1-O-acetyl-ADPribose) (Landry *et al.*, 2000b; Tanner *et al.*, 2000; Tanny and Moazed, 2001). The formation of an enzyme-ADP-ribose intermediate through NAD hydrolysis may be critical for this chemical reaction (Landry *et al.*, 2000b). Since nicotinamide is the first product from hydrolysis of the pyridinium-N-glycosidic bond of NAD, it may function as an inhibitor for its deacetylase activity (Landry *et al.*, 2000b). Nicotinamide is able to inhibit the deacetylase activity of Sir2 α on acetylated p53 *in vitro*.

Similar reactions as described above (Figure 3B), were set up by incubating labeled p53 substrate, recombinant Sir2 α and NAD (50 μ M) alone, or in combination with nicotinamide (5mM). As shown in Figure 4A, ¹⁴C-labeled acetylated p53 was efficiently deacetylated by Sir2 α (lane 2) however, the deacetylation activity was completely inhibited in the presence of nicotinamide (lane 3 vs. lane 2 Figure 4A). As a negative control, 3-AB (3-aminobenzamide), a strong inhibitor of PARP which is involved in another type of NAD-dependent protein modification (Vaziri *et al.*, 1997), showed no significant effect on Sir2 α mediated deacetylation (lane 4 vs. 3, Figure 4A).

To further investigate the role of mammalian Sir2 α -mediated regulation *in vivo*, the effect of Sir2 α expression on p53 acetylation levels during the DNA damage response was determined. Mouse embryonic fibroblast (MEF) cells, which express the wild type of p53, were

infected with either a pBabe-puro retrovirus empty vector or a pBabe-puro retrovirus containing Sir2 α , and cultured for a week under pharmacological selection. The protein levels of p53 activation in response to DNA damage in these cells was determined by Western blot analysis. Similar protein levels of p53 activation were induced in the pBabe vector infected cells and pBabe-Sir2 α infected cells after etoposide treatment for 6 hrs (lanes 3, 4 vs. lanes 1, 2, lower panel, Figure 4B).

In the mock-infected cells, the acetylation level of p53 was significantly enhanced by DNA damage (lane 2 vs. lane 1, Upper panel, Figure 4B). However, DNA damage treatment failed to stimulate the p53 acetylation in the pBabe-Sir2 α infected cells even in the presence of TSA (lane 4 vs. lane 2, Upper panel, Figure 4B), showing that Sir2 α expression results in deacetylation of endogenous p53. This Sir2 α -mediated effect was completely abrogated by nicotinamide treatment (lane 8 vs. lane 6, Figure 4B). Thus, Sir2 α mediated deacetylation of p53 can be inhibited by nicotinamide both *in vitro* and *in vivo*.

Example 4. Maximum induction of p53 acetylation levels in normal cells requires inhibition of endogenous Sir2 α activity

Endogenous Sir2 α in the regulation of p53 acetylation levels in normal cells during the DNA damage response was determined.

As shown in Figure 4C, after the wild-type p53 containing human lung carcinoma cells (H460) were treated by etoposide, acetylation of p53 was induced (lane 2 vs. lane 1). No significant p53 acetylation was detected in the cells treated with a proteasome inhibitor LLNL (lane 6, Figure 4C), indicating that the observed stimulation of p53 acetylation is induced by DNA damage, not through p53 stabilization.

p53 can be deacetylated by a PID/MTA2/HDAC1 complex, whose activity is completely abrogated in the presence of TSA (Luo *et al.*, 2000). The mild enhancement of the acetylation level of p53 by TSA during DNA damage response may be due mainly to its inhibitory effect on endogenous HDAC1-mediated deacetylase activity (lane 3 vs. lane 2, Figure 4C). A super induction of p53 acetylation was showed when the cells were treated with both TSA and nicotinamide (lane 4 vs. lane 3, Figure 4C). In contrast, 3-AB treatment had no effect on the level of p53 acetylation (lane 5 vs. lane 3, Figure 4C), indicating that PARP-mediated poly-ADP ribosylation has no effect on p53 acetylation. Similar results were also observed in other cell

types including either mouse cells (MEFs, F9) or human cells (BL2, HCT116). Thus, maximum induction of p53 acetylation requires inhibitors for both types of deacetylases (HDAC1 and Sir2 α), and endogenous Sir2 α plays a major role in the regulation of the p53 acetylation levels induced by DNA damage.

Example 5. Repression of p53-mediated functions by mammalian Sir2 α requires its deacetylase activity

The functional consequence of mammalian Sir2 α -mediated deacetylation of p53 was determined by testing its effect on p53-mediated transcriptional activation. A mammalian p53 expression vector (CMV-p53), alone or in combination with different amounts of mouse Sir2 α expressing vector (CMV-Sir2 α), was cotransfected into MEF (p53^{-/-}) cells along with a reporter construct containing synthetic p53 binding sites placed upstream of the luciferase gene (PG13-Luc).

As shown in Figure 5A, Sir2 α strongly repressed p53-mediated transactivation in a dose-dependent manner (up to 21 fold), but had no significant effect on the transcriptional activity of the control reporter construct (TK-Luc) (Figure 5B), which has no p53 binding site at the promoter region. Also, expression of human SIRT1 showed a similar effect on the p53 target promoter (Figure 5C). Neither the Sir2 α H355A mutant or SIRT5, both of which are defective in p53 deacetylation (Figure 3C), had any effect on the p53-mediated transactivation (Figure 5C, D). Thus, mammalian Sir2 α specifically represses p53-dependent transactivation, and that this repression requires its deacetylase activity.

The modulation of Sir2 on p53-dependent apoptosis was determined. p53 null cells (H1299) were transfected with p53 alone or cotransfected with p53 and Sir2 α . The transfected cells were fixed, stained for p53, and analyzed for apoptotic cells (SubG1) (Luo *et al.*, 2000). As indicated in Figure 6A, overexpression of p53 alone induced significant apoptosis (32.3% SubG1). However, co-transfection of p53 with Sir2 α significantly reduced the level of apoptosis (16.4% SubG1), while the mutant Sir2 α H355A was impaired in this effect (29.5% SubG1) (Figure 6A, B). Thus, mammalian Sir2 α is involved in the regulation of both p53 mediated transcriptional activation and p53-dependent apoptosis, and deacetylase activity is required for these Sir2 α -mediated effects on p53.

Example 6. The role of mammalian Sir2 α in stress induced apoptotic response

Mammalian Sir2 α can deacetylate p53 both *in vitro* and *in vivo* (Figure 3). Sir2 α can block the induction of endogenous p53 acetylation levels by DNA damage (Figure 4B, 4C). To elucidate the physiological significance for this Sir2 α mediated regulation, the effect on DNA damage-induced apoptotic response was determined.

MEF (p53^{+/+}) cells as described above (Figure 4B), were infected with either a pBabe-puro retrovirus empty vector or a pBabe-puro retrovirus containing Sir2 α . After the DNA damage treatment by etoposide, the cells were stained with PI and analyzed by flow cytometric analysis for apoptotic cells (SubG1) according to DNA content. As shown in Figure 7A, the cells mock infected with the pBabe-vector, were susceptible to etoposide-induced cell death, with about 48% of the cells apoptotic after exposure to 20 μ M of etoposide (3 vs. 1, Figure 7A). In contrast, the pBabe-Sir2 α infected MEF (p53^{+/+}) cells were more resistant to apoptosis induced by the same dose of etoposide, with only 16.4% apoptotic cells (4 vs. 3, Figure 7A). Since no significant apoptosis was detected in MEF (p53^{-/-}) cells by the same treatment, the induced apoptosis observed in MEF (p53^{+/+}) cells is totally p53-dependent. Thus, Sir2 α significantly inhibits p53-dependent apoptosis in response to DNA damage.

The role of mammalian Sir2 α in the oxidative stress response was determined. Recent studies have indicated that oxidative stress-induced cell death is p53-dependent (Yin *et al.*, 1998; Migliaccio *et al.*, 1999). Early-passage normal human fibroblast (NHF) IMR-90 cells were employed for this study since p53-dependent apoptosis can be induced by hydrogen peroxide treatment in these cells (Chen *et al.*, 2000).

IMR-90 cells were infected with either a pBabe-puro retrovirus empty vector or a pBabe-puro retrovirus containing Sir2 α , and cultured for a week under pharmacological selection. By immunofluorescence staining, p53, in these infected cells, was induced significantly after hydrogen peroxide treatment, along with Sir2 α localized in the nuclei detected by immunostaining with specific antibodies (Figure 7C). Sir2 α expression significantly promotes cell survival under oxidative stress. As indicated in Figure 7D, the cells mock infected with the pBabe-vector, were susceptible to H₂O₂-induced cell death, with more than 80% of the cells being killed after 24 hr exposure to 200 μ M H₂O₂ (II vs. I). In contrast, the pBabe-Sir2 α infected cells were much more resistant to death by the same dose of H₂O₂, with about 70% of

the cells surviving after 24 hr of H₂O₂ treatment (IV vs. III, Figure 7D). Mammalian Sir2 α promotes cell survival under stress by inhibiting p53-dependent apoptosis.

Example 7. Mammalian Sir2 α has no effect on p53-independent cell death induced by anti-Fas

The specificity of mammalian Sir2 α -mediated protection of cells from apoptosis was examined by determining whether Sir2 α has any effect of p53-independent, Fas-mediated apoptosis. The MEF (p53^{-/-}) cells were first infected with either a pBabe-puro retrovirus empty vector or a pBabe-puro retrovirus containing Sir2 α , then cultured for a week under pharmacological selection. After the treatment by anti-Fas (100 ng/ml) for 24 hrs, the cells were harvested and further analyzed for apoptotic cells (SubG1).

Cells mock infected with the pBabe vector, were susceptible to anti-Fas induced cell death, with about 31.7% of the cells becoming apoptotic. However, in contrast to the strong protection of p53-dependent apoptosis by Sir2 α during DNA damage response in the MEF (p53^{+/+}) cells (Figure 7A, B), Sir2 α expression had no significant effect on Fas-mediated apoptosis in the MEF (p53^{-/-}) cells. Thus, mammalian Sir2 α regulates p53-mediated apoptosis.

Mammalian Sir2 α has no effect on the Fas mediated apoptosis. (A) Both mock infected cells and pBabe-Sir2 α infected MEF p53(-/-) cells were either not treated (1 and 2) or treated with 100 ng/ml Fas antibody in presence of actinomycin D (0.25 μ g/ml) (3 and 4). The cells were analyzed for apoptotic cells (subG1) according to DNA content (PI staining). The representative results depict the average of three experiments with standard deviations indicated.

Example 8. Physical interaction of hSir2 with p53

p53 protein is acetylated in response to DNA damage and the acetylation contributed to the functional activation of p53 as a transcription factor (Abraham *et al.*, 2000; Sakaguchi *et al.*, 1998). Sir2 is a deacetylase of p53, thereby modulating functioning of p53 as a transcription factor.

In order to study the functional interaction between p53 and hSir2, a full length human hSir2SIRT1 cDNA clone (obtained from the IMAGE consortium (Frye, 1999)) was introduced into a pBabe-based retroviral expression vector which also carries puromycin resistance gene as

a selectable marker. The resulting construct was termed pYESirwt. A retroviral construct bearing a derived, mutant allele of Sir2 and termed pYESirHY was constructed and used in parallel as control. This mutant allele encodes an amino acid substitution at residue 363, at which site the normally present histidine is replaced by tyrosine. This H to Y substitution results in an alteration of the highly conserved catalytic site of the hSir2 protein and subsequent neutralization of its deacetylase activity. These vector constructs were used to transduce the hSIR2SIRT1 gene both by transfection and retroviral infection.

A polyclonal rabbit antibody that specifically recognizes the C-terminal portion of hSir2 was developed and its specificity validated by immunoprecipitation and Western blotting (Figure 8A). Both the endogenous and the ectopically expressed hSir2 proteins were detected as protein species of 120 Kilodalton (Kd) rather than as 80Kd polypeptide predicted from the known primary sequence of hSIR2SIRT1 (Figure 8A). Localization of hSir2 protein by immunofluorescence using the hSir2 antibody showed a punctate nuclear staining pattern (Figure 8B).

The physical interactions between hSir2 and p53 were evaluated by co-transfecting the pYESir2wt plasmid and a vector expressing wt p53 under the control of the cytomegalovirus promoter (pCMV-wtp53) transiently into H1299 human non-small cell lung carcinoma cells (Brower *et al.*, 1986) which have a homozygous deletion of the p53 gene and produce no p53 mRNA or protein (Mitsudomi *et al.*, 1992). Cell lysates were subsequently mixed with the rabbit anti-hSir2 antibody and resulting immune complexes were collected by protein G and analyzed by SDS-PAGE electrophoresis and immunoblotting. The immunoblot was probed with a sheep anti-p53 antibody (Figure 8C) and reprobated it subsequently with an anti-hSir2 antibody (top panel) to verify presence of hSir2 in the complex. As indicated in Figure 8C, immunoprecipitation of hSir2 resulted in co-precipitation of p53.

In the reciprocal experiment, lysates of BJT cells, human fibroblasts into which the telomerase gene has been introduced, were examined. In addition, these cells express either the wild type hSir2 vector or the hSir2HY mutant. Two cell populations were created by infection of mass cultures of BJT cells with the respective vectors and subsequent selection in puromycin. The anti-p53 antibody was employed to immunoprecipitate complexes and subsequently probe the resulting immunoblot with either polyclonal anti-p53 antibodies or an anti-hSir2 antibody. These immunoblots demonstrated a physical interaction between hSir2 and p53 proteins (Figure

8D). Formation of these complexes was unaffected by the H to Y mutation introduced into the hSir2 catalytic site (Figure 8D). Furthermore, radiation used to increase the levels of p53 protein in BJ cells had no effect on the levels of p53:hSir2 complexes. Comparison of the immunoprecipitated p53 to total input p53 resulted in an estimate of approximately 1% of the cells complement of p53 protein was present in physical complexes with hSir2.

Example 9. Deacetylation of p53 by hSir2 *in vitro*

Since hSir2 forms physical complexes with p53, the ability of Sir2 to deacetylate human p53 *in vitro* was evaluated. Since adequate quantities of bacterially produced hSir2 were not available, bacterially expressed mouse SIR2 (mSir2a) enzyme was used in *in vitro* assays (Imai *et al.*, 2000). A 20 residue-long oligopeptide that contains the sequence corresponding to residues 368-386+Cys of the human p53 protein was used as a substrate in these reactions. Lysine residues corresponding to residues 373 and 382 of the p53 protein were synthesized in acetylated form in this oligopeptide substrate. These two residues of p53 are known to be acetylated by p300 (Gu and Roeder, 1997) following γ or UV irradiation (Liu *et al.*, 1999; Sakaguchi *et al.*, 1998) with acetylation of lysine residue 382 being favored in response to ionizing radiation *in vivo* (Abraham *et al.*, 2000). This p53 oligopeptide serves as an excellent surrogate p53 substrate *in vitro* for acetylation studies (Gu and Roeder, 1997).

The deacetylase activity of hSir2 utilizes NAD as a co-factor (Imai *et al.*, 2000; Moazed, 2001; Smith *et al.*, 2000; Tanner *et al.*, 2000; Tanny *et al.*, 1999). In the absence of added NAD, incubation of mSir2 with p53 oligopeptide gave rise to a single prominent peak (peak 1) and a small, minor peak (peak 2) upon high pressure liquid chromatography (HPLC), corresponding to the monomeric and dimeric forms of the peptide, respectively (Figure 9A). However, incubation in the presence of 1mM NAD produced a singly deacetylated species as the major product (peak 3, Figure 9B). Edman sequencing of this singly deacetylated species revealed that mSir2 preferentially deacetylated the residue corresponding to Lys 382 of p53 (Figure 9, C-F), having relatively weak effect on Lys 373. Thus, the acetylated p53 peptide acted as a substrate for hSir2 and indicated that the de-acetylation of p53 at Lys 382 by mammalian Sir2 is specific and not the result of an indiscriminate deacetylase function.

Example 10. Deacetylation of p53 by hSir2 *in vivo*

The ability of hSir2 to deacetylate intact p53 protein *in vivo* was evaluated. To produce acetylated p53 *in vivo*, the p53 expression plasmid was co-transfected with one expressing p300. This protocol leads to acetylation of p53 in the absence of exposure to DNA-damaging agents (Luo *et al.*, 2000). The ability of hSir2 to deacetylate the p53 protein at its K382 residue in H1299 cells that lack endogenous p53 gene was determined. The levels of acetylation of p53 at Lys382 were monitored by using a rabbit polyclonal antibody, termed Ab-1, which had been raised against the acetylated K382 of p53 protein. The specificity of the Ab-1 antibody has been demonstrated (Sakaguchi *et al.*, 1998).

Co-transfection of plasmids expressing wild-type p53 and p300 into H1299 cells showed that p53 protein is readily acetylated at K382, as detected by probing the immunoblot with the Ab-1 antibody (Figure 10A, lane 3). Recognition of this acetylated form of p53 by the Ab-1 antibody was specific, since a mutant p53 protein that was expressed in a parallel culture of H1299 cells and carries an arginine rather than a lysine at residue 382 was not recognized by the Ab-1 antibody, despite ectopic expression of the p300 acetylase. (Figure 10A, lane 6).

Co-transfection of the hSir2-expression plasmid with the p53- and p300-expressing plasmids substantially decreased the acetylated p53 that could be detected by the Ab-1 antibody. (Figure 10A, lane 5). The residual level of acetylated p53 could be further reduced by increasing the amount of co-transfected hSir2 expression plasmid. Thus, hSir2 can deacetylate p53 protein at the Lys382 residue *in vivo*.

The hSir2HY vector, which expresses the mutant-catalytically inactive hSir2, was introduced into these H1299 cells. The mouse equivalent of this hSir2HY mutant lacks 95% of its deacetylase activity (Imai *et al.*, 2000). The hSir2HY mutant failed to deacetylate wt p53 efficiently, indicating that the catalytic activity of the introduced wild type hSir2 gene product was required for specific deacetylation of p53 Lys 382 (Figure 10A, lane 9).

The lysine 320 residue of p53 is also acetylated by PCAF in response to DNA damage (Sakaguchi *et al.*, 1998). Whether the state of acetylation of residue 320 affected the ability of hSir2 to deacetylate residue 382 was determined. A mutant p53 allele that specifies a lysine-to-arginine substitution at residue 320 was expressed. This amino acid substitution did not affect the ability of hSir2 to deacetylate the K382 residue in H1299 cells, indicating that the action of hSir2 on the acetylated K382 residue is independent of the state of acetylation of the K320 residue (Figure 10A, lanes 7, 8).

As a measure of the substrate specificity of hSir2, the effects of hSir2 on histone acetylation, specifically the acetylated residue lysine 9 of histone H3, were determined using cell nuclei from the above experiments. H3 Lys9 acetylation was monitored through the use of the 9671S monoclonal antibody. The 9671S antibody specifically recognizes histone H3 that is acetylated at this position.

Neither wildtype hSir2 nor the catalytically inactive hSir2HY altered the acetylation of histone H3 at this position (Figure 10A, bottom). Thus, de-acetylation of p53 Lys382 *in vivo* reflects a defined substrate specificity of hSir2 and not a non-specific consequence of its over-expression.

Example 11. hSir2 and p53 acetylation in primary and tumor cell lines

Acetylation of lysine residue 382 of p53 accompanies and mediates the functional activation of p53 as a transcription factor following exposure of a cell to ionizing radiation (Sakaguchi *et al.*, 1998). To determine whether hSir2 could antagonize and reverse this activation of p53, by its deacetylase function, either wildtype hSir2 or the mutant form specified by the hSir2HY vector was expressed in BJT human fibroblast cells. Ectopic expression of the telomerase enzyme in these BJT cells, undertaken to extend their lifespan, had no effect on either their activation of p53 protein or their responses to DNA damage (Vaziri *et al.*, 1999).

In order to facilitate detection of *in vivo* acetylated p53 protein, BJT cells were exposed to 6Gy of ionizing radiation in the presence of low trichostatin A (TSA) concentrations. While not directly inhibiting hSir2 catalytic activity (Imai *et al.*, 2000), TSA appears to increase the stability of acetylated p53 protein (Sakaguchi *et al.*, 1998), perhaps by inhibiting non-hSir2 deacetylases that also recognize the acetylated p53 K382 residue. The resulting immunoblot was probed with the polyclonal rabbit antiserum (Ab-1) which specifically recognizes the acetylated K382 form of p53.

Following 6 Gy of ionizing radiation, a 1.5-2 fold increase in the level of acetylated p53 protein was observed, as indicated by the levels of p53 protein recognized by the Ab-1 antiserum (Figure 10B). A four-fold increase in hSir2 levels, achieved through ectopic expression of hSir2, resulted in the reversal of the radiation-induced increase in acetylated K382 p53 protein (Figure 10B). In contrast, expression of the catalytically inactive hSirHY protein at comparable levels increased the radiation-induced levels of p53 acetylated at residue K382 (Figure 10B) suggesting

that the hSir2HY mutant may act in a dominant negative fashion in BJT cells. A re-probing of this immunoblot with a polyclonal anti-p53 antibody showed normal stabilization of p53 in control cells in response to DNA damage and at most, slightly reduced levels of stabilization in the presence of ectopically expressed wild type hSir2 (Figure 10B). Hence, while hSir2 is able to reverse the radiation-induced acetylation of p53 in these cells, it has only minimal effects on the metabolic stabilization of p53 induced by exposure to radiation.

A similar phenomenon was observed in MCF-7 human breast carcinoma line cells, which have retained an apparently intact p53-dependent checkpoint in response to ionizing radiation. Irradiation of these cells led to a three-fold increase in acetylated p53 levels, while a four-fold ectopic expression of wild type hSir2 in irradiated MCF-7 cells led to deacetylation of p53 protein (Figure 10C). In contrast to BJT cells, no significant change in the stability of total p53 protein was observed. However, MCF-7 cells expressing the hSirHY mutant showed a level of radiation-induced acetylation that was comparable to control irradiated cells (Figure 10C). Thus, hSir2 is able to reverse the radiation-induced acetylation in both BJT and MCF-7 cells, suggesting that hSir2 acts as an antagonist of p53 function *in vivo*.

The differences observed in deacetylation activities of hSir2HY in MCF7 and BJT cells may reflect the ability of hSir2HY to act as a dominant-negative allele in BJT cells. BJT cells do express significantly lower levels of endogenous hSir2 when compared with MCF7 cells. These lower levels of hSir2 in BJT cells may enable hSir2HY to form inhibitory complexes with endogenous wild type hSir2 or with other proteins required for its function. In this context, evidence in yeast suggests that H363Y mutant does indeed act as a potent dominant-negative (Tanny *et al.*, 1999).

Example 12. Effects of hSir2 on the transcriptional activity of p53 protein

The effects of hSir2 on the transcriptional activity of p53 were determined by co-transfecting H1299 cells transiently with a p53 expression plasmid and a reporter construct in which the promoter of the p21WAF1 gene (el-Deiry *et al.*, 1993), a known target of transcriptional activity by p53, is able to drive expression of a luciferase reporter gene (Vaziri *et al.*, 1997). As indicated in Figure 11A, luciferase activity increased in response to increasing amount of co-transfected wtp53 expression vector. Conversely, the transcriptional activity of p53 protein was suppressed by co-expression of wild type hSir2 in a dose-dependent fashion.

The catalytically inactive hSir2HY mutant had no effect on p53 transcriptional activity (Figure 11A). The specificity of hSir2 in affecting promoter activity was determined using a constitutively active SV40 promoter linked to the luciferase gene. Expression of this control construct was not affected by increasing amounts of hSir2 expression vector at any level (Figure 11B).

The above observations were confirmed in a more physiologic context using a subline of MCF-7L cells. The subline of MCF-7 cells was stably transfected with a p21WAF1 promoter-reporter construct. In addition, these cells were infected stably with retroviral vector constructs expressing either the wild type hSir2 or the mutant hSir2HY. These cells were exposed to 6 Gy of ionizing radiation and subsequently measured total p53 and p21WAF1 protein levels (Figure 11C).

p53 protein levels increased normally in all cell populations in response to irradiation of these cells. However, the levels of p21WAF1 protein were reduced in cells expressing wild type hSir2 (Figure 11C). Moreover, MCF-7L cells expressing the mutant hSir2HY protein had a higher level of p21WAF1 when compared with the irradiated controls and with the wild type hSir2-overexpressing cells (Figure 11C) showing that the hSir2HY mutant may act in a dominant-negative fashion in these cells. Thus, hSir2 can antagonize the transcriptional activities of p53 that enable it to exert cytostatic effects via transcriptional activation of the p21WAF1 gene.

Example 13. Inhibition of p53-dependent apoptosis by hSir2

hSir2 can antagonize the ability of p53 to act in a cytostatic fashion through induction of p21WAF1 synthesis. The ability of hSir2 to blunt the pro-apoptotic functions of p53 was determined. Restoration of wild-type p53 function in H1299 cells, achieved via introduction of a wt p53-expressing vector, induces apoptosis, as indicated by the expression of the cell surface annexin V antigen (Figure 12A). Co-transfection of a p300 vector with the p53 gene increased this p53-dependent apoptosis (Figure 12A). This apoptotic response was abolished in a dose-dependent manner in cells co-transfected with increasing amounts of the wt hSir2 expression plasmid (Figure 12A). Hence, hSir2 antagonizes both the cytostatic effects of p53 (as mediated by p21WAF1) and its pro-apoptotic effects.

Example 14. Effects of mutant hSir2HY on radiosensitivity of human fibroblasts

In contrast to the behavior of many other murine or human cell lines, human fibroblasts become relatively radioresistant upon inactivation of p53 function (Tsang *et al.*, 1995). This behavior suggested an additional test of the ability of hSir2 to antagonize p53 function, which depended on measuring the long-term survival of human BJT fibroblasts cells following exposure to various doses of low-level ionizing radiation.

Ectopic expression of wild type hSir2 in these cells led to a greater long-term survival (Figure 12B, triangles), while expression of the mutant hSir2HY in BJT cells led to a radiosensitive phenotype (Figure 12B, diamonds) consistent with hSir2HY constructs acting in a dominant-negative fashion in BJT cells. A positive control cell line derived from an individual with ataxia telangiectasia (AT) was highly radiosensitive (Figure 12B, circles). The central role of p53 in these various responses was also shown in the behavior of a subline of BJT fibroblasts that express a dominant-negative form of p53 and also have acquired a measure of radioresistance (Figure 12B, open square). Thus, wt hSir2 antagonizes p53 activity while the hSir2HY mutant potentiates its activity.

REFERENCES

Appella E. and Anderson C.W. (2000) Signaling to p53: breaking the posttranslational modification code. *Pathol. Biol.*, 48:227-245.

Avantaggiati M.L., Ogryzko V., Gardner K., Giordano A., Levine A.S. and Kelly K. (1997) Recruitment of p300/CBP in p53-dependent signal pathways. *Cell*, 89:1175-1184.

Bernstein B.E., Tong J.K. and Schreiber S.L. (2000) Genomewide studies of histone deacetylase function in yeast. *Proc. Natl. Acad. Sci. USA*, 97:13708-13713.

Butler L.M., Agus D.B., Scher H.I., Higgins B., Rose A., Cordon-Cardo C., Thaler H.T., Rifkind R.A., Marks P.A. and Richon V.M. (2000) Suberoylanilide hydroxamic acid, an inhibitor of histone deacetylase suppresses the growth of prostate cancer cells in vitro and in vivo. *Cancer Res.*, 60:5165-5170.

Campisi J. (2000) Aging, chromatin, and food restriction—connecting the dots. *Science*, 289:2062-2063.

Chao C., Saito S., Kang J., Anderson C.W., Appella E. and Xu Y. (2000) p53 transcriptional activity is essential for p53-dependent apoptosis following DNA damage. *EMBO J.*, 19:4967-4975.

Chen Q.M., Liu J. and Merrett J.B. (2000) Apoptosis or senescence-like growth arrest: influence of cell-cycle position, p53, p21 and bax in H₂O₂ response of normal human fibroblasts. *Biochem. J.*, 347:543-551.

Chresta C.M. and Hickman J.A. (1996) Oddball p53 in testicular tumors. *Nat. Med.*, 2:745-746.

Di Cristofano A., Kotsi P., Peng Y.F., Cordon-Cardo C., Elkon K.B. and Pandolfi P.P. (1999) Impaired Fas response and autoimmunity in Pten^{+/-} mice. *Science*, 285:2122-2125.

Ferbeyre G., de Stanchina E., Querido E., Baptiste N., Prives C. and Lowe S.W. (2000) PML is induced by oncogenic ras and promotes premature senescence. *Genes Dev.*, 14:2015-2027.

Frye R.A. (1999) Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity. *Biochem. Biophys. Res. Commun.*, 260:273-279.

Frye R.A. (2000) Phylogenetic classification of prokaryotic and eukaryotic Sir2-like proteins. *Biochem. Biophys. Res. Commun.*, 273:793-798.

Gu W. and Roeder R.G. (1997) Activation of p53 sequence-specific DNA binding by acetylation of the p53 C-terminal domain. *Cell*, 90:595-606.

Gu W., Shi X.L. and Roeder R.G. (1997) Synergistic activation of transcription by CBP and p53. *Nature*, 387:819-823.

Gu W., Malik S., Ito M., Yuan C.X., Fondell J.D., Zhang X., Martinez E., Qin J. and Roeder R.G. (1999) A novel human SRB/MED-containing cofactor complex, SMCC, involved in transcription regulation. *Mol. Cell*, 3:97-108.

Guarente L. (2000) Sir2 links chromatin silencing, metabolism, and aging. *Genes Dev.*, 14:1021-1026.

Guo A., Salomoni P., Luo J., Shih A., Zhong S., Gu W. and Pandolfi P.P. (2000) The function of PML in p53-dependent apoptosis. *Nat. Cell Biol.*, 2:730-736.

Imai S., Armstrong C.M., Kaeberlein M. and Guarente L. (2000) Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature*, 403:795-800.

Ito A., Lai C., Zhao X., Saito S., Hamilton M., Appella E. and Yao T. (2001) p300/CBP-mediated p53 acetylation is commonly induced by p53-activating agents and inhibited by MDM2. *EMBO J.*, 20:1331-1340.

Jimenez G.S., Nister M., Stommel J.M., Beeche M., Barcarse E.A., Zhang X.Q., O'Gorman S. and Wahl G.M. (2000) A transactivation-deficient mouse model provides insights into Trp53 regulation and function. *Nat. Genet.*, 26:37-43.

Kouzarides T. (2000) Acetylation: a regulatory modification to rival phosphorylation? *EMBO J.*, 19:1176-1179.

Landry J., Slama J.T. and Sternglanz R. (2000) Role of NAD(+) in the deacetylase activity of the SIR2-like proteins. *Biochem. Biophys. Res. Commun.*, 278:685-690.

Landry J., Sutton A., Tafrov S.T., Heller R.C., Stebbins J., Pillus L. and Sternglanz R. (2000) The silencing protein SIR2 and its homologs are NAD-dependent protein deacetylases. *Proc. Natl. Acad. Sci. USA*, 97:5807-5811.

Levine A.J. (1997) p53, the cellular gatekeeper for growth and division. *Cell*, 88:323-331.

Lill N.L., Grossman S.R., Ginsberg D., DeCaprio J. and Livingston D.M. (1997) Binding and modulation of p53 by p300/CBP coactivators. *Nature*, 387:823-827.

Lin S.J., Defossez P.A. and Guarente L. (2000) Requirement of NAD and SIR2 for life-span extension by calorie restriction in *Saccharomyces cerevisiae*. *Science*, 289:2126-2128.

Luo J., Su F., Chen D., Shiloh A. and Gu W. (2000) Deacetylation of p53 modulates its effect on cell growth and apoptosis. *Nature*, 408:377-381.

Lutzker S.G. and Levine A.J. (1996) A functionally inactive p53 protein in teratocarcinoma cells is activated by either DNA damage or cellular differentiation. *Nat. Med.*, 2:804-810.

Marks P.A., Rifkind R.A., Richon V.M. and Breslow R. (2001) Inhibitors of histone deacetylase are potentially effective anticancer agents. *Clin. Cancer Res.*, 7:759-760.

Maya R., Balass M., Kim S.T., Shkedy D., Leal J.F., Shifman O., Moas M., Buschmann T., Ronai Z. and Shiloh Y. et al. (2001) ATM-dependent phosphorylation of Mdm2 on serine 395: role in p53 activation by DNA damage. *Genes Dev.*, 15:1067-1077.

Migliaccio E., Giorgio M., Mele S., Pelicci G., Reboldi P., Pandolfi P.P., Lanfrancone L. and Pelicci P.G. (1999) The p66shc adaptor protein controls oxidative stress response and life span in mammals. *Nature*, 402:309-313.

Nakamura S., Roth J.A. and Mukhopadhyay T. (2000) Multiple lysine mutations in the C-terminal domain of p53 interfere with MDM2-dependent protein degradation and ubiquitination. *Mol. Cell. Biol.*, 20:9391-9398.

Nakano K. and Vousden K. (2001) PUMA, a novel proapoptotic gene, is induced by p53. *Mol. Cell*, 7:683-694.

Pearson M., Carbone R., Sebastiani C., Cioce M., Fagioli M., Saito S., Higashimoto Y., Appella E., Minucci S., Pandolfi P.P. and Pelicci P.G. (2000) PML regulates p53 acetylation and premature senescence induced by oncogenic Ras. *Nature*, 406:207-210.

Prives C. and Hall P.A. (1999) The p53 pathway. *Pathol. J.*, 187:112-126.

Rodriguez M.S., Desterro J.M., Lain S., Lane D.P. and Hay R.T. (2000) Multiple C-terminal lysine residues target p53 for ubiquitin-proteasome-mediated degradation. *Mol. Cell. Biol.*, 20:8458-8467.

Shieh S.Y., Ikeda M., Taya Y. and Prives C. (1997) DNA damage-induced phosphorylation of p53 alleviates inhibition MDM2. *Cell*, 91:325-334.

Shore D. (2000) The Sir2 protein family: A novel deacetylase for gene silencing and more. *Proc. Natl. Acad. Sci. USA*, 97:14030-14032.

Smith J.S., Brachmann C.B., Celic I., Kenna M.A., Muhammad S., Starai V.J., Avalos J.L., Escalante-Semerena J.C., Grubmeyer C., Wolberger C. and Boeke J.D. (2000) A phylogenetically conserved NAD-dependent protein deacetylase activity in the Sir2 protein family. *Proc. Natl. Acad. Sci. USA*, 97:6658-6663.

Sterner D.E. and Berger S.L. (2000) Acetylation of histones and transcription-related factors. *Microbiol. Mol. Biol.*, 64:435-459.

Tanner K.G., Landry J., Sternglanz R. and Denu J.M. (2000) Silent information regulator 2 family of NAD-dependent histone/protein deacetylases generates a unique product, 1-O-acetyl-ADP-ribose. *Proc. Natl. Acad. Sci. USA*, 97:14178-14182.

Tanny J.C. and Moazed D. (2001) Coupling of histone deacetylation to NAD breakdown by the yeast silencing protein Sir2: Evidence for acetyl transfer from substrate to an NAD breakdown product. *Proc. Natl. Acad. Sci. USA*, 98:415-420.

Tissenbaum H.A. and Guarente L. (2001) Increased dosage of a sir-2 gene extends lifespan in *Caenorhabditis elegans*. *Nature*, 410:227-230.

Vaziri H., West M.D., Allsopp R.C., Davison T.S., Wu Y.S., Arrowsmith C.H., Poirier G.G. and Benchimol S. (1997) ATM-dependent telomere loss in aging human diploid fibroblasts and DNA damage lead to the post-translational activation of p53 protein involving poly(ADP-ribose) polymerase. *EMBO J.*, 16:6018-6033.

Vaziri H., Dessain S.K., Ng-Eaton E., Imai S.I., Frye R.A., Pandita T.K., Guarente L. and Weinberg R.A. (2001) hSIR2SIRT1 functions as an NAD-dependent p53 deacetylase. *Cell*, 107:149-159.

Vogelstein B., Lane D. and Levine A.J. (2000) Surfing the p53 network. *Nature*, 408:307-310.

Yin Y., Terauchi Y., Solomon G.G., Aizawa S., Rangarajan P.N., Yazaki Y., Kadowaki T. and Barrett J.C. (1998) Involvement of p85 in p53-dependent apoptotic response to oxidative stress. *Nature*, 391:707-710.

Yoshida M., Horinouchi S. and Beppu T. (1995) Trichostatin A and trapoxin: novel chemical probes for the role of histone acetylation in chromatin structure and function. *Bioessays*, 5:423-430.

Yu A., Fan H., Lao D., Bailey A.D. and Weiner A.M. (2000) Activation of p53 or loss of the Cockayne syndrome group B repair protein causes metaphase fragility of human U1, U2, and 5S genes. *Mol. Cell*, 5:801-810.

Yu J., Zhang L., Hwang P., Kinzler K. and Vogelstein B. (2001) PUMA induces the rapid apoptosis of colorectal cancer cells. *Mol. Cell*, 7:673-682.

All patents, patent applications, and published references cited herein are hereby incorporated by reference.

A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention. Accordingly, other embodiments are within the scope of the following claims.

5

WHAT IS CLAIMED IS:**CLAIMS**

1. A method of screening a compound, comprising the steps of:
 - (a) providing a reaction mixture comprising Sir2, a transcription factor, and the compound; and
 - (b) determining if the compound modulates Sir2 interaction with the transcription factor, thereby screening the compound.
2. The method of claim 1, wherein the Sir2 interaction with the transcription factor is direct binding, covalent modification in one or both of the Sir2 or transcription factor, a change in cellular location of the test compound, Sir2 or the transcription factor, or an alteration in activity, stability, or structure.
3. The method of claim 2, wherein the determining includes comparing the binding of Sir2 to the transcription factor at a first concentration of the compound and at a second concentration of the compound.
4. The method of claim 3, wherein the first or second concentration of the compound is zero.
5. The method of claim 1, wherein the reaction mixture further comprises a Sir2 cofactor.
6. The method of claim 5, wherein the Sir2 cofactor is NAD or an NAD analog.
7. The method of claim 1 wherein the Sir2 is a Sir2 variant that has reduced deacetylase activity.
8. The methods of claim 1, wherein the Sir2 is human.
9. The method of claim 8, wherein the Sir2 is human SIRT1.
10. The method of claim 1, wherein the Sir2 is murine.
11. The method of claim 10, wherein the Sir2 is murine Sir2 α .

12. The method of claim 1, wherein the Sir2 is exogenous and expressed from a heterologous nucleic acid.
13. The method of claim 1, wherein the transcription factor is exogenous and expressed from a heterologous nucleic acid.
14. The method of claim 1, further comprising the steps of:
 - (c) repeating steps (a) and (b) to confirm a modulatory effect of the compound on Sir2 interaction with the transcription factor, and
 - (d) contacting or administering the compound with or to a cell or animal to evaluate the effect of the compound on the cell or animal.
15. A method of screening a compound, comprising the steps of:
 - (a) providing a reaction mixture comprising Sir2, a transcription factor, and the compound; and
 - (b) determining if the compound modulates Sir2-mediated deacetylation of the transcription factor, thereby screening the compound.
16. The method of claim 15, wherein the determining includes comparing the acetylation status of the transcription factor, at a first concentration of the compound and at a second concentration of the compound.
17. The method of claim 16, wherein the first or second concentration of the compound is zero.
18. The method of claim 17, wherein the reaction mixture further comprises a Sir2 cofactor.
19. The method of claim 18, wherein the Sir2 cofactor is NAD or an NAD analog.
20. The method of claim 15, wherein the Sir2 is a Sir2 variant that has reduced deacetylase activity.
21. The methods of claim 15, wherein the Sir2 is human.
22. The method of claim 21, wherein the Sir2 is human SIRT1.
23. The method of claim 15, wherein the Sir2 is murine.
24. The method of claim 23, wherein the Sir2 is murine Sir2 α .
25. The method of claim 15, wherein Sir2 is exogenous and expressed from a heterologous nucleic acid.

26. The method of claim 15, wherein the transcription factor is exogenous and expressed from a heterologous nucleic acid.
27. The method of claim 15, further comprising the steps of:
- (c) repeating steps (a) and (b) to confirm a modulatory effect of the compound on Sir2-mediated deacetylation of the transcription factor, and
 - (d) contacting or administering the compound with or to a cell or animal to evaluate the effect of the compound on the cell or animal.
28. A method of screening a compound, comprising the steps of:
- (a) providing a compound that interacts with Sir2;
 - (b) contacting the compound with a cell or a system; and
 - (c) determining if the compound modulates transcription of a transcription factor-regulated gene,
- thereby screening the compound.
29. The method of claim 28, wherein the compound binds Sir2 directly.
30. The method of claim 28, wherein the determining includes comparing the modulation of transcription of a transcription factor-regulated gene at a first concentration of the compound and at a second concentration of the compound.
31. The method of claim 30, wherein the first or second concentration of the compound is zero.
32. The method of claim 15, further comprising the steps of:
- (c) repeating steps (a) and (b) to confirm a modulatory effect of the compound on transcription of transcription factor-regulated genes, and
 - (d) contacting or administering the compound with or to a cell or animal to evaluate the effect of the compound on the cell or animal.
33. A method of modifying the acetylation status of a transcription factor binding site on histone or DNA, the method comprising the steps of:
- (a) providing a Sir2-transcription factor complex;
 - (b) allowing the transcription factor to target the Sir2-transcription factor to the transcription factor binding site; and
 - (c) allowing the Sir2 to modify the acetylation status of the transcription factor binding site.

34. The method of claim 33, wherein the method is performed *in vitro* or *in vivo*.
35. The method of claim 34, wherein the method is performed in cell culture..
36. The method of claim 35, wherein the method is performed in an animal.
37. The method of claim 34, wherein the Sir2-transcription factor complex is supplied at concentrations greater than those which occur naturally *in vitro* or *in vivo*.
38. The method of claim 33, wherein the Sir2-transcription factor complex is supplied at a different stage of development than occurs naturally *in vitro* or *in vivo*.
39. The method of claim 33, wherein the Sir2-transcription factor complex is expressed from one or more exogenous genes.
40. The method of claim 33, wherein the Sir2-transcription factor complex is supplied as exogenous Sir2-transcription factor complex.
41. The method of claim 33, wherein the Sir2-transcription factor complex is supplied by inducing endogenous expression of one or more of Sir2 or a transcription factor complex.




Figure 1




Figure 2




Figure 3






Figure 4

**Figure 5**

A



B




Figure 6

A





Figure 7




Figure 8







Figure 9

A**B****C****Figure 10**




Figure 11




Figure 12

p53 harlow -> Translate • 1-frame

DNA sequence 1546 bp atggaggagccg ... actgttgaattc linear

```

1/1                               31/11
atg gag gag ccg cag tca gat cct agc gtc gag ccc oct ctg agt cag gaa aca ttt tca
M E E P Q S D P S V E F P L S Q E T F S
61/21                               91/31
gac cta tgg aaa cta ctt cct gaa aac aac gtt ctg tcc ccc ttg ccg tcc caa gca atg
D L W K L L P E N N V L S P L P S Q A M
121/41                               151/51
gat gat ttg atg ctg tcc ccg gac gat att gaa caa tgg ttc act gaa gac cca ggt cca
D D L M L S P D D I E Q W F T E D P G P
181/61                               211/71
gat gaa gct ccc aga atg cca gag gct gct ccc ccc gtg gcc cct gca cca gca gct cct
D E A P R M P E A A P P V A P A P A A P
241/81                               271/91
aca cag gcg gcc cct gca cca gcc ccc tcc tgg ccc ctg tca tct tct gtc cct tcc cag
T P A A P A P A P S W P L S S S V P S Q
301/101                               331/111
aaa acc tac cag ggc agc tac ggt ttc cgt ctg ggc ttc ttg cat tct ggg aca gcc aag
K T Y Q G S Y G F R L G F L H S G T A K
361/121                               391/131
cct gtg act tgc acg tac tcc cct gcc ctc aac aag atg ttt tgc caa ctg gcc aag acc
S V T C T Y S P A L N K M F C Q L A K T
421/141                               451/151
tgc cct gtg cag ctg tgg gtt gat tcc aca ccc ccc gcc acc cgc gtc cgc gcc atg
C P V Q L W V D S T P P P G T R V R A M
481/161                               511/171
gcc atc tac aag cag tca cag cac atg acg gag gtt gtg agg cgc tgc ccc cac cat gag
A I Y K Q S Q H M T E V V R R C P H H E
541/181                               571/191
cgc tgc tca gat agc gat ggt ctg gcc cct cct cag cat ctt atc cga gtg gaa gga aat
R C S D S D G L A P P Q H L I R V E G N
601/201                               631/211
ttg cgt gtg gag tat ttg gat gac aga aac act ttt cga cat agt gtg gtg gtg ccc tat
L R V E Y L D D R N T F R H S V V V P Y
661/221                               691/231
gag cag cct gag gtt ggc tct gac tgt acc acc atc cac tac aac tac atg tgt aac agt
E P P E V G S D C T T I H Y N Y M C N S
721/241                               751/251
tcc tgc atg ggc ggc atg aac cgg agg ccc atc ctc acc atc atc aca ctg gaa gac tcc
S C M G G M N R R P I L T I I T L E D S
781/261                               811/271
agt ggt aat cta ctg gga cgg aac agc ttt gag gtg cat gtt tgt gcc tgt cct ggg aga
S G N L L G R N S F E V H V C A C P G R
841/281                               871/291
gac cgg cgc aca gag gaa gag aat ctc cgc aag aaa ggg gag cct cac cac gag ctg ccc
D R R T E E E N L R K K G E P H H E L P
901/301                               931/311
cca ggg agc act aag cga gca ctg ccc aac aac acc agc tcc tct ccc cag cca aag aag
P G S T K R A L P N N T S S S P Q P K K
961/321                               991/331
aaa cca ctg gat gga gaa tat ttc acc ctt cag atc cgt ggg cgt gag cgc ttc gag atg
K P L D G E Y F T L Q I R G R E R F E M

```

Figure 13A

p53 harlow -> Translate • 1-frame

```

1021/341      1051/351
ttc cga gag ctg aat gag gcc ttg gaa ctg aag gat gcc cag gct ggg aag gag cca ggg
F R E L N E A L E L K D A Q A G K E F G
1081/361      1111/371
ggg agc agg gct cac tcc agc cac ctg aag tcc aaa aag ggt cag tct acc tcc cgc cat
G S R A H S S H L K S K K G Q S T S R H
1141/381      1171/391
aaa aaa ctg atg ttc aag aca gaa ggg cct gac tca gac tga cat tct cca ctt ctt gtt
K K L M F K T E G P D S D * K S P L L V
1201/401      1231/411
ccc cac tga cag cct ccc acc ccc atc tct ccc tcc cct gcc att ttg ggt ttt ggg tct
P H * Q P P T P I S P S P A I L G F G S
1261/421      1291/431
ttg aac cct tgc ttg caa tag gtg tgc gtc aga agc acc cag gac ttc cnt ttg ctt tgt
L N P C L Q * V C V R S T Q D F H L L C
1321/441      1351/451
ccc ggg gct cca ctg aac aag ttg gcc tgc act ggt gtt ctg ttg tgg gga gga gga tgg
P G A P L N K L A C T G V L L W G G G W
1381/461      1411/471
gga gta gga cat acc agc tta gat ttt aag gtt ttt act gtg agg gat gtt tgg gag atg
G V G H T S L D F K V F T V R D V W E M
1441/481      1471/491
taa gaa atg ttc ttg cag tta agg gtt agt tta caa tca gcc aca ttc tag gta ggg acc
* E M F L Q L R V S L Q S A T F * V G T
1501/501      1531/511
cac ttc acc gta cta acc agg gaa gct gtc cct cac tgt tga att c
H F T V L T R E A V P H C * I

```

Figure 13B

BASE COUNT 401 a 513 c 460 g 386 t
 ORIGIN 1 bp upstream of Sall site; Chromosome 17p13 [Unpublished (1985) C.
 1 gtcgaccctt tccacccctg gaagatggaa ataaacctgc gtgtgggtgg agtgttagga
 61 caaaaaaaaaa aaaaaaaaaaag tctagagcca ccgtccaggg agcaggtagc tgcitgggctc
 121 cggggacact ttgcgttcgg gcitgggagcg tgcitccac gacggtgaca cgcttccctg
 181 gatggcagc cagactgcct tccgggtcac tgcctggag gacccgcagt cagatccctag
 241 cgtcgagccc cctctgagtc aggaacatt ttcagacctc tggaaactac ttctgaaaa
 301 caacgttcgt tccccctgc cgtcccaagc aatggatgat ttgatgctgt ccccgacga
 361 tatgaacaa tggttactg aagacccagg tccagatgaa gctccagaa tgcagagggc
 421 tgcctcccc cgtggccctg caccagcagc tccacaccg gcggccctg caccagcccc
 481 cctctggccc ctgtacatt ctgtcccttc ccagaaaacc taccagggca gctacgggtt
 541 ccgtctgggc ttctgcatt ctgggacagc caagtctgt actgcacgt actccctgc
 601 cctcaacaag atgtttgccc aactggccaa gacctgcct gtgcagctgt ggggtgatic
 661 cacacccccg cccggcaccg gcgtccgcgc catggccatc tacaagcagt cacagcacat
 721 gacggagggt gtgaggcgct gccccacca tgagcgctgc tcagatagcg atggtctggc
 781 cctctctcag catctatcc gagtggagg aaatttgcgt gtggagtatt tggatgacag
 841 aaacactttt cgacatagtg tgggtgtgcc ctatgagccg cctgagggtg gctctgactg
 901 taccaccatc cactacaact acatgtgtaa cagtctctgc atgggcggca tgaaccggag
 961 gcccacctc accatcatca cactggaaga ctccagtgtt aatctactgg gacggaacag
 1021 ctgtgagggt catgtttgt cctgtctgg gagagaccg cgcacagagg aagagaatct
 1081 ccgcaagaaa ggggagcctc accacgagct gccccaggg agcactaagc gagcactgcc
 1141 caacaacacc agctctctc cccagccaaa gaagaaacca ctggatggag aatatttacc
 1201 ccttcagatc cgtgggcgtg agcgcttca gatgtccga gagctgaatg aggccttggg
 1261 actcaaggat gccaggctg ggaaggagcc aggggggagc agggctcact ccagccacct
 1321 gaagtccaaa aagggtcagt ctacctccg ccataaaaaa ctcatgttca agacagaagg
 1381 gccigactca gactgacatt ctccactct tgttccccc tgcagacctc ccacccccat
 1441 ctctccctcc cctgccatt tgggtttgg gctttgaac cctgtctgc aatagggtgt
 1501 cgtcagaagc acccaggact tccatttct tgcctggg gctccactga acaagttggc
 1561 ctgactgggt gttttgtgt ggggaggagg atggggagta ggacatacca gcttagattt
 1621 taagggtttt actgtgagg atgtttggga gatgaagaa atgttctgc agttaagggt
 1681 tagtttaca tccagccat tctaggtagg gacccactc accgtactaa ccaggaagc
 1741 tgcctcac tgtgaattc

Figure 14

```

      10      20      30      40      50      60
GCGGAGCAGAGAGGCGAGGGCGGAGGGGCCAGAGAGGCAGTTGGAAGATGGCGGACGAGG
                                M A D E V

      70      80      90      100     110     120
TGGCGCTCGCCCTTCAGGCCGCGGGCTCCCTTCGCGCGCGGCCCGCCATGGAGGCCCGGT
  A L A L Q A A G S P S A A A A M E A A S

      130     140     150     160     170     180
CGCAGCCCGGCGGACGAGCCGCTCCGCAAGAGGCCCGCGAGACGGGCCTGGCCTCGGGC
  Q P A D E P L R K R P R R D G P G L G R

      190     200     210     220     230     240
GCAGCCCGGGCGAGCCGAGCGCAGCAGTGGCGCGCGCGCGCGGGTGTGAGCGCGCGA
  S P G E P S A A V A P A A A G C E A A S

      250     260     270     280     290     300
CGCGCCCGCGCCCGCGCGCGCTGTGGCGGGAGGCGGCAGGGCGCGCGCGAGCGCGGAGC
  A A A P A A L W R E A A G A A A S A E R

      310     320     330     340     350     360
GGGAGGCCCGCGCGACCGCCGTGGCGCGGGACGGAGACAATGGGTCCGGCCTCGCGCGGG
  E A P A T A V A G D G D N G S G L R R E

      370     380     390     400     410     420
AGCCGAGGGCGCGCTGACGACTTCGACGACGACGAGGGCGAGGAGGAGGACGAGGCGCGG
  P R A A D D F D D D E G E E E D E A A A

      430     440     450     460     470     480
CGGCAGCGGGCGGGCGGCGAGCGATCGGCTACCGAGACAACCTCCTGTGACCGATGGACTCC
  A A A A A A I G Y R D N L L L T D G L L

      490     500     510     520     530     540
TCACTAATGGCTTTTCATTCCTGTGAAAGTGATGACGATGACAGAACGTCACACGCCAGCT
  T N G F H S C E S D D D D R T S H A S S

      550     560     570     580     590     600
CTAGTGACTGGACTCCGCGGCCGCGGATAGGTCATATACITTTTGTTCAGCAACATCTCA
  S D W T P R P R I G P Y T F V Q Q H L M

      610     620     630     640     650     660
TGATTGGCACCGATCCTCGAACAATTCTTAAAGATTATTACCAGAAACAATTCTCCAC
  I G T D P R T I L K D L L P E T I P P P

      670     680     690     700     710     720
CTGAGCTGGATGATATGACGCTGTGGCAGATTTGTTATTATATCCTTTTCAGAACCCACAA
  E L D D M T L W Q I V I N I L S E P P K

      730     740     750     760     770     780
AGCGGAAAAAAGAAAAGATATCAATACAAATTGAAGATGCTGTGAAGTTACTGCAGGAGT
  R K K R K D I N T I E D A V K L L Q E C

      790     800     810     820     830     840
GTAAAAAGATAATAGTTCTGACTGGAGCTGGGGTTTCTGTCTCCTGTGGGATTCTGACT
  K K I I V L T G A G V S V S C G I P D F

```

Figure 15A

850 860 870 880 890 900
 TCAGATCAAGAGACCGGTATCTATGCTCGCCTTGCGGTGGACTTCCCAGACCTCCCAGACC
 R S R D G I Y A R L A V D F P D L P D P
 910 920 930 940 950 960
 CTCAGGCCATGTTTGATATTGAGTATTTTAGAAAAGACCCPAGACCAITTCITCAAGTTTG
 Q A M F D I E Y F R K D P R P F F K F A
 970 980 990 1000 1010 1020
 CAAAGGAAATATATCCCGGACAGTTCCAGCCGTCTCTGTGTGCACAAATTCATAGCTTTGT
 K E I Y P G Q F Q P S L C H K F I A L S
 1030 1040 1050 1060 1070 1080
 CAGATAAGGAAGGAAACTACTTTCGAAATTATACTCAAAATATAGATACCTTCGAGCAGG
 D K E G K L L R N Y T Q N I D T L E Q V
 1090 1100 1110 1120 1130 1140
 TTGCAGGAATCCAAAGGATCCCTTCAGTGTCTATGGTTCTTTTGCAACAGCATCTTGCCTGA
 A G I Q R I L Q C H G S F A T A S C L I
 1150 1160 1170 1180 1190 1200
 TTTGTAAATACAAAGTTGATTGTGAAGCTGTTTCGTGGAGACATTTTAAATCAGGTAGTTTC
 C K Y K V D C E A V R G D I F N Q V V P
 1210 1220 1230 1240 1250 1260
 CTCGGTGCCTTAGGTGCCAGCTGATGAGCCACTTGCCATCATGAAGCCAGAGATTGTCT
 R C P R C P A D E P L A I M K P E I V F
 1270 1280 1290 1300 1310 1320
 TCTTTGGTGAAGAACTTACCAGAACAGTTTCATAGAGCCATGAAGTATGACAAAGATGAAG
 F G E N L P E Q F H R A M K Y D K D E V
 1330 1340 1350 1360 1370 1380
 TTGACCTCCCTCATTTGTTATTTGGATCTTCTCTGAAAGTGAGACCAGTAGCACTAATTCCAA
 D L L I V I G S S L K V R P V A L I P S
 1390 1400 1410 1420 1430 1440
 GTTCTATACCCCATGAAGTGCCTCAAAATATTAAATAAATAGGGAACCTTTGCCTCATCTAC
 S I P H E V P Q I L I N R E P L P H L H
 1450 1460 1470 1480 1490 1500
 ATTTTGATGTAGAGCTCCTTGGAGACTGCGATGTTTAAATTAATGAGTTGTGTGCATAGGC
 F D V E L L G D C D V I I N E L C H R L
 1510 1520 1530 1540 1550 1560
 TAGGTGGTGAATATGCCAAACTTTGTTGTAACCCCTGTAAAGCTTTTCAGAAATTACTGAAA
 G G E Y A K L C C N P V K L S E I T E K
 1570 1580 1590 1600 1610 1620
 AACCTCCACGCCCAACAAAGGAATTGGTTTCATTTATCAGAGTTGCCACCAACACCTCTTC
 P P R P Q K E L V H L S E L P P T P L H
 1630 1640 1650 1660 1670 1680
 ATATTTTCGGAAGACTCAAGTTTCACCTGAAAGAACTGTACCACAAGACTCTTCTGTGATTG
 I S E D S S S P E R T V P Q D S S V I A
 1690 1700 1710 1720 1730 1740
 CTACACTTGTAGACCAAGCAACAAACAAATGTTAATGATTTAGAAGTATCTGATCAA
 T L V D Q A T N N N V N D L E V S E S S

Figure 15B

1750 1760 1770 1780 1790 1800
GTTGCTGCGGAGAAAACCACAGAGTACAGACTAGTAGGAATGTTGAGAACATTAAATG
C V E E K P Q E V Q T S R N V E N I N V

1810 1820 1830 1840 1850 1860
TGGAAAATCCAGATTTTAAGGCTGTTGGTTCCAGTACTGCAGACAAAATGAAAGAACTT
E N P D F K A V G S S T A D K N E R T S

1870 1880 1890 1900 1910 1920
CAGTTGCAGAAACAGTGCAGAAAATGCTGGCCTAATAGACTTGCAGAGGAGCAGATTAGTA
V A E T V R K C W P N R L A K E Q I S K

1930 1940 1950 1960 1970 1980
AGCGGCTTGAGGGTAATCAATACCTGTTTGTACCACCAATCGTTACATATTCCACGGTG
R L E G N Q Y L F V P P N R Y I F H G A

1990 2000 2010 2020 2030 2040
CTGAGGTATACTCAGACTCTGAAGATGACGCTCTTGTCTCTTAGTTCCTGTGGCAGTAACA
E V Y S D S E D D V L S S S S C G S N S

2050 2060 2070 2080 2090 2100
GTGACAGTGGCACATGCCAGAGTCCAGTTTAGAAGAACCCCTTGGAGATGAAAGTCAAA
D S G T C Q S P S L E E P L E D E S E I

2110 2120 2130 2140 2150 2160
TTGAAGAATTCTACATATGGCTTGGAGATGATACGGAGAGGCCCGAATGTGCTGGAGGAT
E E F Y N G L E D D T E R P E C A G G S

2170 2180 2190 2200 2210 2220
CTGGATTTGGAGCTGATGGAGGGGATCAAGAGGTTGTTAATGAAGCTATAGCTACAAGAC
G F G A D G G D Q E V V N E A I A T R Q

2230 2240 2250 2260 2270 2280
AGGAATTGACAGATGTAACTATCCATCAGACAAATCATAACACTATTGAAGCTGTCCGG
E L T D V N Y P S D K S *

2290 2300 2310 2320 2330 2340
ATTCAGGAATTGCTCCACCAGCATTTGGGAACCTTTAGCATGTCAAAAAATGAATGTTTAC

2350 2360 2370 2380 2390 2400
TTGTGAACCTTGAACAAGGAAATCTGAAAGATGTATTATTATAGACTGGAAAATAGATTG

2410 2420 2430 2440 2450 2460
TCTTCTTGGATAATTTCTAAAGTTCCATCATTTCTGTTTGTACTTGTACATTCAACACTG

2470 2480 2490 2500 2510 2520
TTGGTTGACTTCATCTTCCTTTCAAGGTTTCATTTGTATGATACATTTCGTATGTATGTATA

2530 2540 2550 2560 2570 2580
ATTTTGTTTTTTGCCTAATGAGTTTCAACCTTTTAAAGTTTTTCAAAGCCATTGGAATGT

2590 2600 2610 2620 2630 2640
TAATGTAAGGGGACAGCTTATCTAGACCAAGAAATGGTATTTTCACTTTTTTGTGTGT

2650 2660 2670 2680 2690 2700
AACATTGADTAGTTTAAAGCCCTCAATTTCTGTTCTGCTGAACTTTATTTTTAGGACAG



2710 2720 2730 2740 2750 2760
TTAACTTTTTTAAACACTGGCATTTTCCAAAACCTGTGGCAGCTAACTTTTTTAAATCACA

Figure 15C

2770 2780 2790 2800 2810 2820
 GATGACTTGTAAATGTGAGGAGTCAGCACCGTGTCTGGAGCACTCAAAAACCTTGGGCTCAGT
 2830 2840 2850 2860 2870 2880
 GTGTGAAGCGTACTTACTGCATCGTTTTTGTACTTGCTGCAGACGTGGTAATGTCCAAAC
 2890 2900 2910 2920 2930 2940
 AGGCCCCCTGAGACTAATCTGATAAATGATTTGGAAATGTGTTCAGTTGTTCTAGAAACA
 2950 2960 2970 2980 2990 3000
 ATAGTGCCCTGTCTATATAGTCCCCCTTAGTTTTGAATATTTGCCATTGTTTTAATTAAATAC
 3010 3020 3030 3040 3050 3060
 CTATCACTGTGGTAGAGCCGTGCATAGATCTTCACCACAAATACTGCCAAGATGTGAATAT
 3070 3080 3090 3100 3110 3120
 GCAAAGCCCTTTCTGAATCTAATAATGGTACTTCTACTGGGGAGAGTGTAAATATTTTGGAC
 3130 3140 3150 3160 3170 3180
 TGCTGTTTTTCCATTAAATGAGGAAAGCAATAGGCCTCTTAATTAAAGTCCCAAAGTCATA
 3190 3200 3210 3220 3230 3240
 AGATAAATTGTAGCTCAACCAGAAAGTACACTGTTGCCTGTTGAGGATTTGGTGTAATGT
 3250 3260 3270 3280 3290 3300
 ATCCCCAAGGTGTTAGCCTTGATATATGGAGATGAATACAGATCCAATAGTCAAATGAAAC
 3310 3320 3330 3340 3350 3360
 TAGTTCCTTAGTTATTTAAAAGCTTAGCTTGCCTTAAACTAGGGATCAATTTTCTCAACT
 3370 3380 3390 3400 3410 3420
 GCAGAAACCTTTAGCCCTTCAAAACAGTTTCACACCTCAGAAAGTCAGTATTTATTTTACAG
 3430 3440 3450 3460 3470 3480
 ACTTCCTTTGGAACATTGCCCCCAAATTTAAATATTCATGTGGGTTTAGTATTTATTACAA
 3490 3500 3510 3520 3530 3540
 AAAAAATGATTTGAAATATAGCTGTTCTTTATGCATAAAATACCCAGTTAGGACCATTACT
 3550 3560 3570 3580 3590 3600
 GCCCAGGAGAAAGTATTAAAGTAGCTCATTTCCCTACCTAAAAGATAACTGAATTTATT
 3610 3620 3630 3640 3650 3660
 TGGCTACACTAAAGAATGCAGTATATTTAGTTTTCATTTCATGATGTGTTTGTGCTAT
 3670 3680 3690 3700 3710 3720
 AGACAAATATTTTAAATTGAAAAATTTGTTTTTAATTTATTTTACAGTGAAGACTGTTTTTC
 3730 3740 3750 3760 3770 3780
 AGCTCTTTTATATTGTACATAGACTTTTATGTAATCTGGCATATGTTTTGTAGACCGTT
 3790 3800 3810 3820 3830 3840
 TAATGACTGGAATTATCTTCCTCCAACCTTTTGAATACAAAAACAGTGTTTTATACTAAAA
 3850 3860 3870
 AAAAAAAAAGTCGACGCGGCCGCGAATTC

Figure 15D

NCBI Sequence Viewer

PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM

Search for

Limits Index History Clipboard

as

☐ 1: [AF214646](#) **Mus musculus Sir2alpha protein (Sir2alpha) mRNA, complete cds** PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

LOCUS	AF214646	3849 bp	mRNA	ROD	26-OCT-2000
DEFINITION	Mus musculus Sir2alpha protein (Sir2alpha) mRNA, complete cds.				
ACCESSION	AF214646				
VERSION	AF214646.1 GI:6693710				
KEYWORDS	.				
SOURCE	house mouse.				
ORGANISM	<u>Mus musculus</u>				
	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Mus.				
REFERENCE	1 (bases 1 to 3849)				
AUTHORS	Imai, S., Armstrong, C.M., Kaerberlein, M. and Guarente, L.				
TITLE	Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase				
JOURNAL	Nature 403 (6771), 795-800 (2000)				
MEDLINE	20155478				
PUBMED	10693811				
REFERENCE	2 (bases 1 to 3849)				
AUTHORS	Imai, S., Armstrong, C.M. and Guarente, L.				
TITLE	Direct Submission				
JOURNAL	Submitted (10-DEC-1999) Dept. of Biology, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA				
FEATURES	Location/Qualifiers				
source	1..3849				
	/organism="Mus musculus"				
	/strain="Swiss Webster/NIH"				
	/db_xref="taxon:10090"				
	/dev_stage="15-day embryo"				
gene	1..3849				
	/gene="Sir2alpha"				

Figure 16A

NCBI Sequence Viewer

CDS 48..2261
 /gene="Sir2alpha"
 /function="have NAD-dependent histone deacetylase and putative mono-ADP-ribosyltransferase activities"
 /note="closest homolog to the yeast silencing and aging protein, Sir2p"
 /codon_start=1
 /product="Sir2alpha protein"
 /protein_id="AAF24983.1"
 /db_xref="GI:6693711"
 /translation="MADEVALLALQAAGSPSAAAAMEAASQPADEPLRKRPRRDGPGLGRSPGEPSAAVAPAAAGCEAASAAAPALWREAAGAAASAEREAPATAVAGDGDNGSGLRREPRAADDFDDDEGEEDDEAAAAAIAIGYRDNLLLTGGLLTNGFHSCESDDDDRTHASSSDWTFRPRIGPYTFVQQHLMIGTDPRTILKDLLPETIPPPPELDDMTLWQIVINILSEPPKRKRKRDINTIEDAVKLLQECKKIIVLTGAGVSVSCGIPDFRSRDGIYARLAYTQNIIDTLEQVAGIQRILOCHGSFATASCLICKYKVDCEAVRGDI FNQVVRCPRCPADEPLAIMKPEIVFFGENLPEQFHRAMKYDKDEVDLLIVIGSSLKVRPVALIPSSIPHEVPQILINREPLPHLHFDVELLGDGCDVI INELCHRLGGEYAKLCCNPVKLSEITEKPPRPQKELVHLSELPPFTPLHISEDSSSPERTVFPQDSSVIATLVDQATNNNVNDLEVSSESSVEEKPQEVQTSRNVENINVENPDFKAVGSSSTADKNERTSVAETVRKCWPNRLAKEQISKRLGNQYLFVPPNRYIFHGAEVYSDSEDDVLSSSSCGSNSDSGTCQSPSLEEPLDESEIEEFYNGLEDDTERPECAGGSGFGADGGDQEVVNEAIATRQELTDVNYPDSKS"

BASE COUNT 1118 a 761 c 889 g 1081 t
 ORIGIN

1	gcggagcaga	ggaggcgagg	gcggagggcc	agagaggcag	ttggaagatg	gcggacgagg
61	tggcgctcgc	ccttcaggcc	gcgggctccc	cttcgcggc	ggccgccatg	gaggccgcgt
121	cgcagccggc	ggacgagccg	ctccgcaaga	ggccccgcg	agacgggcct	ggcctcgggc
181	gcagcccggg	cgagccgagc	gcagcagtgg	cgccggcggc	cgcggggtgt	gaggcgccga
241	gcgccgcggc	cccggcgggc	ctgtggcggg	aggcggcagg	ggcgccggcg	agcgccgagc
301	gggaggcccc	ggcgacggcc	gtggccgggg	acggagacaa	tggttcgggc	ctgcggcggg
361	agccgagggc	ggctgacgac	ttcgacgacg	acgagggcga	ggaggaggac	gaggcgccgg
421	cggcagcggc	ggcggcagcg	atcggtacc	gagacaaact	cctgttgacc	ctgtggacct
481	tcactaatgg	ctttcattcc	tgtgaaagtg	atgacgatga	cagaacgtca	cacgccagct
541	ctagtgactg	gactccgcgg	ccgcggatag	gtccatatac	ttttgttcag	caacatctca
601	tgattggcac	cgatccctcg	acaattctta	aagatttatt	accagaaaac	attcctccac
661	ctgagctgga	tgatatgacg	ctgtggcaga	ttgttattaa	tatcctttca	gaaccaccaa
721	agcggaaaaa	aagaaaagat	atcaatacaa	ttgaagatgc	tgtgaagtta	ctgcaggagt
781	gtaaaaagat	aatagttctg	actggagctg	gggtttctgt	ctcctgtggg	attcctgact
841	tcagatcaag	agacggtatc	tatgctcgcc	ttgcggtgga	cttcccagac	ctcccagacc
901	ctcaagccat	gtttgatatt	gagtatttta	gaaaagaccc	aagaccattc	ttcaagtttg
961	caaaggaaat	atatcccggg	cagttccagc	cgtctctgtg	tcacaaatc	atagctttgt
1021	cgataaagga	aggaaaacta	cttcgaaatt	atactcaaaa	tatagatacc	ttggagcagg
1081	ttgcaggaat	ccaaaggatc	cttcagtgtc	atggttcctt	tgcaacagca	tcttgccctga
1141	tttgtaaata	caaagttgat	tgtgaagctg	ttcgtggaga	catttttaat	caggtagttc
1201	ctcggtgccc	taggtgcccc	gotgatgagc	cacttgccat	catgaagcca	gagattgtct
1261	tctttggtga	aaacttacca	gaacagtttc	atagagccat	gaagtatgac	aaagatgaag
1321	ttgacctcct	cattgttatt	ggatcttctc	tgaaagtgag	accagtagca	ctaattccaa
1381	gttctatacc	cctatgaagtg	cctcaaatat	taataaatag	ggaacctttg	cctcatctac
1441	attttgatgt	agagctcctt	ggagactgcg	atgttataat	taatgagttg	tgtcataggg
1501	taggtgggtga	atatgccaaa	ctttgttgta	accctgtaaa	gctttcagaa	attactgaaa
1561	aacctccacg	cccacaaaag	gaattggttc	atztatcaga	gttgccacca	acacctcttc

Figure 16B

NCBI Sequence Viewer

```




1621 atattttcggga agactcaagt tcacctgaaa gaactgtacc acaagactct tctgtgattg
1681 ctacacttgt agaccaagca acaacaaca atgttaatga tttagaagta tctgaatcaa
1741 gttgtgttga agaaaaacca caagaagtac agactagtag gaatgttgag aacattaatg
1801 tggaaaaatcc agatttttaag gctgttggtt ccagtactgc agacaaaaat gaaagaactt
1861 cagttgcaga aacagtggga aaatgctggc ctaatagact tgcaaaggag cagattagta
1921 agcggcttga gggtaatcaa tacctgtttg taccacaaa tcgttacata ttccacgggtg
1981 ctgagggtata ctacagactct gaagatgacg tcttgtcctc tagttcctgt ggcagtaaca
2041 gtgacagtgg cacatgccag agtccaagtt tagaagaacc cttggaagat gaaagtgaaa
2101 ttgaagaatt ctacaatggc ttggaagatg atacggagag gcccgaaatgt gctggaggat
2161 ctggatttgg agctgatgga ggggatcaag aggttggtta tgagctata gctacaagac
2221 aggaattgac agatgtaaac tatccatcag acaaatcata aactattga agctgtccgg
2281 attcaggaat tgctccacca gcattgggaa ctttagcatg tcaaaaaaat gaatgtttac
2341 ttgtgaactt gaacaaggaa atctgaaaga tgtattattt atagactgga aaatagattg
2401 tcttcttggga taatttctaa agttccatca tttctgtttg tacttgtaaa ttcaacactg
2461 ttggttgact tcattcttct ttcaagggtt atttgtatga tacattcgta tgtatgtata
2521 attttgtttt ttgcctaatt agtttcaacc ttttaaagtt ttcaaaagcc attggaatgt
2581 taatgtaaag ggaacagctt atctagacca aagaatggta tttcacactt tttgtttgt
2641 aacattgaat agttttaaag cctcaatttc tgttctgctg aacttttatt tttaggacag
2701 ttaacttttt aaacactggc attttccaaa acttgtggca gctaactttt taaaatcaca
2761 gatgacttgt aatgtgagga gtcagcaccg tgtctggagc actcaaaact tgggctcagt
2821 gtgtgaagcg tacttactgc atcgtttttg taettgctgc agacgtggta atgtccaaac
2881 agggccctga gactaatctg ataaatgatt tggaaatgtg tttcagttgt tctagaaaca
2941 atagtgcctg tctatatagg tccccttagt ttgaatattt gccattgttt aattaaatc
3001 ctatcactct aataaagcct gcatagatct tcaccacaaa tactgccaa agtggaatat

3061 gcaaaagcctt tctgaatcta ataattggtac ttctactggg gagagtgtaa tattttggac
3121 tgctgttttt ccattaatga ggaaagcaat aggcctctta attaaagtcc caaagtcata
3181 agataaatfg tagctcaacc agaaagtaca ctgttgccctg ttgaggattt ggtgtaatgt
3241 atcccaaggt gttagccttg tattatggag atgaatacag atccaatagt caaatgaaac
3301 tagttcttag ttatttaaaa gcttagcttg ccttaaaact agggatcaat tttctcaact
3361 gcagaaactt ttagcctttc aaacagttca cacctcagaa agtcagtatt tattttacag
3421 acttcttttg aacattgccc ccaaatttaa atattcatgt gggtttagta tttattacaa
3481 aaaaatgatt tgaaatatag ctgttcttta tgcataaaat acccagttag gaccattact
3541 gccagaggag aaaagtatta agtagctcat ttccctacct aaaagataac tgaatttatt
3601 tggctacact aaaggaatgca gtatatattag ttttccattt gcatgatgtg tttgtgctat
3661 agacaatatt ttaaattgaa aaatttgttt taaattattd ttacagtga gactgttttc
3721 agctcttttt atattgtaca tagactttta tgtaatctgg catatgtttt gtagaccgtt
3781 taatgactgg attatcttcc tocaactttt gaaatacaaa aacagtgttt tataactaaa
3841 aaaaaaaaaa

```

Figure 16C

NCBI Sequence Viewer

PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM

Search ☒ Nucleotide for

as

☐ 1: [AF083107](#) *Homo sapiens sirtuin type 2 (SIRT2) mRNA, complete cds* PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

LOCUS	AF083107	1963 bp	mRNA	PRI	21-MAR-2001
DEFINITION	Homo sapiens sirtuin type 2 (SIRT2) mRNA, complete cds.				
ACCESSION	AF083107				
VERSION	AF083107.2 GI:13400019				
KEYWORDS	.				
SOURCE	human.				
ORGANISM	<u>Homo sapiens</u> Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.				
REFERENCE	1 (bases 1 to 1963)				
AUTHORS	Frye, R.A.				
TITLE	Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity				
JOURNAL	Biochem. Biophys. Res. Commun. 260 (1), 273-279 (1999)				
MEDLINE	99310604				
REFERENCE	2 (bases 1 to 1963)				
AUTHORS	Frye, R.A.				
TITLE	Direct Submission				
JOURNAL	Submitted (10-AUG-1998) Pathology, VA Med. Cent. (132L), Univ. of Pittsburgh, University Drive C, Pittsburgh, PA 15240, USA				
COMMENT	On Mar 21, 2001 this sequence version replaced gi: 5225319 .				
FEATURES	Location/Qualifiers				
source	1..1963 /organism="Homo sapiens" /db_xref="taxon:9606" /chromosome="19" /map="19q13" /tissue_type="testis"				
gene	1..1963 /gene="SIRT2"				
CDS	201..1370 /gene="SIRT2" /note="yeast silent information regulator 2 (Sir2p) homolog" /codon_start=1 /product="sirtuin type 2" /protein_id="AAD40850.2" /db_xref="GI:13400020"				

Figure 17A

NCBI Sequence Viewer




```

/translation="MAEPDPSHPLETQAGKVQEAQDSDSSEGGGAAGGEADMDFLRNL
FSQTLNLSGSKERLLDELTLBGVARYMQSERCRRVICLVGAGISTSAGIPDFRSPSTG
LYDNLEKYHLPYPEAIFEISYFKKHPEPFFALAKELYPGQFKPTICHYFMRLLKDKGL
LLRCYTONIDTLERIAGLEQEDLVEAHGTFTYTSHCVSASCRHEYPLSWMKEKIFSEVT
PKCEDCQSLVKPDIVFFGESLPARFFSCMQSDFLKVDLLVMGTSLQVQPFASLISKÅ
PLSTPRLINKEKAGQSDPFLGMIMGLGGGMDFDSSKKAYRDVAWLGECDQGCLALAEI
LGWKKLELDLVRREHASIDAQSGAGVNPSTSSASPKKSPPPAKDEARTTEREKPQ"
BASE COUNT      432 a      616 c      541 g      374 t
ORIGIN
1  gtgttgtaacg aaagcgcgctc tgcggccgca atgtctgctg agagttgtag ttctgtgccc
61  tatcacggcc actcccattt ctggtgccgt cacgggacag agcagtcggt gacaggacag
121 agcagtcggt gacgggacac agtggttggt gacgggacag agcggtcggt gacagcctca
181 agggcttcag caccgcgccc atggcagagc cagacccctc tcaccctctg gagaccagg
241 cagggaaggt gcaggaggct caggactcag attcagactc tgaggaggga gccgctggtg
301 gagaagcaga catggacttc ctgcggaact tattctccca gacgctcagc ctgggcagcc
361 agaaggagcg tctgctggac gagctgacct tgggaaggggt ggcccgggtac atgcagagcg
421 aacgctgtcg cagagtcatc tgtttggtgg gagctggaat ctccacatcc gcaggcatcc
481 ccgactttcg ctctccatcc accggcctct atgacaacct agagaagtac catcttccct
541 acccagaggc catctttgag atcagctatt tcaagaaaca tccggaacct ttcttcgccc
601 tcgccaagga actctatcct gggcagttca agccaacct ctgtcactac ttctgcgcc
661 tgctgaagga caaggggcta ctctgcgctc gctacacgca gaacatagat aacctggagc
721 gaatagccgg gctggaacag gaggacttgg tggaggcgca cggcaccttc tacacatcac
781 actgcgtcag cgccagctgc cggcacgaat acccgctaag ctggatgaaa gagaagatct
841 tctctgaggt gacgcccagg tgtgaagact gtcagagcct ggtgaagcct gatatcgtct
901 tttttggtga gagcctccca gcgcgtttct tctcctgtat gcagtcagac ttctgaagg
961 tggacctcct cctggtcatg ggtacctctc tgcaggtgca gccctttgac tccctcatca
1021 gcaaggcacc cctctccacc cctcgccctc tcatcaacaa ggagaaagct ggccagtcgg
1081 accctttcct ggggatgatt atgggcctcg gaggaggcat ggactttgac tccaagaagg
1141 cctacaggga cgtggcctgg ctgggtgaat gcgaccaggg ctgcctggcc cttgctgagc
1201 tccttgatg gaagaaggag ctggaggacc ttgtccggag ggagcacgcc agcatagatg
1261 ccagtcggg ggcgggggtc cccaacccca gcacttcagc ttcccccaag aagtccccgc
1321 cacctgcca ggacgaggcc aggacaacag agagggagaa accccagtga cagctgcac
1381 tccaggcg gâtggcgagc tctcaggga cagctgagcc ccaaccgggc ctggcccccct
1441 cttaaccagc agttcttgtc tggggagctc agaacatccc ccaatctctt acagctccct
1501 cccaaaact ggggtcccag caaccctggc ccccaacccc agcaaattct taacacctcc
1561 tagaggccaa ggcttaaaaca ggcattctta ccagccccac tgtctctaac cactcctggg
1621 ctaaggagta acctccctca tctctaactg ccccaacggg gccagggcta cccagaact
1681 ttttaactct ccaggacagg gagcttcggg ccccaactct gtctcctgcc ccggggggcc
1741 tgtggctaag taaaccatac ctaacctacc ccagtgtggg tgtgggcctc tgaatataac
1801 ccacaccagc cgtaggggga gtctgagccg ggagggtcc cgagtctctg ccttcagctc
1861 ccaaagtggg tgggtgggcc ccttcacgtg ggacccactt cccatgctgg atgggcagaa
1921 gacattgctt attggagaca aattaaaaaac aaaaacaact aac

```

//

Figure 17B

PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM

Search ☒ Nucleotide for

as

☒ 1: AF083106 **Homo sapiens sirtuin type 1 (SIRT1) mRNA, complete cds** PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

LOCUS	AF083106	4086 bp	mRNA	PRI	14-APR-2000
DEFINITION	Homo sapiens sirtuin type 1 (SIRT1) mRNA, complete cds.				
ACCESSION	AF083106				
VERSION	AF083106.2 GI:7555470				
KEYWORDS	.				
SOURCE	human.				
ORGANISM	Homo sapiens				
	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.				
REFERENCE	1 (bases 1 to 4086)				
AUTHORS	Frye,R.A.				
TITLE	Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity				
JOURNAL	Biochem. Biophys. Res. Commun. 260 (1), 273-279 (1999)				
MEDLINE	99310604				
REFERENCE	2 (bases 1 to 4086)				
AUTHORS	Frye,R.A.				
TITLE	Direct Submission				
JOURNAL	Submitted (10-AUG-1998) Pathology, VA Med. Cent. (132L), Univ. of Pittsburgh, University Drive C, Pittsburgh, PA 15240, USA				
REFERENCE	3 (bases 1 to 4086)				
AUTHORS	Frye,R.A.				
TITLE	Direct Submission				
JOURNAL	Submitted (14-APR-2000) Pathology, VA Med. Cent. (132L), Univ. of Pittsburgh, University Drive C, Pittsburgh, PA 15240, USA				
REMARK	Sequence update by submitter				
COMMENT	On Apr 14, 2000 this sequence version replaced gi:5225317 .				
FEATURES	Location/Qualifiers				
source	1..4086				
	/organism="Homo sapiens"				
	/db_xref="taxon:9606"				
	/tissue_type="testis"				
gene	1..4086				
	/gene="SIRT1"				
CDS	54..2297				
	/gene="SIRT1"				
	/note="yeast silent information regulator 2 (Sir2p) homolog"				
	/codon_start=1				
	/product="sirtuin type 1"				
	/protein_id="AAD40849.2"				
	/db_xref="GI:7555471"				
	/translation="MADEAALALQPGGSPSAAGADREAASSPAGEPLRKPRRDGPGLERSPGEPGGAAPEREPVPAARGCPGAAAAALWREAEEAAAAGGEQEAQATAAAGEGD"				

Figure 18A

NGPGLQGSPREPPLADNLYDEDDDDDEGEEEEAAAAAIGYRDNLLFGDEIITNGFHSC
 ESDEEDRA\$HASSSDWTPRPRIGPYTFVQOHLMI GTDPRTILKDLLPETIPPPELDDM
 TLWQIVINILSEPPKRRKRDINTIEDAVKLLQECKKIIIVLTGAGVSVSCGIPDFR\$R
 DGIYARLAVDFPDLDPQAMFDIEYFRKDFRPF\$KFAKEIYPGQFQPSLCHKFIALSD
 KEGKLLRNYTQNI DTLEQVAGIQRI IQCHGSFATASCLICKYKVDCEAVRGDIFNQVV
 PRCPRCPADEPLAIMKPEIVFFGENLPEQFHRAMKYDKDEVDLLIVIGSSLKVRPVAL
 IPSSIPHEVPQILINREPLPHLHFDVELLGD CDVIINELCHRLGGEYAKLCCNPFVKLS
 * EITEKPPRTQKELAYLSELPPTPLHVSEDS\$SPERTSPPDSSVIVTLLDQAAKSNDDL
 DVSESKGCM\$EKPQEVQTSRNVESIAEQMENPDLKNVGSSTGEKNERTSVAGTVRKCW
 * PNRVAKEQISRRLDGNQYLFLLPNRYIFHGAEVYSDEDDVLSSSSCGSNSDSGTCQS
 PSLEBPMEDESEIEEFYNGLEDEPDVPERAGGAGFGTGDGDDQEAINEAISVKQEV TDM
 NYPSNKS"
 misc feature 750..767
 /gene="SIRT1"
 /note="encodes putative nuclear localization signal"
 BASE COUNT 1225 a 732 c 925 g 1204 t
 ORIGIN
 1 gtcgagcgagg agcagaggag gcgaggaggagg agggccagag aggcagttgg aagatggcgg
 61 acgaggcgagg cctcgccctt cagcccgccg gctccccctc ggcggcgagg gccgacaggg
 121 agggcgcgctc gtcccccgcc ggggagccgc tccgcaagag gccgaggaga gatgggtccg
 181 gcctcgagcgc gagccccggc gagccccgtg gggcgccccc agagcgtgag gtgccggcgg
 241 cggccagggg ctgccccggg gcgccggcgg cggcgctgtg gcgggaggcg gaggcagagg
 301 cggcgggcggc agggcgggag caagaggccc aggcgactgc ggcggctggg gaaggagaca
 361 atggggccggg cctgcagggc ccatctcggg agccaccgct ggcgcacaac ttgtacgacg
 421 aagacgacga cgacgagggc gaggaggagg aagaggcggc ggcggcgcg attgggtacc
 481 gagataacct tctgttcggg gatgaaatta tcactaatgg ttttcattcc tgtgaaagtg
 541 atgaggagga tagagcctca catgcaagct ctagtgactg gactccaagg ccacggatag
 601 gtccatatac ttttgttcag caacatctta tgattggcac agatcctcga acaattctta
 661 aagattttatt gccggaaaca atacctccac ctgagttgga tgatatgaca ctgtggcaga
 721 ttgtttattaa tatcctttca gaaccaccaa aaaggaaaaa aagaaaagat attaatataa
 781 ttgaagatgc tgtgaaatta ctgcaagagt gcaaaaaaat tatagttcta actggagctg
 841 ggggtgtctgt ttcattgtga atacctgact tcagggtcaag ggatgggtatt tatgctcgcc
 901 ttgctgtaga cttcccagat cttccagatc ctcaagcgat gtttgatatt gaattttca
 961 gaaaagattcc aagaccattc ttcaagtttg caaaggaaat atatcctgga caattccagc
 1021 catctctctg tcacaaattc atagccttgt cagataagga aggaaaaacta cttcgcaact
 1081 ataccacgaa catagacacg ctggaccaca ttgcggsaat ccaaaaggata attcagtgct
 1141 atgggttcctt tgeaacagca tcttgccctga tttgtaaata caaagttgac tgtgaagctg
 1201 tacgaggaga tatttttaac caggtagttc ctgatgtcc taggtgcccga gctgatgaac
 1261 cgcttgctat catgaaacca gagattgtgt tttttgggtga aaattttacca gaacagtttc
 1321 atagagccat gaagtatgac aaagatgaag ttgacctcct cattgtttatt ggggtcttccc
 1381 tcaagtgtag accagtagca ctaattccaa gttccatacc ccatgaagtgt cctcagatat
 1441 taattaatag agaacccttg cctcatctgc attttgatgt agagcttctt ggagactgtg
 1501 atgtcataat taatgaattg tgtcatagggt taggtgggtga atatgccaaa ctttgctgta
 1561 accctgtaaa gctttcagaa attactgaaa aacctccacg aacacaaaaa gaattggctt
 1621 atttgtcaga gttgccaccc acaoctcttc atgtttcaga agactcaagt tcaccagaaa
 1681 gaacttcacc accagattct tcagtgattg tcacactttt agaccaagca gctaagagta
 1741 atgatgattt agatgtgtct gaatcaaaag gttgtatgga agaaaaacca caggaagtac
 1801 aaacttctag gaatgttgaa agtatgtctg aacagatgga aaatccggat ttgaagaatg
 1861 ttgggtctag tactggggag aaaaatgaaa gaacttcagt ggctggaaca gtgagaaaat
 1921 gctggccctaa tagagtggca agggagcaga ttagtagggc gcttgatggg aatcagtatc
 1981 tgtttttgcc accaaatcgt tacattttcc atggcgctga ggtatattca gactctgaag
 2041 atgacgtctt atcctctagt tcttgtggca gtaacagtga tagtgggaca tgccagagtc
 2101 caagttttaga agaaccatg gaggatgaaa gtgaaattga agaattctac aatggcttag
 2161 aagatgagcc tgaatgtcca gagcgagctg gagggagctg atttgggact gatggagatg
 2221 atcaagagggc aattaatgaa cctatatctg tgaaacagga agtaacagac atgaactatc
 2281 catcaaacaa atcatagtgt aataattgtg caggtacagg aattgttcca ccagcattag
 2341 gaacttttagc atgtcaaaat gaatgtttac ttgtgaactc gatagagcaa ggaaaccaga
 2401 aagggtgtaat atttataggt tggtaaaata gattgttttt catggataat ttttaacttc
 2461 attatttctg tacttgtaca aactcaacac taactttttt ttttttaaaa aaaaaaaggt
 2521 actaagtatc ttcaatcago tgttgggtca agactaactt tcttttaaa gttcatttgt

Figure 18 B

```

2581 atgataaatt catatgtgta tatataatTT tttttgtttt gtctagttag tttcaacatt
2641 ttttaaagttt tcaaaaagcc atcggaatgt taaattaatg taaagggaca gctaattctag
2701 accaaagaat ggtatTTTca cttttctttg taaattgaa tggtttgaag tactcaaaat
2761 ctgttacgct aaacttttga ttctttaaca caattatTTT taaacactgg cattttccaa
2821 aactgtggca gctaactTTT taaaatctca aatgacatgc agtgtgagta gaaggaagtc
2881 aacaatatgt ggggagagca ctcggttgtc tttactTTTa aaagtaatac ttggtgctaa
2941 gaatttcagg attattgtat ttacgttcaa atgaagatgg cttttgtact tcctgtggac
3001 atgtagtaat,gtctatattg gtcataaaaa ctaacctgaa aaacaaataa atgctttgga
3061 aatgtttcag ttgctttaga aacatttagtg cctgcctgga tccccctagt tttgaaatat
3121 ttgccattgt, tgttttaaata cctatcactg tggtagagct tgcattgac ttttccacaa
3181 gtattaaact, gccaaaatgt gaatatgcaa agcctttctg aatctataat aatgggtactt
3241 ctactgggga gagtgttaata ttttggactg ctgttttcca ttaatgagga gagcaacagg
3301 cccctgatta tacagttcca aagtaataag atgttaattg taattcagcc agaaagtaca
3361 tgtctcccat tgggaggatt tgggtgttaa taccaaactg ctagccctag tattatggag
3421 atgaacatga tgatgtaact tgtaatagca gaatagttaa tgaatgaaac tagttcttat
3481 aattttatctt tattttaaaag cttagcctgc cttaaaacta gagatcaact ttctcagctg
3541 caaaagcttc tagtctttca agaagtctcat actttatgaa attgcacagt aagcatttat
3601 ttttcagacc atttttgaac atcactccta aattaataaa gtattcctct gttgctttag
3661 tattttattac aataaaaagg gtttgaaata tagctgttct ttatgcataa aacaccagc
3721 taggaccatt actgccagag aaaaaaatcg tattgaatgg ccatttccct acttataaga
3781 tgtctcaatc tgaattttatt tggctacact aaagaatgca gtatatTTtag ttttccattt
3841 gcatgatgtt tgtgtgctat agatgatatt ttaaattgaa aagtttgttt taaattattt
3901 ttacagtga gactgttttc agctcttttt atattgtaca tagtctttta tgtaatttac
3961 tggcatatgt tttgtagact gtttaatgac tggatatctt ccttcaactt ttgaaataca
4021 aaaccagtgt tttttacttg tacactgttt taaagtctat taaaattgtc atttgacttt
4081 tttctg

```

//

Figure 18C

NCBI Sequence Viewer

```

/translation="MAFWGWRAAAALRLWGRVVERVEAGGGVGPFPQACGCRLLVLGGRD
DVSAGLRGSHGARGEPLDPA RPLQRP RPPEVPRAFRRQPRAAAP SFFSSIKGGRRSI
SFSVGASSVVGSGSSDKGLSLQDVAELIRARACQRVVVMVGAGISTPSGIPDFRSP
GSGLYSNLQQYDLPYPEAIFELPFFFFHNP KPFFTLAKELYPGNYKPNVTHYFLRL LHD
KGLLLRLYTQNI DGLERVSGIPASKLVEAHGTFASATCTVCQRPFPGEDIRADVMADR
VPRCPVCTGVVKPDIVFFGEPLPQRFL LHVDFPMADLLLILGTSLEVEPFASLTEAV
RSSVPRL LINRDLVGLAWHPRSRDVAQLGDVVHGVESLVELLGWTEEMRDLVQRETG
KLDGPKD"
BASE COUNT      361 a      507 c      593 g      408 t
ORIGIN
   1 ggcgcgcgggg gcgggggtgg gagggcggagg cggggccgggg gcgcgcgcggg cggggcgccg
  61 ggggcgcggggc gaggcccgag gactcctcgg actgcgcgga acatggcggt ctgggggttg
121 cgcgcgcgcgg cagccctccg gctgtggggc cgggtagttg aacgggtcga ggccggggga
181 ggcgtggggc cgtttcaggc ctgcggctgt cggctgggtg ttggcggcag ggacgatgtg
241 agtgcggggc tgagaggcag ccatggggcc cgcgggtgag ccttggaccc ggcgcgcccc
301 ttgcagaggc ctcccagacc cgaggtgccc agggcattcc ggaggcagcc gagggcagca
361 gctccaggtt tcttcttttc gagtattaaa ggtggaagaa ggtccatata ttttctgtg
421 ggtgcttcaa gtgtgttggt aagtggaggc agcagtgaca aggggaagct ttccctgcag
481 gatgtagctg agctgattcg ggccagagcc tgccagaggg tgggtggtcat ggtgggggcc
541 ggcattcagca caccagtggt cattccagac ttccagatcg cggggagtggt cctgtacagc
601 aacctccagc agtacgatct cccgtacccc gaggccattt ttgaactccc attcttcttt
661 cacaacccca agcccttttt cactttggcc aaggagctgt accctggaaa ctacaagccc
721 aacgtcactc actactttct ccggctgctt catgacaagg ggctgcttct gcggctctac
781 acgcagaaca tcgatgggct tgagagagtg tcgggcatcc ctgctcaca gctggttgaa
841 gctcatggaa cctttgcctc tgccacctgc acagtctgcc aaagaccctt cccaggggag
901 gacattcggg ctgacgtgat ggcagacagg gttccccgct gcccggctcg caccggcggt
961 gtgaagcccc acattgtgtt ctttggggag ccgctgcccc agaggttctt gctgcatgtg
1021 gttgatttcc ccatggcaga tctgtctctc atccttggga cctccctgga ggtggagcct
1081 tttgccagct tgaccgaggc cgtgcgagac tcagttcccc gactgctcat caaccgggag
1141 ttggtggggc ccttggcttg gcctcctcgc agcagggagc tggcccagct gggggacgtg
1201 gttcacggcg tggaaagcct agtggagctt ctgggctgga cagaagagat gcgggacott
1261 gtgcagcggg aaactgggaa gcttgatgga ccagacaaat aggatgatgg ctgccccac
1321 acaataaatg gtaacatagg agacatccac atcccaattc tgacaagacc tcatgcctga
1381 agacagcttg ggcaggtgaa accagaatat gtgaactgag tggacacccg aggctgccac
1441 tggaatgtct tctcaggcca tgagctgcag tgactggtag ggctgtgttt acagtcaggg
1501 ccaccccgct acatatacaa aggagctgcc tgcctgtttg ctgtgttgaa ctcttcactc
1561 tgctgaagct cctaattgga aaagctttct tctgactgtg accctcttga actgaatcag
1621 accaactgga atcccagacc gagtctgctt tctgtgcta gttgaacggc aagctcggca
1681 tctgttggtt acaagatcca gacttgggac gagcgggtccc cagccctctt catgttccga
1741 agtgtagtct tgaggccctg gtgcgcact tctagcatgt tggctcctt tagtggggct
1801 atttttaatg agagaaaatc tgttctttcc agcatgaaat acatttagtc tcctcaaaaa
1861 aaaaaaaca

```

//

Figure 19

NCBI Sequence Viewer

ORIGIN



```

1  gtccgtagag ctgtgagaga atgaagatga gctttgcggt gactttcagg tcagcaaaaag
61  gccgttggat cgcaaaacccc agccagccgt gctcgaaagc ctccattggg ttatttgtgc
121 cagcaagtcc tcctctggac cctgagaagg tcaaagagtt acagcgcttc atcacccttt
181 ccaagagact ccttgtgatg actggggcag gaatctccac cgaatcgggg ataccagact
241 acaggtcaga aaaagtgggg ctttatgccc gcactgaccg caggcccatc cagcatggtg
301 attttgtccg gagtgcacca atccgccagc ggtactgggc gagaaacttc gtaggctggc
361 ctcaattctc ctcccaccag cctaaccctg cactctgggc tttgagcacc tgggagaaac
421 tcggaaagct gtactggttg gtgacccaaa atgtggatgc tttgcacacc aaggcgggga
481 gtcggcgccg gacagagctc cacggatgca tggacagggt cctgtgcttg gattgtgggg
541 aacagactcc ccgggggggt ctgcaagagc gtttccaagt cctgaacccc acctggagtg
601 ctgaggccca tggcctggct cctgatggtg acgtctttct ctcagaggag caagtccgga
661 gctttcaggt cccaacctgc gttcaatgtg gaggccatct gaaaccagat gtcgttttct
721 tcggggacac agtgaacctc gacaagggtg attttgtgca caagcgtgta aaagaagccg
781 actccctctt ggtggtggga tcatccttgc aggtatactc tggttacagg tttatcctca
841 ctgcctggga gaagaagctc ccgattgcaa tactgaacat tgggccaca cggtcggatg
901 acttggcggt tctgaaactg aattctcgtt gtggagagtt gctgcctttg atagacccat
961 gctgaccaca gcctgatatt ccagaacctg gaacagggac tttcacttga atcttgctgc
1021 taaatgtaaa tgccttctca aatgacagat tccagttccc attcaacaga gtagggtgca
1081 ctgacaaagt atagaagggt ctaggtatct taatgtgtgg atattcttaa ttaaaaactca
1141 ttttttttaa ataaaaaatt gttcagcttt aaaa

```

//

Figure 20A

PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM

Search for






as

☐ 1: [AF083109](#) **Homo sapiens sirtuin type 4 (SIRT4) mRNA, complete cds** PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

LOCUS AF083109 1174 bp mRNA PRI 26-JUN-1999
 DEFINITION Homo sapiens sirtuin type 4 (SIRT4) mRNA, complete cds.
 ACCESSION AF083109
 VERSION AF083109.1 GI:5225323
 KEYWORDS
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 1174)
 AUTHORS Frye, R.A.
 TITLE Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity
 JOURNAL Biochem. Biophys. Res. Commun. 260 (1), 273-279 (1999)
 MEDLINE [99310604](#)
 REFERENCE 2 (bases 1 to 1174)
 AUTHORS Frye, R.A.
 TITLE Direct Submission
 JOURNAL Submitted (10-AUG-1998) Pathology, VA Med. Cent. (132L), Univ. of Pittsburgh, University Drive C, Pittsburgh, PA 15240, USA
 FEATURES
 source Location/Qualifiers
 1..1174
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="12"
 /map="12q"
 /tissue_type="testis"
 gene 1..1174
 /gene="SIRT4"
 CDS 21..965
 /gene="SIRT4"
 /note="yeast silent information regulator 2 (Sir2p) homolog; similar to prokaryotic sirtuins"
 /codon_start=1
 /product="sirtuin type 4"
 /protein_id="[AAD40852.1](#)"
 /db_xref="GI:5225324"
 /translation="MKMSFALTFRSAKGRWIANPSQPCSKASIGLFVPASPPLDPEKV
 KELQRFITLSKRLVMTGAGISTESGIPDYRSEKVGLYARTDRRPIQHGDVFVRSAPIR
 QRYWARNFVGWPQFSSHPNPAHWALSTWEKLGKLYWLVTQNVDAHTKAGSRRITEL
 HGCMDRVLCLDCGEQTPRGVLQERFQVLNPTWSAEAHGLAPDGDVFLSEEQVRSFQVP
 TCVQCGGHLKPDVVFVFGDTVNPDKVDFVHKRVKEADSLVVGSSSLQVSGYRFILTAW
 EKKLPAILNIGPTRSDDDLACLKLNRCGELLPLIDPC"

BASE COUNT 283 a 292 c 310 g 289 t

Figure 20B

PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM

Search for

as

☐ 1: [AF083110](#) **Homo sapiens sirtuin type 5 (SIRT5) mRNA, complete cds** PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

LOCUS AF083110 1633 bp mRNA PRI 26-JUN-1999
 DEFINITION Homo sapiens sirtuin type 5 (SIRT5) mRNA, complete cds.
 ACCESSION AF083110
 VERSION AF083110.1 GI:5225325
 KEYWORDS
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 1633)
 AUTHORS Frye, R.A.
 TITLE Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity
 JOURNAL Biochem. Biophys. Res. Commun. 260 (1), 273-279 (1999)
 MEDLINE 99310604

REFERENCE 2 (bases 1 to 1633)
 AUTHORS Frye, R.A.
 TITLE Direct Submission
 JOURNAL Submitted (10-AUG-1998) Pathology, VA Med. Cent. (132L), Univ. of Pittsburgh, University Drive C, Pittsburgh, PA 15240, USA

FEATURES
 source 1..1633
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /tissue_type="testis"
 gene 1..1633
 /gene="SIRT5"
 CDS 274..1206
 /gene="SIRT5"
 /note="yeast silent information regulator 2 (Sir2p) homolog; similar to prokaryotic sirtuins"
 /codon_start=1
 /product="sirtuin type 5"
 /protein_id="A40853.1"
 /db_xref="GI:5225325"
 /translation="MRPLQIVPSRLISQLYCGLKPPASTRNQICLKMARPSSSMADFR
 KPPAKAKHIVIISGAGVSAESGVPTFRGAGGYWRKWQAQDLATPLAFHNPSRVWEFY
 HYRREVMGSKEPNAGHRAIAECETRLGKQGRVVVITQNIDELHRKAGTKNLLIHS
 LFKTRCTSCGVVAENYKSPICPALSGKGAPEPGTQDASIPVEKLPREEAGCGGLLRP
 HVVWFGENLDPAILEEVDRELAHCDLCLVVGTSVVVPAAMFAPQVAARGVPVAEFNT
 ETTPATNRFRFHFQGPCGTTLPALACHENETVS"

BASE COUNT 455 a 357 c 422 g 399 t
 ORIGIN
 1. cgccctctagg agaaagcctg gaacgcgtac cggagggtac cagagctctt agcggggccgg

Figure 21A

NCBI Sequence Viewer

```

61 cagcatgtgc ggggcccaagt aaatggaaat gttttctaac atataaaaac ctacagaaga
121 agaaaataat tttctggatc aaattagaag tctgtattat attgatgtct ccagattcaa
181 atatattaga aagcagccgt ggagacaacc atcttcattt tgggagaaat aactaaagcc
241 cgctcaagc attagaacta cagacaaacc ctgatgcgac ctctccagat tgtcccaagt
301 cgattgattt cccagctata ttgtggcctg aagcctccag cgtccacacg aaaccagatt
361 tgctgaaaa tggctcggcc aagttcaagt atggcagatt ttcgaaagtt ttttgcaaaa
421 gcaaagcaca tagtcatcat ctcaggagct ggtgttagtg cagaaagtggt tgttccgacc
481 ttcagaggag ctggagggtta ttggagaaaa tggcaagccc aggacctggc gactcccctg
541 gcctttgccc acaaccgcgc cggggtgtgg gatttctacc actaccggcg ggagggtcatg
601 gggagcaagg dgcccaacgc cgggcaccgc gccatagccg agtgtgagac ccggctgggc
661 aagcagggcc ggcgagtcgt ggtcatcacc cagaacatcg atgagctgca ccgcaaggct
721 ggcaccaaga accttctgga gatccatggt agcttattta aaactcgatg tacctcttgt
781 ggagttgtgg ctgagaatta caagagtcca atttgtccag ctttatcagg aaaagggtgt
841 ccagaacctg gaactcaaga tgccagcatc ccagttgaga aacttceccg gtgtgaagag
901 gcaggctgcg ggggcttgct gcgacctcac gtcgtgtggt ttggagaaaa cctggatcct
961 gccattcttg aggaggttga cagagagctc gccactgtg atttatgtct agtgggtggc
1021 acttctctg tgggtgtacc agcagccatg tttgcccccc aggtggctgc caggggcgtg
1081 ccagtggctg aatttaacac ggagaccacc ccagctacga acagattcag gtttcatttc
1141 cagggaacct gtggaacgac tcttctgaa gcccttgccg gtcattgaaa tgaaactggt
1201 tcttaagtgt cctggggaag aaagaaatta cagtatatct aagaactagg ccacacgcag
1261 aggagaaatg gtcttatggg tgggtgagctg agtactgaac aatctaaaaa tagcctctga
1321 ttccctcgct ggaatocaac ctgttgataa gtgatggggg tttagaagta gcaaagagca
1381 ccacattca aaagtccacg aactggaag ttaattcata ttatttggtt tgaactgaaa
1441 cgtgaggtat ctttgatgtg tatggttggt tattgggagg gaaaaatttt gtaaattaga
1501 ttgtctaaaa aaaatagtta ttctgattat atttttgtta tctgggcaaa gtagaagtca
1561 aggggtaaaa accctactat tctgattttt gcacaagttt tagtggaaaa taaaatcaca
1621 ctctacagta ggt

```

//

Figure 21B

5

SEQUENCE LISTING

<110> Guarente, Leonard
Vaziri, Homayoun
Imai, Shin-Ichiro
Gu, Wei

10

<120> Sir2 and p53 Deacetylation

<130> 13407-015001

15

<150> US 60/303,456

<151> 2001-07-06

<150> US60/303,370

20

<151> 2001-07-06

<160> 18

<170> PatentIn version 3.1

25

<210> 1

<211> 18

<212> PRT

<213> homo sapiens

30

<400> 1

His Leu Lys Ser Lys Gly Gln Ser Thr Ser Arg His Lys Leu Met Phe

1

5

10

15

35

Lys Cys

40

5 <210> 2
 <211> 1546
 <212> DNA
 <213> homo sapiens

10 <400> 2
 atggaggagc cgcagtcaga tcctagcgtc gagccccctc tgagtcagga aacattttca
60
 gacctatgga aactacttcc tgaaaacaac gttctgtccc ccttgccgtc ccaagcaatg
15 120
 gatgatttga tgctgtcccc ggacgatatt gaacaatggt tcactgaaga ccaggtcca
180
 gatgaagctc ccagaatgcc agaggctgct cccccgtgg cccctgcacc agcagctcct
20 240
 acaccggcgg cccctgcacc agccccctcc tggcccctgt catcttctgt cccttcccag
300
25 360
 aaaacctacc agggcagcta cggtttccgt ctgggcttct tgcattctgg gacagccaag
 tctgtgactt gcacgtactc ccctgccctc aacaagatgt ttgccaact ggccaagacc
30 420
 tgccctgtgc agctgtgggt tgattccaca cccccgccg gcaccgcgt ccgcgccatg
480
 gccatctaca agcagtcaca gcacatgacg gaggttgtga ggcgtgccc ccacatgag
35 540
 cgctgctcag atagcgatgg tctggcccct cctcagcatc ttatccgagt ggaaggaaat
600
40

5 ttgogtgtgg agtattttgga tgacagaaac actttttcgac atagtgtggt ggtgccctat
660

 gagccgcctg aggttggctc tgactgtacc accatccact acaactacat gtgtaacagt
720
10 tcctgcatgg gcggcatgaa ccggaggccc atcctcacca tcatcacact ggaagactcc
780

 agtggtaatc tactgggacg gaacagcttt gaggtgcatg tttgtgcctg tcctgggaga
15 840

 gaccggcgca cagaggaaga gaatctccgc aagaaagggg agcctcacca cgagctgccc
900

20 ccaggggagca ctaagcgagc actgccaac aacaccagct cctctcccca gccaaagaag
960

 aaaccactgg atggagaata tttcacccct cagatccgtg ggcgtgagcg cttcgagatg
1020
25 ttccgagagc tgaatgaggc cttggaactc aaggatgccc aggctgggaa ggagccaggg
1080

 gggagcaggg ctcactccag ccacctgaag tccaaaaagg gtcagtctac ctcccgccat
30 1140

 aaaaaactca tgttcaagac agaagggcct gactcagact gacattctcc acttcttggt
1200

35 cccactgac agcctccac cccatctct cctcccctg ccattttggg ttttgggtct
1260

 ttgaaccctt gcttgcaata ggtgtgcgtc agaagcacc aggacttcca tttgctttgt
1320
40

5 cccggggctc cactgaacaa gttggcctgc actggtgttt tgttgtgggg aggaggatgg
 1380

 ggagtaggac ataccagctt agattttaag gtttttactg tgagggatgt ttgggagatg
 1440
 10 taagaaatgt tcttgcagtt aagggttagt ttacaatcag ccacattcta ggtagggacc
 1500

 cacttcaccg tactaaccag ggaagctgtc cctcactgtt gaattc
 15 1546

 <210> 3
 <211> 393
 20 <212> PRT
 <213> homo sapiens

 <400> 3

 25 Met Glu Glu Pro Gln Ser Asp Pro Ser Val Glu Pro Pro Leu Ser Gln
 1 5 10 15

 Glu Thr Phe Ser Asp Leu Trp Lys Leu Leu Pro Glu Asn Asn Val Leu
 30 20 25 30

 Ser Pro Leu Pro Ser Gln Ala Met Asp Asp Leu Met Leu Ser Pro Asp
 35 40 45
 35

 Asp Ile Glu Gln Trp Phe Thr Glu Asp Pro Gly Pro Asp Glu Ala Pro
 50 55 60

 40

5 Arg Met Pro Glu Ala Ala Pro Pro Val Ala Pro Ala Pro Ala Ala Pro
 65 70 75 80

10 Thr Pro Ala Ala Pro Ala Pro Ala Pro Ser Trp Pro Leu Ser Ser Ser
 85 90 95

15 Val Pro Ser Gln Lys Thr Tyr Gln Gly Ser Tyr Gly Phe Arg Leu Gly
 100 105 110

20 Phe Leu His Ser Gly Thr Ala Lys Ser Val Thr Cys Thr Tyr Ser Pro
 115 120 125

25 Ala Leu Asn Lys Met Phe Cys Gln Leu Ala Lys Thr Cys Pro Val Gln
 130 135 140

30 Leu Trp Val Asp Ser Thr Pro Pro Pro Gly Thr Arg Val Arg Ala Met
 145 150 155 160

35 Ala Ile Tyr Lys Gln Ser Gln His Met Thr Glu Val Val Arg Arg Cys
 165 170 175

40 Pro His His Glu Arg Cys Ser Asp Ser Asp Gly Leu Ala Pro Pro Gln
 180 185 190

45 His Leu Ile Arg Val Glu Gly Asn Leu Arg Val Glu Tyr Leu Asp Asp
 195 200 205

5 Arg Asn Thr Phe Arg His Ser Val Val Val Pro Tyr Glu Pro Pro Glu
 210 215 220

10 Val Gly Ser Asp Cys Thr Thr Ile His Tyr Asn Tyr Met Cys Asn Ser
 225 230 235 240

15 Ser Cys Met Gly Gly Met Asn Arg Arg Pro Ile Leu Thr Ile Ile Thr
 245 250 255

20 Leu Glu Asp Ser Ser Gly Asn Leu Leu Gly Arg Asn Ser Phe Glu Val
 260 265 270

25 His Val Cys Ala Cys Pro Gly Arg Asp Arg Arg Thr Glu Glu Glu Asn
 275 280 285

30 Lys Arg Ala Leu Pro Asn Asn Thr Ser Ser Ser Pro Gln Pro Lys Lys
 305 310 315 320

35 Lys Pro Leu Asp Gly Glu Tyr Phe Thr Leu Gln Ile Arg Gly Arg Glu
 325 330 335

40 Arg Phe Glu Met Phe Arg Glu Leu Asn Glu Ala Leu Glu Leu Lys Asp
 340 345 350

5 Ala Gln Ala Gly Lys Glu Pro Gly Gly Ser Arg Ala His Ser Ser His
 355 360 365

10 Leu Lys Ser Lys Lys Gly Gln Ser Thr Ser Arg His Lys Lys Leu Met
 370 375 380

15 Phe Lys Thr Glu Gly Pro Asp Ser Asp
 385 390

20 <210> 4
 <211> 1760
 <212> DNA
 <213> Homo sapiens

60 <400> 4
 gtcgaccctt tccaccctg gaagatggaa ataaacctgc gtgtgggtgg agtgtttagga

25 caaaaaaaaa aaaaaaaaaag tctagagcca ccgtccaggg agcaggtage tgctgggctc

120 cggggacact ttgcgttcgg gctgggagcg tgctttccac gacggtgaca cgcttccctg

30 180 gattggcagc cagactgcct tcogggtcac tgccatggag gagccgcagt cagatcctag

240 cgtcgagccc cctctgagtc aggaaacatt ttcagaccta tggaaactac ttcttgaaaa

35 300 caacgttctg tcccccttgc cgtcccaagc aatggatgat ttgatgctgt ccccggaoga

360

40

5 tattgaacaa tggttcactg aagaccacagg tccagatgaa gctcccagaa tgccagaggc
420

 tgctcccccc gtggcccctg caccagcagc tcctacaccg gcggcccctg caccagcccc
480

10 ctcctggccc ctgtcatctt ctgtcccttc ccagaaaacc taccagggca gctacggttt
540

 ccgtctgggc ttcttgcatt ctgggacagc caagtctgtg acttgcacgt actcccctgc
15 600

 cctcaacaag atgttttgcc aactggccaa gacctgccct gtgcagctgt gggttgattc
660

20 cacacccccg cccggcaccc gcgtccgcgc catggccatc tacaagcagt cacagcacat
720

 gacggagggt gtgaggeget gccccacca tgagcgtgc tcagatagcg atggtctggc
780

25 ccctcctcag catcttatcc gagtgaagg aaatttgctg gtggagtatt tggatgacag
840

 aaacactttt cgacatagtg tggtggtgcc ctatgagccg cctgaggttg gctctgactg
30 900

 taccaccatc cactacaact acatgtgtaa cagttcctgc atgggcggca tgaaccggag
960

35 gcccatcctc accatcatca cactggaaga ctccagtggg aatctactgg gacggaacag
1020

 ctttgaggtg catgtttgtg cctgtcctgg gagagaccgg cgcacagagg aagagaatct
1080

40

5 ccgcaagaaa ggggagcctc accacgagct gccccaggg agcactaagc gagcactgcc
1140

 caacaacacc agctcctctc cccagccaaa gaagaaacca ctggatggag aatatttcac
1200

10 ccttcagatc cgtgggcgtg agcgcttcga gatgttcga gagctgaatg aggccttgga
1260

 actcaaggat gcccaggctg ggaaggagcc aggggggagc agggctcact ccagccacct
15 1320

 gaagtccaaa aagggtcagt ctacctcccg ccataaaaaa ctcatgttca agacagaagg
1380

20 gcctgactca gactgacatt ctccacttct tgttccccac tgacagcctc ccacccccat
1440

 ctctccctcc cctgccatth tgggttttgg gtctttgaac ccttgcttgc aatagggtgtg
1500

25 cgtcagaagc acccaggact tccatttgct ttgtcccggg gctccactga acaagttggc
1560

 ctgcactggg gttttgttgt ggggaggagg atggggagta ggacatacca gcttagatth
30 1620

 taaggthttt actgtgaggg atgtttggga gatgtaagaa atgttcttgc agttaagggt
1680

35 tagthttaaa tcagccacat tctaggtagg gaccacttc accgtactaa ccagggaagc
1740

 tgtccctcac tgttgaattc
1760

40

5

<210> 5
<211> 3869
<212> DNA
<213> Mus musculus

10

<400> 5
g c g g a g c a g a g g a g g g c g g a g g g c c a g a g a g g c a g t t g g a a g a t g g c g g a c g a g g

60

15

t g g c g c t c g c c e t t c a g g c c g c c g g c t c c c c t t c c g c g g c g g c c g c c a t g g a g g c c g c g t

120

c g c a g c c g g c g g a c g a g c c g c t c c g c a a g a g g c c c c g c c g a g a c g g g c c t g g c c t c g g g c

180

20

g c a g c c c g g g c g a g c c g a g c g c a g c a g t g g c g c c g g c g g c c g c g g g g t g t g a g g c g g c g a

240

g c g c c g c g g c c c c g g c g g c g c t g t g g c g g g a g g c g g c a g g g g c g g c g g a g c g c g g a g c

25

300

g g g a g g c c c c g g c a c g g c g t g g c c g g g a c g g a g a c a a t g g g t c c g g c c t g c g g c g g g

360

30

a g c c g a g g g c g g c t g a c g a c t t c g a c g a c g a c g a g g g g c g a g g a g g a c g a g g c g g c g g

420

c g g c a g c g g c g g c g g c a g c g a t c g g c t a c c g a g a c a a c c t c c t g t t g a c c g a t g g a c t c c

480

35

t c a c t a a t g g c t t t c a t t c c t g t g a a a g t g a t g a c g a t g a c a g a a c g t c a c a c c a g c t

540

c t a g t g a c t g g a c t c c g c g g c c g c g g a t a g g t c c a t a t a c t t t t g t t c a g c a a c a t c t c a

40

600

5
tgattggcac cgatcctcga acaattctta aagatttatt accagaaaca attcctccac
660
ctgagctgga tgatatgacg ctgtggcaga ttgttattaa tctcctttca gaaccaccaa
10 720
agcggaaaaa aagaaaagat atcaatacaa ttgaagatgc tgtgaagtta ctgcaggagt
780
15 gtaaaaagat aatagtcttg actggagctg gggtttctgt ctctgtggg attcctgact
840
tcagatcaag agacggtatc tatgctcgcc ttgcggtgga ctcccagac ctcccagacc
900
20 ctcaagccat gtttgatatt gagtatttta gaaaagacc aagaccattc ttcaagtttg
960
caaaggaaat atatcccgga cagttccagc cgtctctgtg tcacaaattc atagctttgt
25 1020
cagataagga aggaaaacta cttcgaaatt atactcaaaa tatagatacc ttggagcagg
1080
30 ttgcaggaat ccaaaggatc cttcagtgtc atggttcctt tgcaacagca tcttgctga
1140
tttgtaaata caaagttgat tgtgaagctg ttctgtggaga catttttaac caggtagttc
1200
35 ctcggtgccc taggtgcccc gctgatgagc cacttgccat catgaagcca gagattgtct
1260
tctttggtga aaacttacca gaacagtttc atagagccat gaagtatgac aaagatgaag
40 1320

5
1380 ttgacctcct cattgttatt ggatcttctc tgaaagtgag accagtagca ctaattccaa
10 1440 gttctataacc ccatgaagtg cctcaaatat taataaatag ggaacctttg cctcatctac
1500 attttgatgt agagctcctt ggagactgcg atgttataat taatgagttg tgtcataggc
15 1560 taggtggtga atatgccaaa ctttgttgta accctgtaaa gctttcagaa attactgaaa
1620 aacctccacg cccacaaaag gaattgggtc atttatcaga gttgccacca acacctcttc
20 1680 atatttcgga agactcaagt tcacctgaaa gaactgtacc acaagactct tctgtgattg
1740 ctacacttgt agaccaagca acaaacaaca atgttaatga tttagaagta tctgaatcaa
25 1800 gttgtgtgga agaaaaacca caagaagtac agactagtag gaatgttgag aacattaatg
1860 tggaaaatcc agattttaag gctgttggtt ccagtactgc agacaaaaat gaaagaactt
1920 cagttgcaga aacagtgaga aaatgctggc ctaatagact tgcaaaggag cagattagta
30 1980 agcggcttga gggtaatcaa tacctgtttg taccaccaa tcgttacata ttccacggtg
2040 ctgaggtata ctcagactct gaagatgacg tcttgtcctc tagttcctgt ggcagtaaca
40

5
gtgacagtgg cacatgccag agtccaagtt tagaagaacc cttggaagat gaaagtgaaa
2100
ttgaagaatt ctacaatggc ttggaagatg atacggagag gcccgaatgt gctggaggat
10 2160
ctggatttgg agctgatgga ggggatcaag aggttggttaa tgaagctata gctacaagac
2220
aggaattgac agatgtaaac tatccatcag acaaatcata acactattga agctgtccgg
15 2280
attcaggaat tgctccacca gcattgggaa ctttagcatg tcaaaaaaat gaatgtttac
2340
ttgtgaactt gaacaaggaa atctgaaaga tgtattattt atagactgga aaatagattg
20 2400
tcttcttggg taattttctaa agttccatca tttctgtttg tacttgtaca ttcaacactg
25 2460
ttggttgact tcattcttct ttcaagggtc atttgtatga tacattcgta tgtatgtata
2520
at ttgtgtttt ttgcctaattg agtttcaacc ttttaaagtt ttcaaaagcc attggaatgt
30 2580
taatgtaaag ggaacagctt atctagacca aagaatggta tttcacactt tttgttttgt
2640
aacattgaat agttttaaac cctcaatttc tgttctgctg aacttttatt tttaggacag
35 2700
ttaacttttt aaacactggc attttccaaa acttgtggca gctaactttt taaaatcaca
40 2760

5
gatgacttgt aatgtgagga gtcagcaccg tgtctggagc actcaaaact tgggctcagt
2820
gtgtgaagcg tacttactgc atcgtttttg tacttgctgc agacgtggta atgtccaaac
10 2880
aggccccctga gactaatctg ataaatgatt tggaaatgtg tttcagttgt tctagaaaca
2940
atagtgcctg tctatatagg tccccttagt ttgaatattt gccattgttt aattaaatac
15 3000
ctatcactgt ggtagagcct gcatagatct tcaccacaaa tactgccaaag atgtgaatat
3060
20 gcaaagcctt tctgaatcta ataatggtac ttctactggg gagagtgtaa tattttggac
3120
tgctgttttt ccattaatga ggaaagcaat aggctctta attaaagtcc caaagtcata
25 3180
agataaattg tagctcaacc agaaagtaca ctgttgccctg ttgaggattt ggtgtaatgt
3240
atcccaaggt gttagccttg tattatggag atgaatacag atccaatagt caaatgaaac
30 3300
tagttcttag ttattttaaaa gcttagcttg ccttaaaaact agggatcaat tttctcaact
3360
35 gcagaaactt ttagcctttc aaacagttca cacctcagaa agtcagtatt tattttacag
3420
acttctttgg aacattgccc ccaaatttaa atattcatgt gggtttagta tttattacaa
40 3480

5

aaaaatgatt tgaaatatag ctgttcttta tgcataaaat acccagtttag gaccattact

3540

gccagaggag aaaagtatta agtagctcat ttccctacct aaaagataac tgaatttatt

10 3600

tggctacact aaagaatgca gtatatattag ttttccattt gcatgatgtg tttgtgctat

3660

15

agacaatatt ttaaattgaa aaatttgttt taaattattt ttacagtga gactgttttc

3720

agctcttttt atattgtaca tagactttta tgtaatctgg catatgtttt gtagaccgtt

3780

20

taatgactgg attatcttcc tccaactttt gaaatacaaa aacagtgttt tataactaaaa

3840

aaaaaaaaag tcgacgcggc cgcgaattc

25 3869

<210> 6

<211> 737

30

<212> PRT

<213> Mus musculus

<400> 6

35

Met Ala Asp Glu Val Ala Leu Ala Leu Gln Ala Ala Gly Ser Pro Ser

1

5

10

15

Ala Ala Ala Ala Met Glu Ala Ala Ser Gln Pro Ala Asp Glu Pro Leu

40

20

25

30

5

Arg Lys Arg Pro Arg Arg Asp Gly Pro Gly Leu Gly Arg Ser Pro Gly
 35 40 45

10

Glu Pro Ser Ala Ala Val Ala Pro Ala Ala Ala Gly Cys Glu Ala Ala
 50 55 60

15

Ser Ala Ala Ala Pro Ala Ala Leu Trp Arg Glu Ala Ala Gly Ala Ala
 65 70 75 80

20

Ala Ser Ala Glu Arg Glu Ala Pro Ala Thr Ala Val Ala Gly Asp Gly
 85 90 95

25

Asp Asn Gly Ser Gly Leu Arg Arg Glu Pro Arg Ala Ala Asp Asp Phe
 100 105 110

Asp Asp Asp Glu Gly Glu Glu Glu Asp Glu Ala Ala Ala Ala Ala Ala
 115 120 125

30

Ala Ala Ala Ile Gly Tyr Arg Asp Asn Leu Leu Leu Thr Asp Gly Leu
 130 135 140

35

Leu Thr Asn Gly Phe His Ser Cys Glu Ser Asp Asp Asp Asp Arg Thr
 145 150 155 160

40

Ser His Ala Ser Ser Ser Asp Trp Thr Pro Arg Pro Arg Ile Gly Pro
 165 170 175

5

Tyr Thr Phe Val Gln Gln His Leu Met Ile Gly Thr Asp Pro Arg Thr
 180 185 190

10

Ile Leu Lys Asp Leu Leu Pro Glu Thr Ile Pro Pro Pro Glu Leu Asp
 195 200 205

15

Asp Met Thr Leu Trp Gln Ile Val Ile Asn Ile Leu Ser Glu Pro Pro
 210 215 220

20

Lys Arg Lys Lys Arg Lys Asp Ile Asn Thr Ile Glu Asp Ala Val Lys
 225 230 235 240

25

Leu Leu Gln Glu Cys Lys Lys Ile Ile Val Leu Thr Gly Ala Gly Val
 245 250 255

30

Ser Val Ser Cys Gly Ile Pro Asp Phe Arg Ser Arg Asp Gly Ile Tyr
 260 265 270

Ala Arg Leu Ala Val Asp Phe Pro Asp Leu Pro Asp Pro Gln Ala Met
 275 280 285

35

Phe Asp Ile Glu Tyr Phe Arg Lys Asp Pro Arg Pro Phe Phe Lys Phe
 290 295 300

40

Ala Lys Glu Ile Tyr Pro Gly Gln Phe Gln Pro Ser Leu Cys His Lys
 305 310 315 320

5

Phe Ile Ala Leu Ser Asp Lys Glu Gly Lys Leu Leu Arg Asn Tyr Thr
325 330 335

10

Gln Asn Ile Asp Thr Leu Glu Gln Val Ala Gly Ile Gln Arg Ile Leu
340 345 350

15

Gln Cys His Gly Ser Phe Ala Thr Ala Ser Cys Leu Ile Cys Lys Tyr
355 360 365

20

Lys Val Asp Cys Glu Ala Val Arg Gly Asp Ile Phe Asn Gln Val Val
370 375 380

25

Pro Arg Cys Pro Arg Cys Pro Ala Asp Glu Pro Leu Ala Ile Met Lys
385 390 395 400

30

Pro Glu Ile Val Phe Phe Gly Glu Asn Leu Pro Glu Gln Phe His Arg
405 410 415

Ala Met Lys Tyr Asp Lys Asp Glu Val Asp Leu Leu Ile Val Ile Gly
420 425 430

35

Ser Ser Leu Lys Val Arg Pro Val Ala Leu Ile Pro Ser Ser Ile Pro
435 440 445

40

His Glu Val Pro Gln Ile Leu Ile Asn Arg Glu Pro Leu Pro His Leu
450 455 460

5

His Phe Asp Val Glu Leu Leu Gly Asp Cys Asp Val Ile Ile Asn Glu
 465 470 475 480

10

Leu Cys His Arg Leu Gly Gly Glu Tyr Ala Lys Leu Cys Cys Asn Pro
 485 490 495

15

Val Lys Leu Ser Glu Ile Thr Glu Lys Pro Pro Arg Pro Gln Lys Glu
 500 505 510

20

Leu Val His Leu Ser Glu Leu Pro Pro Thr Pro Leu His Ile Ser Glu
 515 520 525

25

Asp Ser Ser Ser Pro Glu Arg Thr Val Pro Gln Asp Ser Ser Val Ile
 530 535 540

30

Ala Thr Leu Val Asp Gln Ala Thr Asn Asn Asn Val Asn Asp Leu Glu
 545 550 555 560

35

Ser Arg Asn Val Glu Asn Ile Asn Val Glu Asn Pro Asp Phe Lys Ala
 580 585 590

40

Val Gly Ser Ser Thr Ala Asp Lys Asn Glu Arg Thr Ser Val Ala Glu
 595 600 605

5

Thr Val Arg Lys Cys Trp Pro Asn Arg Leu Ala Lys Glu Gln Ile Ser
610 615 620

10

Lys Arg Leu Glu Gly Asn Gln Tyr Leu Phe Val Pro Pro Asn Arg Tyr
625 630 635 640

15

Ile Phe His Gly Ala Glu Val Tyr Ser Asp Ser Glu Asp Asp Val Leu
645 650 655

20

Ser Ser Ser Ser Cys Gly Ser Asn Ser Asp Ser Gly Thr Cys Gln Ser
660 665 670

25

Pro Ser Leu Glu Glu Pro Leu Glu Asp Glu Ser Glu Ile Glu Glu Phe
675 680 685

30

Tyr Asn Gly Leu Glu Asp Asp Thr Glu Arg Pro Glu Cys Ala Gly Gly
690 695 700

Ser Gly Phe Gly Ala Asp Gly Gly Asp Gln Glu Val Val Asn Glu Ala
705 710 715 720

35

Ile Ala Thr Arg Gln Glu Leu Thr Asp Val Asn Tyr Pro Ser Asp Lys
725 730 735

Ser

40

5

<210> 7

<211> 3849

<212> DNA

10

<213> Mus musculus

<400> 7

gcggagcaga ggaggcgagg gcggagggcc agagaggcag ttggaagatg gcggacgagg

60

15

tggegetcgc ctttcaggcc gccggctccc ctccgcggc gcccgccatg gaggcgcgt

120

cgcagccggc ggacgagccg ctccgcaaga ggccccgcg agaaggccct ggcctcgggc

20

180

gcagcccggg cgagccgagc gcagcagtgg cgccggcggc cgcgggggtgt gaggcggcga

240

25

gcgcgcggc cccggcggcg ctgtggcggg aggcggcagg ggccggcggc agcgcgagc

300

gggaggcccc ggcgacggcc gtggccgggg acggagacaa tgggtccggc ctgcggcggg

360

30

agccgagggc ggctgacgac ttcgacgac acgagggcga ggaggaggac gaggcggcgg

420

cggcagcggc ggccggcagc atcggctacc gagacaacct cctgttgacc gatggactcc

35

480

tcactaatgg ctttcattcc tgtgaaagt atgacgatga cagaacgtca caccgcagct

540

5 ctagtgactg gactccgcgg ccgcggatag gtccatatac ttttgttcag caacatctca
600

 tgattggcac cgatcctcga acaattctta aagattttatt accagaaaca attcctccac
660
10 ctgagctgga tgatatgacg ctgtggcaga ttgtttattaa tatcctttca gaaccaccaa
720

 agcggaaaaa aagaaaagat atcaatacaa ttgaagatgc tgtgaagtta ctgcaggagt
15 780

 gtaaaaagat aatagttctg actggagctg gggtttctgt ctctgtggg attcctgact
840
20 tcagatcaag agacggtatc tatgctcgcc ttgcggtgga cttccagac ctccagacc
900

 ctcaagccat gtttgatatt gagtatttta gaaaagacc cagaccattc ttcaagtttg
960
25 caaaggaaat atatcccgga cagttccagc cgtctctgtg tcacaaattc atagctttgt
1020

 cagataagga aggaaaacta cttcgaaatt atactcaaaa tatagatacc ttggagcagg
30 1080

 ttgcaggaat ccaaaggatc cttcagtgtc atggttcctt tgcaacagca tcttgccatga
1140
35 tttgtaaata caaagttgat tgtgaagctg ttcgtggaga catttttaat caggtagttc
1200

 ctcgggtgcc taggtgccc gctgatgagc cacttgccat catgaagcca gagattgtct
1260
40

5 tcttttggtga aaacttacca gaacagtttc atagagccat gaagtatgac aaagatgaag
1320

 ttgaacctct cattgtttatt ggatctttctc tgaaagtgag accagtagca ctaattccaa
1380

10 gtttctatacc ccatgaagtg cctcaaatat taataaatag ggaacctttg cctcatctac
1440

 atthttgatgt agagctcctt ggagactgcg atgtttataat taatgagttg tgtcataggg
15 1500

 taggtggtga atatgccaaa ctttgttgta accctgtaaa gctttcagaa attactgaaa
1560

20 aacctccacg cccacaaaag gaattgggtc atttatcaga gttgccacca acacctcttc
1620

 atattttcgga agactcaagt tcacctgaaa gaactgtacc acaagactct tctgtgattg
1680

25 ctacacttgt agaccaagca acaaacaaca atgttaatga tttagaagta tctgaatcaa
1740

 gttgtgtgga agaaaaacca caagaagtac agactagtag gaatgttgag aacattaatg
30 1800

 tggaatatcc agatttttaag gctgttggtt ccagtactgc agacaaaaat gaaagaactt
1860

35 cagttgcaga aacagtgaga aaatgctggc ctaatagact tgcaaaggag cagattagta
1920

 agcggccttga gggtaatcaa tacctgtttg taccaccaa tcgttacata ttccacgggtg
1980

40

5 ctgaggtata ctcagactct gaagatgacg tcttgtcctc tagttcctgt ggcagtaaca
2040

 gtgacagtgg cacatgccag agtccaagtt tagaagaacc cttggaagat gaaagtgaag
2100

10 ttgaagaatt ctacaatggc ttggaagatg atacggagag gccogaatgt gctggaggat
2160

 ctggatttgg agctgatgga ggggatcaag aggttggtta tgaagctata gctacaagac
15 2220

 aggaattgac agatgtaaac tatccatcag acaaatacata acaactattga agctgtccgg
2280

20 attcaggaat tgctccacca gcattgggaa ctttagcatg tcaaaaaaat gaatgtttac
2340

 ttgtgaactt gaacaaggaa atctgaaaga tgtattatatt atagactgga aaatagattg
2400

25 tcttcttggg taattttctaa agttccatca tttctgtttg taattgtaca ttcaaacactg
2460

 ttggttgact tcatcttcct ttcaaggttc atttgtatga tacattcgta tgtatgtata
30 2520

 attttgtttt ttgcctaattg agtttcaacc ttttaaagtt ttcaaaagcc attggaatgt
2580

35 taatgtaaag ggaacagctt atctagacca aagaatggta tttcacactt ttttgtttgt
2640

 aacattgaat agtttaaagc cctcaatttc tgttctgctg aactttttatt tttaggacag
2700

40

5 ttaacttttt aaacactggc attttc meta acttggtggca gctaactttt taaaatcaca
2760

 gatgacttgt aatgtgagga gtcagcacgg tgtctggagc actcaaaact tgggctcagt
2820

10 gtgtgaagcg tacttactgc atcgtttttg tacttgctgc agacgtggta atgtccaaac
2880

 aggcccttga gactaatctg ataaatgatt tggaaatgtg tttcagttgt tctagaaaca
15 2940

 atagtgcctg tctatatagg tccccttagt ttgaatattt gccattgttt aattaaatac
3000

20 ctatcactgt ggtagagcct gcatagatct tcaccacaaa tactgccaag atgtgaatat
3060

 gcaaagcctt tctgaatcta ataatggtac ttctactggg gagagtgtaa tattttggac
3120

25 tgctgttttt ccattaatga ggaaagcaat aggctcttta attaaagtcc caaagtcata
3180

 agataaattg tagctcaacc agaaagtaca ctgttgcttg ttgaggattt ggtgtaatgt
30 3240

 atccaaggt gttagccttg tattatggag atgaatacag atccaatagt caaatgaaac
3300

35 tagttcttag ttattttaaaa gcttagcttg ccttaaaaact agggatcaat tttctcaact
3360

 gcagaaaactt ttagcctttc aaacagttca cacctcagaa agtcagtatt tattttacag
3420

40

5 acttcttttg aacattgcc ccaaatttaa atattcatgt gggtttagta tttattacaa
3480

 aaaaatgatt tgaaatatag ctgttcttta tgcataaaat acccagtttag gaccattact
3540
10 gccagaggag aaaagtatta agtagctcat ttccctacct aaaagataac tgaatttatt
3600

 tggctacact aaagaatgca gtatatattag ttttccattt gcatgatgtg tttgtgotat
15 3660

 agacaatatt ttaaattgaa aaatttggtt taaattattt ttacagtga gactgttttc
3720

20 agctcttttt atattgtaca tagactttta tgtaatctgg catatgtttt gtagacogtt
3780

 taatgactgg attatcttcc tocaaactttt gaaatacaaa aacagtgttt tataactaaaa
3840
25 aaaaaaaaa
3849

30 <210> 8
 <211> 737
 <212> PRT
 <213> Mus musculus

35 <400> 8

Met Ala Asp Glu Val Ala Leu Ala Leu Gln Ala Ala Gly Ser Pro Ser
1 5 10 15

40

5 Ala Ala Ala Ala Met Glu Ala Ala Ser Gln Pro Ala Asp Glu Pro Leu
 20 25 30

10 Arg Lys Arg Pro Arg Arg Asp Gly Pro Gly Leu Gly Arg Ser Pro Gly
 35 40 45

15 Glu Pro Ser Ala Ala Val Ala Pro Ala Ala Ala Gly Cys Glu Ala Ala
 50 55 60

20 Ser Ala Ala Ala Pro Ala Ala Leu Trp Arg Glu Ala Ala Gly Ala Ala
 65 70 75 80

25 Ala Ser Ala Glu Arg Glu Ala Pro Ala Thr Ala Val Ala Gly Asp Gly
 85 90 95

30 Asp Asn Gly Ser Gly Leu Arg Arg Glu Pro Arg Ala Ala Asp Asp Phe
 100 105 110

35 Asp Asp Asp Glu Gly Glu Glu Glu Asp Glu Ala Ala Ala Ala Ala Ala
 115 120 125

40 Ala Ala Ala Ile Gly Tyr Arg Asp Asn Leu Leu Leu Thr Asp Gly Leu
 130 135 140

45 Leu Thr Asn Gly Phe His Ser Cys Glu Ser Asp Asp Asp Asp Arg Thr
 145 150 155 160

5 Ser His Ala Ser Ser Ser Asp Trp Thr Pro Arg Pro Arg Ile Gly Pro
 165 170 175

10 Tyr Thr Phe Val Gln Gln His Leu Met Ile Gly Thr Asp Pro Arg Thr
 180 185 190

15 Ile Leu Lys Asp Leu Leu Pro Glu Thr Ile Pro Pro Pro Glu Leu Asp
 195 200 205

20 Asp Met Thr Leu Trp Gln Ile Val Ile Asn Ile Leu Ser Glu Pro Pro
 210 215 220

25 Lys Arg Lys Lys Arg Lys Asp Ile Asn Thr Ile Glu Asp Ala Val Lys
 225 230 235 240

30 Leu Leu Gln Glu Cys Lys Lys Ile Ile Val Leu Thr Gly Ala Gly Val
 245 250 255

35 Ser Val Ser Cys Gly Ile Pro Asp Phe Arg Ser Arg Asp Gly Ile Tyr
 260 265 270

40 Ala Arg Leu Ala Val Asp Phe Pro Asp Leu Pro Asp Pro Gln Ala Met
 275 280 285

45 Phe Asp Ile Glu Tyr Phe Arg Lys Asp Pro Arg Pro Phe Phe Lys Phe
 290 295 300

5 Ala Lys Glu Ile Tyr Pro Gly Gln Phe Gln Pro Ser Leu Cys His Lys
 305 310 315 320

10 Phe Ile Ala Leu Ser Asp Lys Glu Gly Lys Leu Leu Arg Asn Tyr Thr
 325 330 335

15 Gln Asn Ile Asp Thr Leu Glu Gln Val Ala Gly Ile Gln Arg Ile Leu
 340 345 350

20 Gln Cys His Gly Ser Phe Ala Thr Ala Ser Cys Leu Ile Cys Lys Tyr
 355 360 365

25 Lys Val Asp Cys Glu Ala Val Arg Gly Asp Ile Phe Asn Gln Val Val
 370 375 380

30 Pro Glu Ile Val Phe Phe Gly Glu Asn Leu Pro Glu Gln Phe His Arg
 405 410 415

35 Ala Met Lys Tyr Asp Lys Asp Glu Val Asp Leu Leu Ile Val Ile Gly
 420 425 430

40 Ser Ser Leu Lys Val Arg Pro Val Ala Leu Ile Pro Ser Ser Ile Pro
 435 440 445

5 His Glu Val Pro Gln Ile Leu Ile Asn Arg Glu Pro Leu Pro His Leu
 450 455 460

10 His Phe Asp Val Glu Leu Leu Gly Asp Cys Asp Val Ile Ile Asn Glu
 465 470 475 480

15 Leu Cys His Arg Leu Gly Gly Glu Tyr Ala Lys Leu Cys Cys Asn Pro
 485 490 495

20 Val Lys Leu Ser Glu Ile Thr Glu Lys Pro Pro Arg Pro Gln Lys Glu
 500 505 510

25 Leu Val His Leu Ser Glu Leu Pro Pro Thr Pro Leu His Ile Ser Glu
 515 520 525

30 Asp Ser Ser Ser Pro Glu Arg Thr Val Pro Gln Asp Ser Ser Val Ile
 530 535 540

35 Ala Thr Leu Val Asp Gln Ala Thr Asn Asn Asn Val Asn Asp Leu Glu
 545 550 555 560

40 Val Ser Glu Ser Ser Cys Val Glu Glu Lys Pro Gln Glu Val Gln Thr
 565 570 575

45 Ser Arg Asn Val Glu Asn Ile Asn Val Glu Asn Pro Asp Phe Lys Ala
 580 585 590

5 Val Gly Ser Ser Thr Ala Asp Lys Asn Glu Arg Thr Ser Val Ala Glu
 595 600 605

10 Thr Val Arg Lys Cys Trp Pro Asn Arg Leu Ala Lys Glu Gln Ile Ser
 610 615 620

15 Lys Arg Leu Glu Gly Asn Gln Tyr Leu Phe Val Pro Pro Asn Arg Tyr
 625 630 635 640

20 Ile Phe His Gly Ala Glu Val Tyr Ser Asp Ser Glu Asp Asp Val Leu
 645 650 655

25 Pro Ser Leu Glu Glu Pro Leu Glu Asp Glu Ser Glu Ile Glu Glu Phe
 675 680 685

30 Tyr Asn Gly Leu Glu Asp Asp Thr Glu Arg Pro Glu Cys Ala Gly Gly
 690 695 700

35 Ser Gly Phe Gly Ala Asp Gly Gly Asp Gln Glu Val Val Asn Glu Ala
 705 710 715 720

40 Ile Ala Thr Arg Gln Glu Leu Thr Asp Val Asn Tyr Pro Ser Asp Lys
 725 730 735

5 Ser

<210> 9

10 <211> 1963

<212> DNA

<213> Homo sapiens

<400> 9

15 gtgttgtagc aaagcgcgtc tgcggccgca atgtctgctg agagttgtag ttctgtgcc

60

tatcacggcc actccattt ctggtgccgt cacgggacag agcagtcggt gacaggacag

120

20

agcagtcggt gacgggacac agtggttggg gacgggacag agcggtcggt gacagcctca

180

agggcttcag caccgcgccc atggcagagc cagaccctc tcaccctctg gagaccag

25 240

cagggaaggt gcaggaggct caggactcag attcagactc tgaggaggga gccgctggtg

300

30 gagaagcaga catggacttc ctgcggaact tattctccca gacgtcagc ctgggcagcc

360

agaaggagcg tctgctggac gagctgacct tggaaggggt ggcccggtac atgcagagcg

420

35

aacgctgtcg cagagtcata tgtttggtgg gagctggaat ctccacatcc gcaggcatcc

480

ccgactttcg ctctccatcc aocggcctct atgacaacct agagaagtac catcttccct

40 540

5
accagagggc catctttgag atcagctatt tcaagaaaca tccggaaccc ttcttggccc
600
tcgccaagga actctatcct gggcagttca agccaaccat ctgtcactac ttcattggcc
10 660
tgctgaagga caaggggcta ctcttgcgct gctacacgca gaacatagat accctggagc
720
gaatagccgg gctggaacag gaggacttgg tggaggcgca cggcaccttc tacacatcac
15 780
actgcgtcag cgccagctgc cggcacgaat acccgctaag ctggatgaaa gagaagatct
840
tctctgaggt gacgoccaaag tgtgaagact gtcagagcct ggtgaagcct gatatcgtct
20 900
tttttgggtga gagcctccca ggcggtttct tctctgtat gcagtcagac ttcctgaagg
25 960
tggaacctct cctggtcatt ggtacctct tgcaggtgca gccctttgcc tccctcatca
1020
gcaaggcacc cctctccacc cctcgccctgc tcatcaacaa ggagaaagct ggccagtcgg
30 1080
accctttcct ggggatgatt atgggcctcg gaggaggcat ggactttgac tccaagaagg
1140
cctacaggga cgtggcctgg ctgggtgaat gcgaccaggg ctgcctggcc cttgctgagc
35 1200
tccttggatg gaagaaggag ctggaggacc ttgtccggag ggagcacgcc agcatagatg
40 1260

5
cccagtcggg ggcgggggtc cccaaccca gcacttcagc ttccccaag aagtcctcgc
1320
cacctgccaa ggacgaggcc aggacaacag agaggagaa accccagtga cagctgcac
10 1380
tcccaggcgg gatgccgagc tcctcaggga cagctgagcc ccaaccgggc ctggccccct
1440
cttaaccagc agttcttgtc tggggagctc agaacatccc ccaatctctt acagctccct
15 1500
ccccaaaact ggggtcccag caaccctggc cccaacccc agcaaattct taacacctcc
1560
20 tagaggccaa ggcttaaaca ggcattctta ccagccccac tgtctctaac cactcctggg
1620
ctaaggagta acctccctca tctctaactg cccccacggg gccagggcta cccagaaact
25 1680
tttaactctt ccaggacagg gagcttcggg ccccaactct gtctcctgcc cccggggggc
1740
30 tgtggctaag taaaccatac ctaacctacc ccagtgtggg tgtgggcctc tgaatataac
1800
ccacaccag cgtaggggga gtctgagccg ggagggtcc cgagtctctg ccttcagctc
1860
35 ccaaagtggg tgggtgggcc ccttcacgtg ggaccactt cccatgctgg atgggcagaa
1920
gacattgctt attggagaca aattaaaaac aaaaacaact aac
40 1963

5

<210> 10

<211> 389

<212> PRT

10

<213> Homo sapiens

<400> 10

15

Met Ala Glu Pro Asp Pro Ser His Pro Leu Glu Thr Gln Ala Gly Lys
 1 5 10 15

20

Val Gln Glu Ala Gln Asp Ser Asp Ser Asp Ser Glu Gly Gly Ala Ala
 20 25 30

25

Gly Gly Glu Ala Asp Met Asp Phe Leu Arg Asn Leu Phe Ser Gln Thr
 35 40 45

30

Leu Ser Leu Gly Ser Gln Lys Glu Arg Leu Leu Asp Glu Leu Thr Leu
 50 55 60

Glu Gly Val Ala Arg Tyr Met Gln Ser Glu Arg Cys Arg Arg Val Ile
 65 70 75 80

35

Cys Leu Val Gly Ala Gly Ile Ser Thr Ser Ala Gly Ile Pro Asp Phe
 85 90 95

40

Arg Ser Pro Ser Thr Gly Leu Tyr Asp Asn Leu Glu Lys Tyr His Leu
 100 105 110

5

Pro Tyr Pro Glu Ala Ile Phe Glu Ile Ser Tyr Phe Lys Lys His Pro
 115 120 125

10

Glu Pro Phe Phe Ala Leu Ala Lys Glu Leu Tyr Pro Gly Gln Phe Lys
 130 135 140

15

Pro Thr Ile Cys His Tyr Phe Met Arg Leu Leu Lys Asp Lys Gly Leu
 145 150 155 160

20

Leu Leu Arg Cys Tyr Thr Gln Asn Ile Asp Thr Leu Glu Arg Ile Ala
 165 170 175

Gly Leu Glu Gln Glu Asp Leu Val Glu Ala His Gly Thr Phe Tyr Thr
 180 185 190

25

Ser His Cys Val Ser Ala Ser Cys Arg His Glu Tyr Pro Leu Ser Trp
 195 200 205

30

Met Lys Glu Lys Ile Phe Ser Glu Val Thr Pro Lys Cys Glu Asp Cys
 210 215 220

35

Gln Ser Leu Val Lys Pro Asp Ile Val Phe Phe Gly Glu Ser Leu Pro
 225 230 235 240

Ala Arg Phe Phe Ser Cys Met Gln Ser Asp Phe Leu Lys Val Asp Leu
 245 250 255

40

5
 Leu Leu Val Met Gly Thr Ser Leu Gln Val Gln Pro Phe Ala Ser Leu
 260 265 270

10
 Ile Ser Lys Ala Pro Leu Ser Thr Pro Arg Leu Leu Ile Asn Lys Glu
 275 280 285

15
 Lys Ala Gly Gln Ser Asp Pro Phe Leu Gly Met Ile Met Gly Leu Gly
 290 295 300

20
 Gly Gly Met Asp Phe Asp Ser Lys Lys Ala Tyr Arg Asp Val Ala Trp
 305 310 315 320

25
 Leu Gly Glu Cys Asp Gln Gly Cys Leu Ala Leu Ala Glu Leu Leu Gly
 325 330 335

30
 Trp Lys Lys Glu Leu Glu Asp Leu Val Arg Arg Glu His Ala Ser Ile
 340 345 350

35
 Asp Ala Gln Ser Gly Ala Gly Val Pro Asn Pro Ser Thr Ser Ala Ser
 355 360 365

40
 Pro Lys Lys Ser Pro Pro Pro Ala Lys Asp Glu Ala Arg Thr Thr Glu
 370 375 380

Arg Glu Lys Pro Gln
 385

5

<210> 11
<211> 4086
<212> DNA
<213> Homo sapiens

10

<400> 11
gtcgcagcggg agcagaggag gcgagggagg agggccagag aggcagttgg aagatggcgg

60

15

acgaggcggc cctcgccctt cagcccggcg gctccccctc ggccggcggg gccgacaggg

120

aggccgcgtc gtcccccgcc ggggagccgc tccgcaagag gccgcggaga gatggtcccc

180

20

gcctcgagcg gagccccggc gagcccggcg gggcgggccc agagcgtgag gtgccggcgg

240

cggccagggg ctgcccgggt gcggcggcgg cggcgctgtg gcgggaggcg gaggcagagg

25

300

cggcggcggc aggcggggag caagaggccc aggcgactgc ggccgctggg gaaggagaca

360

30

atgggcccgg cctgcagggc ccatctcggg agccaccgct ggccgacaac ttgtacgacg

420

aagacgacga cgacgagggc gaggaggagg aagaggcggc ggccggcgcg attgggtacc

480

35

gagataacct tctgttcggt gatgaaatta tcactaatgg ttttcattcc tgtgaaagtg

540

atgaggagga tagagcctca catgcaagct ctagtgactg gactccaagg ccacggatag

40

600

5
gtccatatac ttttgttcag caacatctta tgattggcac agatcctcga acaattctta
660
aagatatttatt gccggaaaca atacctccac ctgagttgga tgatatgaca ctgtggcaga
10 720
ttgtttattaa tatcctttca gaaccaccaa aaaggaaaaa aagaaaagat attaatacaa
780
ttgaagatgc tgtgaaatta ctgcaagagt gcaaaaaaat tatagttcta actggagctg
840
gggtgtctgt ttcattgtgga atacctgact tcaggtcaag ggatgggtatt tatgctcgcc
900
20
ttgctgtaga cttcccagat cttccagatc ctcaagcgat gtttgatatt gaatatttca
960
gaaaagatcc aagaccattc ttcaagtttg caaaggaaat atatcctgga caattccagc
25 1020
catctctctg tcacaaattc atagccttgt cagataagga aggaaaacta cttcgcaact
1080
ataccagaa catagacacg ctggaacagg ttgcgggaat ccaaaggata attcagtgtc
30 1140
atggttcctt tgcaacagca tcttgacctga tttgtaaata caaagttgac tgtgaagctg
1200
35
tacgaggaga tattttttaat caggtagttc ctcgatgtcc taggtgccca gctgatgaac
1260
cgcttgctat catgaaacca gagattgtgt tttttggtga aaatttacca gaacagtttc
40 1320

5
atagagccat gaagtatgac aaagatgaag ttgacctcct cattgttatt gggctcttccc
1380

10 1440
tcaaagtaag accagtagca ctaattccaa gttccatacc ccatgaagtg cctcagatat

1500
taattaatag agaacctttg cctcatctgc attttgatgt agagcttctt ggagactgtg

15 1560
atgtcataat taatgaattg tgtcataggt taggtgggtga atatgccaaa ctttgctgta

1620
accctgtaaa gctttcagaa attactgaaa aacctccacg aacacaaaaa gaattggctt

20 1680
atttgtcaga gttgccaccc acacctcttc atgtttcaga agactcaagt tcaccagaaa

1740
gaacttcacc accagattct tcagtgattg tcacactttt agaccaagca gctaagagta

25 1800
atgatgattt agatgtgtct gaatcaaaag gttgtatgga agaaaaacca caggaagtac

30 1860
aaacttctag gaatgttgaa agtattgctg aacagatgga aaatccggat ttgaagaatg

1920
ttggttctag tactggggag aaaaatgaaa gaacttcagt ggctggaaca gtgagaaaat

35 1980
gctggcctaa tagagtggca aaggagcaga ttagtaggcg gcttgatggg aatcagtatc

2040
tgtttttgcc accaaatcgt tacattttcc atggcgctga ggtatattca gactctgaag

40

5
atgacgtctt atcctctagt tcttgtggca gtaacagtga tagtgggaca tgccagagtc
2100
caagtttaga agaaccatg gaggatgaaa gtgaaattga agaattctac aatggcttag
10 2160
aagatgagcc tgatgttcca gagagagctg gaggagctgg atttgggact gatggagatg
2220
atcaagaggc aattaatgaa gctatatctg tgaaacagga agtaacagac atgaactatc
15 2280
catcaaaca atcatagtgt aataattgtg caggtacagg aattgttcca ccagcattag
2340
gaactttagc atgtcaaaat gaatgtttac ttgtgaactc gatagagcaa ggaaaccaga
20 2400
aagggtgaat atttataggt tggtaaaata gattgttttt catggataat ttttaacttc
25 2460
attatttctg tacttgtaca aactcaacac taactttttt ttttttaaaa aaaaaaagg
2520
actaagtatc ttcaatcagc tgttgggtca agactaactt tcttttaaag gttcatttgt
30 2580
atgataaatt catatgtgta tatataattt tttttgtttt gtctagttag tttcaacatt
2640
tttaaagttt tcaaaaagcc atcggaatgt taaattaatg taaagggaca gctaacttag
35 2700
accaaagaat ggtattttca cttttctttg taacattgaa tggtttgaag tactcaaaat
40 2760

5
ctgttacgct aaacttttga ttctttaaca caattatitt taaacactgg catitttccaa
2820
aactgtggca gctaactttt taaaatctca aatgacatgc agtgtgagta gaaggaagtc
10 2880
aacaatatgt ggggagagca ctcggttgtc ttacttttta aaagtaatac ttggtgctaa
2940
gaatttcagg attattgtat ttacgttcaa atgaagatgg cttttgtact tcctgtggac
15 3000
atgtagtaat gtctatatattg gtcataaaaa ctaacctgaa aaacaaataa atgctttgga
3060
20 aatgtttcag ttgctttaga aacattagtg cctgcctgga tccccttagt tttgaaatat
3120
ttgccattgt tgtttaaata cctatcactg tggtagagct tgcattgac ttttccacaa
25 3180
gtattaaact gccaaaatgt gaatatgcaa agcctttctg aatctataat aatggtaactt
3240
30 ctactgggga gagtgttaata ttttggactg ctgttttcca ttaatgagga gagcaacagg
3300
cccctgatta tacagttcca aagtaataag atgttaattg taattcagcc agaaagtaca
3360
35 tgtctcccat tgggaggatt tgggtgttaa taccaaactg ctagccctag tattatggag
3420
atgaacatga tgatgtaact tgtaatagca gaatagttaa tgaatgaaac tagttcttat
40 3480

5
aatttatctt tattttaaag cttagcctgc cttaaacta gagatcaact ttctcagctg
3540
caaaagcttc tagtctttca agaagttcat actttatgaa attgcacagt aagcatttat
10 3600
ttttcagacc atttttgaac atcactccta aattaataaa gtattcctct gttgctttag
3660
tatttattac aataaaaagg gtttgaaata tagctgttct ttatgcataa aacaccagc
15 3720
taggaccatt actgccagag aaaaaaatcg tattgaatgg ccatttcctt acttataaga
3780
20
tgtctcaatc tgaatttatt tggctacact aaagaatgca gtatatttag ttttccattt
3840
gcatgatgtt tgtgtgctat agatgatatt ttaaattgaa aagtttgttt taaattattt
25 3900
ttacagtgaa gactgttttc agctcttttt atattgtaca tagtctttta tgtaatttac
3960
30
tggcatatgt tttgtagact gtttaatgac tggatatctt ctttcaactt ttgaaataca
4020
aaaccagtgt tttttacttg tacactgttt taaagtctat taaaattgtc atttgacttt
4080
35
tttctg
4086
40 <210> 12

5 <211> 747
 <212> PRT
 <213> Homo sapiens

 <400> 12
 10
 Met Ala Asp Glu Ala Ala Leu Ala Leu Gln Pro Gly Gly Ser Pro Ser
 1 5 10 15

 15 Ala Ala Gly Ala Asp Arg Glu Ala Ala Ser Ser Pro Ala Gly Glu Pro
 20 25 30

 Leu Arg Lys Arg Pro Arg Arg Asp Gly Pro Gly Leu Glu Arg Ser Pro
 20 35 40 45

 Gly Glu Pro Gly Gly Ala Ala Pro Glu Arg Glu Val Pro Ala Ala Ala
 50 55 60
 25

 Arg Gly Cys Pro Gly Ala Ala Ala Ala Ala Leu Trp Arg Glu Ala Glu
 65 70 75 80

 30 Ala Glu Ala Ala Ala Ala Gly Gly Glu Gln Glu Ala Gln Ala Thr Ala
 85 90 95

 35 Ala Ala Gly Glu Gly Asp Asn Gly Pro Gly Leu Gln Gly Pro Ser Arg
 100 105 110

 Glu Pro Pro Leu Ala Asp Asn Leu Tyr Asp Glu Asp Asp Asp Asp Glu
 40 115 120 125

5

Gly Glu Glu Glu Glu Glu Ala Ala Ala Ala Ala Ile Gly Tyr Arg Asp
130 135 140

10

Asn Leu Leu Phe Gly Asp Glu Ile Ile Thr Asn Gly Phe His Ser Cys
145 150 155 160

15

Glu Ser Asp Glu Glu Asp Arg Ala Ser His Ala Ser Ser Ser Asp Trp
165 170 175

20

Thr Pro Arg Pro Arg Ile Gly Pro Tyr Thr Phe Val Gln Gln His Leu
180 185 190

25

Met Ile Gly Thr Asp Pro Arg Thr Ile Leu Lys Asp Leu Leu Pro Glu
195 200 205

30

Thr Ile Pro Pro Pro Glu Leu Asp Asp Met Thr Leu Trp Gln Ile Val
210 215 220

Ile Asn Ile Leu Ser Glu Pro Pro Lys Arg Lys Lys Arg Lys Asp Ile
225 230 235 240

35

Asn Thr Ile Glu Asp Ala Val Lys Leu Leu Gln Glu Cys Lys Lys Ile
245 250 255

40

Ile Val Leu Thr Gly Ala Gly Val Ser Val Ser Cys Gly Ile Pro Asp
260 265 270

5

Phe Arg Ser Arg Asp Gly Ile Tyr Ala Arg Leu Ala Val Asp Phe Pro
 275 280 285

10

Asp Leu Pro Asp Pro Gln Ala Met Phe Asp Ile Glu Tyr Phe Arg Lys
 290 295 300

15

Asp Pro Arg Pro Phe Phe Lys Phe Ala Lys Glu Ile Tyr Pro Gly Gln
 305 310 315 320

20

Phe Gln Pro Ser Leu Cys His Lys Phe Ile Ala Leu Ser Asp Lys Glu
 325 330 335

25

Gly Lys Leu Leu Arg Asn Tyr Thr Gln Asn Ile Asp Thr Leu Glu Gln
 340 345 350

30

Val Ala Gly Ile Gln Arg Ile Ile Gln Cys His Gly Ser Phe Ala Thr
 355 360 365

Ala Ser Cys Leu Ile Cys Lys Tyr Lys Val Asp Cys Glu Ala Val Arg
 370 375 380

35

Gly Asp Ile Phe Asn Gln Val Val Pro Arg Cys Pro Arg Cys Pro Ala
 385 390 395 400

40

Asp Glu Pro Leu Ala Ile Met Lys Pro Glu Ile Val Phe Phe Gly Glu
 405 410 415

5

Asn Leu Pro Glu Gln Phe His Arg Ala Met Lys Tyr Asp Lys Asp Glu
 420 425 430

10

Val Asp Leu Leu Ile Val Ile Gly Ser Ser Leu Lys Val Arg Pro Val
 435 440 445

15

Ala Leu Ile Pro Ser Ser Ile Pro His Glu Val Pro Gln Ile Leu Ile
 450 455 460

20

Asn Arg Glu Pro Leu Pro His Leu His Phe Asp Val Glu Leu Leu Gly
 465 470 475 480

25

Asp Cys Asp Val Ile Ile Asn Glu Leu Cys His Arg Leu Gly Gly Glu
 485 490 495

30

Tyr Ala Lys Leu Cys Cys Asn Pro Val Lys Leu Ser Glu Ile Thr Glu
 500 505 510

Lys Pro Pro Arg Thr Gln Lys Glu Leu Ala Tyr Leu Ser Glu Leu Pro
 515 520 525

35

Pro Thr Pro Leu His Val Ser Glu Asp Ser Ser Ser Pro Glu Arg Thr
 530 535 540

40

Ser Pro Pro Asp Ser Ser Val Ile Val Thr Leu Leu Asp Gln Ala Ala
 545 550 555 560

5

Lys Ser Asn Asp Asp Leu Asp Val Ser Glu Ser Lys Gly Cys Met Glu
 565 570 575

10

Glu Lys Pro Gln Glu Val Gln Thr Ser Arg Asn Val Glu Ser Ile Ala
 580 585 590

15

Glu Gln Met Glu Asn Pro Asp Leu Lys Asn Val Gly Ser Ser Thr Gly
 595 600 605

20

Glu Lys Asn Glu Arg Thr Ser Val Ala Gly Thr Val Arg Lys Cys Trp
 610 615 620

25

Pro Asn Arg Val Ala Lys Glu Gln Ile Ser Arg Arg Leu Asp Gly Asn
 625 630 635 640

30

Gln Tyr Leu Phe Leu Pro Pro Asn Arg Tyr Ile Phe His Gly Ala Glu
 645 650 655

Val Tyr Ser Asp Ser Glu Asp Asp Val Leu Ser Ser Ser Ser Cys Gly
 660 665 670

35

Ser Asn Ser Asp Ser Gly Thr Cys Gln Ser Pro Ser Leu Glu Glu Pro
 675 680 685

40

Met Glu Asp Glu Ser Glu Ile Glu Glu Phe Tyr Asn Gly Leu Glu Asp
 690 695 700

5

Glu Pro Asp Val Pro Glu Arg Ala Gly Gly Ala Gly Phe Gly Thr Asp
 705 710 715 720

10

Gly Asp Asp Gln Glu Ala Ile Asn Glu Ala Ile Ser Val Lys Gln Glu
 725 730 735

15

Val Thr Asp Met Asn Tyr Pro Ser Asn Lys Ser
 740 745

<210> 13

20

<211> 1869

<212> DNA

<213> Homo sapiens

<400> 13

25

ggcgccgggg gggggggtgg gaggcggagg cggggccggg gcgcgcggg cggggcgccg

60

ggggcggggc gaggccggag gactcctcgg actgcgcgga acatggcggt ctgggggttg

120

30

cgcgccgcgg cagccctccg gctgtggggc cgggtagttg aacgggtcga ggccggggga

180

ggcgtggggc cgtttcaggc ctgcggctgt cggctggtgc ttggcggcag ggacgatgtg

35

240

agtgcggggc tgagaggcag ccatggggcc cgcggtgagc ccttggaccc ggcgcgcccc

300

5 ttgcagagggc ctcccagacc cgaggtgccc agggcattcc ggaggcagcc gagggcagca
360

 gctcccagtt tcttcttttc gagtattaaa ggtggaagaa ggtccatata tttttctgtg
420

10 ggtgcttcaa gtgttggttg aagtggaggg agcagtgaca aggggaagct ttccctgcag
480

 gatgtagctg agctgattcg ggccagagcc tgccagaggg tgggtggatcat ggtggggggc
15 540

 ggcatcagca caccagtggt cattccagac ttcagatcgc cggggagtggt cctgtacagc
600

20 aacctccagc agtacgatct cccgtacccc gagggcattt ttgaactccc atttctcttt
660

 cacaacccca agcccttttt cactttggcc aaggagctgt accctggaaa ctacaagccc
720

25 aacgtcactc actactttct ccggctgctt catgacaagg ggctgctttct gcggctctac
780

 acgcagaaca tcgatgggct tgagagagtg tcgggcatcc ctgcctcaaa gctggttgaa
30 840

 gctcatggaa cctttgcctc tgccacctgc acagtctgcc aaagaccctt cccagggggag
900

35 gacattcggg ctgacgtgat ggcagacagg gttccccgct gcccggtctg caccggcggt
960

 gtgaagcccg acattgtggt ctttggggag ccgctgcccc agaggttctt gctgcatgtg
1020

40

5 gttgattttcc ccatggcaga tctgctgctc atccttgga cctccctgga ggtggagcct
1080

 tttgccagct tgaccgaggc cgtgctggagc tcagttcccc gactgctcat caaccgggac
1140

10 ttggtggggc ccttggttg gcctcctgc agcagggacg tggcccagct gggggacgtg
1200

 gttcacggcg tggaaagcct agtggagctt ctgggctgga cagaagagat gcgggacctt
15 1260

 gtgcagcggg aaactgggaa gcttgatgga ccagacaaat aggatgatgg ctgccccac
1320

20 acaataaatg gtaacatagg agacatccac atcccaattc tgacaagacc tcatgctga
1380

 agacagcttg ggcaggtgaa accagaatat gtgaactgag tggacacccg aggtgccac
1440

25 tggaatgtct tctcaggcca tgagctgcag tgactggtag ggctgtgttt acagtcaggg
1500

 ccaccccgtc acatatacaa aggagctgcc tgctgtttg ctgtgttgaa ctcttcactc
30 1560

 tgctgaagct cctaattggaa aaagctttct tctgactgtg accctcttga actgaatcag
1620

35 accaactgga atcccagacc gagtctgctt tctgtgcta gttgaacggc aagctcggca
1680

 tctgttggtt acaagatcca gacttgggcc gagcggctccc cagccctctt catgttccga
1740

40

5 agtgtagtct tgaggccctg gtgccgcact tctagcatgt tggctctcctt tagtggggct
1800

attttttaatg agagaaaatc tgttctttcc agcatgaaat acatttagtc tcctcaaaaa
1860

10 aaaaaaaca
1869

15 <210> 14
<211> 399
<212> PRT
<213> Homo sapiens

20 <400> 14

Met Ala Phe Trp Gly Trp Arg Ala Ala Ala Ala Leu Arg Leu Trp Gly
1 5 10 15

25 Arg Val Val Glu Arg Val Glu Ala Gly Gly Gly Val Gly Pro Phe Gln
20 25 30

30 Ala Cys Gly Cys Arg Leu Val Leu Gly Gly Arg Asp Asp Val Ser Ala
35 40 45

Gly Leu Arg Gly Ser His Gly Ala Arg Gly Glu Pro Leu Asp Pro Ala
35 50 55 60

Arg Pro Leu Gln Arg Pro Pro Arg Pro Glu Val Pro Arg Ala Phe Arg
65 70 75 80

40

5

Arg Gln Pro Arg Ala Ala Ala Pro Ser Phe Phe Phe Ser Ser Ile Lys
 85 90 95

10

Gly Gly Arg Arg Ser Ile Ser Phe Ser Val Gly Ala Ser Ser Val Val
 100 105 110

15

Gly Ser Gly Gly Ser Ser Asp Lys Gly Lys Leu Ser Leu Gln Asp Val
 115 120 125

20

Ala Glu Leu Ile Arg Ala Arg Ala Cys Gln Arg Val Val Val Met Val
 130 135 140

Gly Ala Gly Ile Ser Thr Pro Ser Gly Ile Pro Asp Phe Arg Ser Pro
 145 150 155 160

25

Gly Ser Gly Leu Tyr Ser Asn Leu Gln Gln Tyr Asp Leu Pro Tyr Pro
 165 170 175

30

Glu Ala Ile Phe Glu Leu Pro Phe Phe Phe His Asn Pro Lys Pro Phe
 180 185 190

35

Phe Thr Leu Ala Lys Glu Leu Tyr Pro Gly Asn Tyr Lys Pro Asn Val
 195 200 205

40

Thr His Tyr Phe Leu Arg Leu Leu His Asp Lys Gly Leu Leu Leu Arg
 210 215 220

5

Leu Tyr Thr Gln Asn Ile Asp Gly Leu Glu Arg Val Ser Gly Ile Pro
 225 230 235 240

10

Ala Ser Lys Leu Val Glu Ala His Gly Thr Phe Ala Ser Ala Thr Cys
 245 250 255

15

Thr Val Cys Gln Arg Pro Phe Pro Gly Glu Asp Ile Arg Ala Asp Val
 260 265 270

20

Met Ala Asp Arg Val Pro Arg Cys Pro Val Cys Thr Gly Val Val Lys
 275 280 285

Pro Asp Ile Val Phe Phe Gly Glu Pro Leu Pro Gln Arg Phe Leu Leu
 290 295 300

25

His Val Val Asp Phe Pro Met Ala Asp Leu Leu Leu Ile Leu Gly Thr
 305 310 315 320

30

Ser Leu Glu Val Glu Pro Phe Ala Ser Leu Thr Glu Ala Val Arg Ser
 325 330 335

35

Ser Val Pro Arg Leu Leu Ile Asn Arg Asp Leu Val Gly Pro Leu Ala
 340 345 350

40

Trp His Pro Arg Ser Arg Asp Val Ala Gln Leu Gly Asp Val Val His
 355 360 365

5
 Gly Val Glu Ser Leu Val Glu Leu Leu Gly Trp Thr Glu Glu Met Arg
 370 375 380

10
 Asp Leu Val Gln Arg Glu Thr Gly Lys Leu Asp Gly Pro Asp Lys
 385 390 395

15
 <210> 15
 <211> 1174
 <212> DNA
 <213> Homo sapiens

20
 60
 gtccgtagag ctgtgagaga atgaagatga gctttgogtt gactttcagg tcagcaaaag
 120
 gccgttggat cgcaaaccac agccagccgt gctcgaaagc ctccattggg ttattttgtgc
 25
 180
 cagcaagtcc tcctctggac cctgagaagg tcaaagagtt acagcgcttc atcacccttt
 240
 ccaagagact ccttgtgatg actggggcag gaatctccac cgaatcgggg ataccagact
 30
 300
 acaggtcaga aaaagtgggg ctttatgccc gcaactgaccg caggcccatc cagcatggtg
 360
 attttgtccg gagtgcccca atccgccagc ggtactgggc gagaaacttc gtaggctggc
 420
 ctcaattctc ctcccaccag cctaaccctg cacactgggc tttgagcacc tgggagaaac
 40

5 tcggaaagct gtactggttg gtgacccaaa atgtggatgc tttgcacacc aaggcgggga
480

 gtcgggcgcct gacagagctc cacggatgca tggacagggt cctgtgcttg gattgtgggg
540

10 aacagactcc ccgggggggtg ctgcaagagc gtttccaagt cctgaacccc acctggagtg
600

 ctgaggccca tggcctggct cctgatgggt acgtctttct ctcagaggag caagtccgga
15 660

 gctttcaggt cccaacctgc gttcaatgtg gaggccatct gaaaccagat gtcgtttttct
720

20 tcggggacac agtgaaccct gacaagggtg attttgtgca caagcgtgta aaagaagccg
780

 actccctctt ggtggtggga tcatccttgc aggtatactc tggttacagg tttatcctca
840

25 ctgcctggga gaagaagctc ccgattgcaa tactgaacat tgggccacac cggtcggatg
900

 acttggcgtg tctgaaactg aattctcgtt gtggagagtt gctgcctttg atagacccat
30 960

 gctgaccaca gcctgatatt ccagaacctg gaacagggac tttcacttga atcttgctgc
1020

35 taaagttaaa tgcccttctca aatgacagat tccagttccc attcaacaga gtaggggtgca
1080

 ctgacaaagt atagaagggt ctaggtatct taatgtgtgg atattcttaa ttaaaactca
1140

40

5 ttttttttaa ataaaaaatt gttcagcttt aaaa
1174

<210> 16

10 <211> 314

<212> PRT

<213> Homo sapiens

<400> 16

15

Met Lys Met Ser Phe Ala Leu Thr Phe Arg Ser Ala Lys Gly Arg Trp
1 5 10 15

20

Ile Ala Asn Pro Ser Gln Pro Cys Ser Lys Ala Ser Ile Gly Leu Phe
20 25 30

25

Val Pro Ala Ser Pro Pro Leu Asp Pro Glu Lys Val Lys Glu Leu Gln
35 40 45

30

Arg Phe Ile Thr Leu Ser Lys Arg Leu Leu Val Met Thr Gly Ala Gly
50 55 60

Ile Ser Thr Glu Ser Gly Ile Pro Asp Tyr Arg Ser Glu Lys Val Gly
65 70 75 80

35

Leu Tyr Ala Arg Thr Asp Arg Arg Pro Ile Gln His Gly Asp Phe Val
85 90 95

40

Arg Ser Ala Pro Ile Arg Gln Arg Tyr Trp Ala Arg Asn Phe Val Gly

5		100		105		110											
	Trp	Pro	Gln	Phe	Ser	Ser	His	Gln	Pro	Asn	Pro	Ala	His	Trp	Ala	Leu	
			115					120					125				
10																	
	Ser	Thr	Trp	Glu	Lys	Leu	Gly	Lys	Leu	Tyr	Trp	Leu	Val	Thr	Gln	Asn	
		130					135					140					
15																	
	Val	Asp	Ala	Leu	His	Thr	Lys	Ala	Gly	Ser	Arg	Arg	Leu	Thr	Glu	Leu	
	145					150					155					160	
20																	
	His	Gly	Cys	Met	Asp	Arg	Val	Leu	Cys	Leu	Asp	Cys	Gly	Glu	Gln	Thr	
					165					170					175		
25																	
	Pro	Arg	Gly	Val	Leu	Gln	Glu	Arg	Phe	Gln	Val	Leu	Asn	Pro	Thr	Trp	
				180					185					190			
30																	
	Ser	Ala	Glu	Ala	His	Gly	Leu	Ala	Pro	Asp	Gly	Asp	Val	Phe	Leu	Ser	
			195					200					205				
35																	
	Glu	Glu	Gln	Val	Arg	Ser	Phe	Gln	Val	Pro	Thr	Cys	Val	Gln	Cys	Gly	
		210					215					220					
40																	
	Gly	His	Leu	Lys	Pro	Asp	Val	Val	Phe	Phe	Gly	Asp	Thr	Val	Asn	Pro	
	225				230						235					240	
40																	
	Asp	Lys	Val	Asp	Phe	Val	His	Lys	Arg	Val	Lys	Glu	Ala	Asp	Ser	Leu	

```

5          245          250          255

Leu Val Val Gly Ser Ser Leu Gln Val Tyr Ser Gly Tyr Arg Phe Ile
          260          265          270

10

Leu Thr Ala Trp Glu Lys Lys Leu Pro Ile Ala Ile Leu Asn Ile Gly
          275          280          285

15

Pro Thr Arg Ser Asp Asp Leu Ala Cys Leu Lys Leu Asn Ser Arg Cys
          290          295          300

20

Gly Glu Leu Leu Pro Leu Ile Asp Pro Cys
305          310

25

<210> 17
<211> 1633
<212> DNA
<213> Homo sapiens

30

<400> 17
cgcctctagg agaaagcctg gaacgcgtac cggaggggtac cagagctctt agcggggccgg
60

cagcatgtgc ggggcccaagt aaatggaaat gttttctaac atataaaaac ctacagaaga
120

35

agaaaataat tttctggatc aaattagaag tctgtattat attgatgtct ccagattcaa
180

atatattaga aagcagccgt ggagacaacc atcttcattt tgggagaaat aactaaagcc
40 240

```


5
cgctcaagc attagaacta cagacaaacc ctgatgacac ctctccagat tgtccaagt
300
cgattgattt ccagctata ttgtggcctg aagcctccag cgtccacacg aaaccagatt
10 360
tgctgaaaa tggctcggcc aagttcaagt atggcagatt ttcgaaagt ttttgcaaaa
420
gcaaagcaca tagtcatcat ctgaggagct ggtgttagtg cagaaagtgg tgttccgacc
15 480
ttcagaggag ctggagggtta ttggagaaaa tggcaagccc aggacctggc gactcccctg
540
gcctttgccc acaaccogtc ccgggtgtgg gagttctacc actaccggcg ggagggtcatg
20 600
gggagcaagg agcccaacgc cgggcaccgc gccatagccg agtgtgagac ccggctgggc
25 660
aagcagggcc ggcgagtcgt ggtcatcacc cagaacatcg atgagctgca ccgcaaggct
720
ggcaccaaga accttctgga gatccatggt agcttattta aaactcgatg tacctcttgt
30 780
ggagttgtgg ctgagaatta caagagtcca atttgtccag ctttatcagg aaaagggtgct
840
ccagaacctg gaactcaaga tgccagcatc ccagttgaga aacttccccg gtgtgaagag
35 900
gcaggctgcg ggggcttget ggcacctcac gtcgtgtggt ttggagaaaa cctggatcct
40 960

5
gccattcttg aggaggttga cagagagctc gccactgtg atttatgtct agtgggtgggc
1020
acttcctctg tgggtgtacc agcagccatg ttgcccccc aggtggctgc caggggctg
10 1080
ccagtggctg aatttaacac ggagaccacc ccagctacga acagattcag gtttcatttc
1140
cagggaccct gtggaacgac tcttcctgaa gcccttgctt gtcataaaaa tgaaactgtt
15 1200
tcttaagtgt cctggggaag aaagaaatta cagtatatct aagaactagg ccacacgcag
1260
20
aggagaaatg gtcttatggg tgggtgagctg agtactgaac aatctaaaaa tagcctctga
1320
ttccctcgct ggaatccaac ctgttgataa gtgatggggg tttagaagta gcaaagagca
25 1380
cccacattca aaagtcacag aactggaaag ttaattcata ttatttggtt tgaactgaaa
1440
30
cgtgaggtat ctttgatgtg tatggttggt tattgggagg gaaaaatttt gttaaattaga
1500
ttgtctaaaa aaaatagtta ttctgattat atttttgtta tctgggcaaa gtagaagtca
1560
35
aggggtaaaa accctactat tctgattttt gcacaagttt tagtggaata taaaatcaca
1620
ctctacagta ggt
40 1633

5

<210> 18

<211> 310

<212> PRT

10

<213> Homo sapiens

<400> 18

15

Met Arg Pro Leu Gln Ile Val Pro Ser Arg Leu Ile Ser Gln Leu Tyr
 1 5 10 15

20

Cys Gly Leu Lys Pro Pro Ala Ser Thr Arg Asn Gln Ile Cys Leu Lys
 20 25 30

25

Met Ala Arg Pro Ser Ser Ser Met Ala Asp Phe Arg Lys Phe Phe Ala
 35 40 45

Lys Ala Lys His Ile Val Ile Ile Ser Gly Ala Gly Val Ser Ala Glu
 50 55 60

30

Ser Gly Val Pro Thr Phe Arg Gly Ala Gly Gly Tyr Trp Arg Lys Trp
 65 70 75 80

35

Gln Ala Gln Asp Leu Ala Thr Pro Leu Ala Phe Ala His Asn Pro Ser
 85 90 95

40

Arg Val Trp Glu Phe Tyr His Tyr Arg Arg Glu Val Met Gly Ser Lys
 100 105 110

5

Glu Pro Asn Ala Gly His Arg Ala Ile Ala Glu Cys Glu Thr Arg Leu
 115 120 125

10

Gly Lys Gln Gly Arg Arg Val Val Val Ile Thr Gln Asn Ile Asp Glu
 130 135 140

15

Leu His Arg Lys Ala Gly Thr Lys Asn Leu Leu Glu Ile His Gly Ser
 145 150 155 160

20

Leu Phe Lys Thr Arg Cys Thr Ser Cys Gly Val Val Ala Glu Asn Tyr
 165 170 175

Lys Ser Pro Ile Cys Pro Ala Leu Ser Gly Lys Gly Ala Pro Glu Pro
 180 185 190

25

Gly Thr Gln Asp Ala Ser Ile Pro Val Glu Lys Leu Pro Arg Cys Glu
 195 200 205

30

Glu Ala Gly Cys Gly Gly Leu Leu Arg Pro His Val Val Trp Phe Gly
 210 215 220

35

Glu Asn Leu Asp Pro Ala Ile Leu Glu Glu Val Asp Arg Glu Leu Ala
 225 230 235 240

His Cys Asp Leu Cys Leu Val Val Gly Thr Ser Ser Val Val Tyr Pro
 245 250 255

40

5

Ala Ala Met Phe Ala Pro Gln Val Ala Ala Arg Gly Val Pro Val Ala
260 265 270

10

Glu Phe Asn Thr Glu Thr Thr Pro Ala Thr Asn Arg Phe Arg Phe His
275 280 285

15

Phe Gln Gly Pro Cys Gly Thr Thr Leu Pro Glu Ala Leu Ala Cys His
290 295 300

20

Glu Asn Glu Thr Val Ser
305 310